

## Determination of *ctxAB* Expression in *Vibrio cholerae* Classical and El Tor Strains using Real-Time PCR

Seyed Mahmoud Amin Marashi<sup>1,2</sup>, Ramazan Rajabnia<sup>3</sup>, Abbas Ali Imani Fooladi<sup>4</sup>, Zohreh Hojati<sup>5</sup>,  
Sharareh Moghim<sup>6</sup>, Bahram Nasr Esfahani<sup>6\*</sup>

1. Cellular and Molecular Biology Research Center, (CMBRC), Babol University of Medical Sciences, Babol, Iran.

2. Department of Microbiology and Immunology, Babol University of Medical Sciences, Babol, Iran.

3. Infectious Diseases & Tropical Medicine Research Center, Babol University of Medical Sciences, Babol, Iran.

4. Applied Microbiology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.

5. Genetics Division, Biology Department, Faculty of Sciences, University of Isfahan, Isfahan, Iran.

6. Department of Microbiology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.

Submitted 5 March 2013; Accepted 18 March 2013


Cholera is an infection of the small intestines caused by the bacterium *V. cholerae*. It is a major cause of health threat and also a major cause of death worldwide and especially in developing countries. The major virulence factor produced by *V. cholerae* during infection is the cholera toxin. Total mRNA extraction and reverse transcription was performed for making *ctxAB* cDNA. Relative Real-Time PCR analysis showed unequal enterotoxin production in *V. cholerae* strains. The results showed that, classical strain produces cholera toxin more than El Tor strain.

**Key words:** *Vibrio cholerae*, RT-qPCR, *ctxAB* expression

Cholera is one of the infectious diseases that still happens in developing countries. The 8th pandemic of cholera spreads from Southeast Asia across the Middle East and into Central America and Africa (1, 2). The important pathogenesis factor in *Vibrio cholerae* is a potent enterotoxin, cholera toxin, which causes the severe diarrhea of cholera (3, 4). The cholera toxin is produced by *V. cholerae* and CTXΦ phage corporation. Control of enterotoxin gene expression seems to be complex, so that environmental factors are very important in

its expression (5, 6). The environmental signals affect *TcpPH* gene and cause its activation and finally affect *ToxT* gene. The ToxT protein is the most important agent for *ctxAB* toxin expression, because ToxT protein attaches to *toxbox* region at upstream of *ctxAB* gene and induces *ctxAB* toxin expression (7, 8) (Fig 1). Beside, other signaling systems such as *ToxR*, *RS1*, *AphAB* and quorum sensing have positive or negative effects on ToxT protein (9, 10). Moreover, H-NS protein has negative effect on *TcpPH* and *ToxT* genes that

\* Corresponding author: Department of Microbiology, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.  
E-Mail: nasr@hlth.mui.ac.ir.



**Fig 1.** Diagram of the *vibrio cholerae* ToxR regulon and *ctxAB* expression, with permission from ASM

finally would decrease the *ctxAB* toxin production (11). The RS<sub>1</sub> region contains *rstA*, *rstB* and *rstC* fragments which have positive effect on *ToxR* gene and therefore increase *ctxAB* toxin production (12). Bakhshi *et al.* reported several *V. cholerae* attacks in Southwest and Southeast of Iran between 2005-2009 (13, 14). As the level of production of a protein is somehow related to its mRNA quantity, we therefore aimed to determine *V. cholerae* strains that can produce more *ctxAB* toxin.

## Material and Method

### Bacterial strains and growth conditions

We used two standard strains named *V. cholerae* O<sub>1</sub> Classic ATCC 14035 & *V. cholerae* O<sub>1</sub> El Tor N16961. The isolates were confirmed by biochemical and immunological tests. Serotyping was performed using monoclonal O<sub>1</sub> antiserum and mono-specific Inaba and Ogawa antisera (Pasteur institute, Paris, France). All selected strains were cultured according to the AKI-SW method and standard growth curve were drawn (15).

### Isolation of RNA and RT-PCR

Approximately,  $2 \times 10^8$  cfu/ml from each strain, was used for total RNA extraction. Total RNA was isolated from the strains isolated randomly from each *V. cholerae* grown in AKI

medium using the RNeasy® Protect Bacteria Mini Kit (Qiagen Inc, GMBH, Germany) and the integrity and purity was checked. Equivalent concentrations of total RNA from each strain were selected as template for RT-PCR. cDNA synthesis and PCR amplification were performed using QuantiTec Reverse Transcription Qiagen kit (Qiagen Inc, GMBH, Germany). RT-PCR was performed in the presence of random primer at 42°C for 10 min. After cDNA synthesis, the *ctxAB* and *recA* genes were PCR amplified for checking. PCR amplification was performed for 35 cycles as follows: initial denaturation at 94°C for 5 min, then denaturation at 94°C for 30 sec, annealing at 60°C for 30 sec, extension at 72°C for 30 sec. At the end of the 35th cycle, reaction mixtures were left at 72°C for another 3 min. Five microliters of each reaction mixture was loaded on a 1% agarose gel and subjected to electrophoresis to confirm that the unique amplified fragment correspond to the expected *ctxAB* gene fragment and *recA* as housekeeping gene (16).

### Real-Time PCR

Prepared cDNA was quantified using SYBR green I dye. Four primers were designed by AlleleID 6 software, 5'-CAGTCAGGTGGTCT-TATGC-3' (*ctxAB*-F) and 5'-ATCGTGCCTAAC-

### Determination of *ctxAB* Expression in *Vibrio cholerae*

AAATCCC-3' (*ctxAB*-R) for gene of interest and 5' -ATTGAAGGCGAAATGGGCGATAG- 3' (*recA*-F) and 5' -TACACATACAGTTGGATTGCTTG AGG- 3' (*recA*-R) for housekeeping gene. Those primers were specific to *ctxAB* and *recA* and amplified a 115 & 106 bp respectively. SYBR green Real-Time PCR assay was performed with a 20  $\mu$ l PCR mixture volume containing 2x QuantiTec SYBR Green PCR Master Mix (Qiagen Inc, GMBH, Germany), 0.25  $\mu$ M specific primer sets, and 2  $\mu$ l of cDNA sample. Amplification of the primers, data acquisition, and relative analysis were carried out in Chromo4 BioRad Real-Time PCR. PCR reactions were performed as followings: one cycle of 95  $^{\circ}$ C for 5 min, then 40 cycles of 95  $^{\circ}$ C for 15 sec, 60  $^{\circ}$ C for 30 sec. Following the amplification, melting temperature analysis of the PCR products was performed to determine the specificity of the PCR. The standard curve was established by using genomic DNA for each studied gene to confirm that the primers amplified at the same rate and to validate the experiment (55-95 $^{\circ}$ C with warming of 0.2 $^{\circ}$ C per sec). Reverse transcription and PCR positive controls (RNA and DNA, respectively) and negative controls (distilled water) were included in each run. The Real-Time PCR reaction was performed twice assayed in


triplicate. Classical *V. cholerae* O<sub>1</sub> ATCC 14035 was used as a standard control.

### Results


The specificity of each primer set for *V. cholerae* was tested by PCR with genomic DNA extracted via boiling. Only one size of amplicon was obtained by PCR reaction for *ctxAB* and *recA* genes when DNA from *V. cholerae* strains was used. The amplicons obtained for each gene were verified by sequencing. The presence of a single PCR product was confirmed by Real-Time PCR for each set of the primers using melting curve analysis that resulted in a single product-specific melting curve (Fig 2). The PCR efficiencies varied between 1.90 and 1.94. The relative expression ratio was calculated for each gene of interest by a mathematical model described by  $\Delta C_T$  method. The Cycle Threshold ( $C_T$ ) results are showed in table 1. Histogram and samples  $C_T$  values are indicated in Fig 3.

### Discussion

In our study, the results are derived by using "relative" method and  $\Delta C_T$  formula. By considering that *ctxAB* primers have been carefully designed, the amount of standard deviation results are close to



**Fig 2.** Melting curves of *ctxAB* and *recA* genes.



**Fig 3.** Level of cholera toxin production in classical and EL Tor strains.

Table 1: Cycle threshold ( $C_t$ ) results for the *V.cholerae* O<sub>1</sub> Classic ATCC 14035 & *V. cholerae* O<sub>1</sub> El Tor 62013

Strain	Mean $C_t$ recA $\pm$ SD	Mean $C_t$ ctxAB $\pm$ SD	Mean $\Delta\Delta C_t$	Mean Ratio*
Classic	25.58 $\pm$ 0.14	24.35 $\pm$ 0.39	1.23	1.0
El Tor	22.85 $\pm$ 0.13	26.02 $\pm$ 0.15	4.35	0.06

\* $\Delta\Delta C_t$  was calculated as:  $\Delta C_t$  (test) -  $\Delta C_t$  (calibrator). Ratio=efficiency<sup>- $\Delta\Delta C_t$</sup> .

zero, the primer-dimer bands were not seen, because the concentration of participating primers in the reaction had been set up. Also, for more accuracy and sensitivity, the PCR efficiency in Real-Time PCR reactions were calculated and replaced with ratio 2 in computational equations.

Dirita *et al.* showed that the level of cholera toxin was higher in the classical strain compared to El Tor strain because of the influence of *toxR* on cholera toxin production in the classical strain (17). In our study, in addition to Dirita *et al.* results, we determined quantitative cholera toxin production between classic and El Tor strains. Both classical and El Tor strains have been shown to express equivalent levels of ToxR. In contrast, the classical strain expresses more ToxT, which has a higher binding affinity to toxbox region, resulting in higher expression of cholera toxin (18).

The comparison of the  $C_T$  of the El Tor strain with classical strain shows that toxin production in El Tor strain is approximately 16-17 times lower than in the classical strain ( $P_{value}<0.05$  for each). This result is consistent with other reports because the amount of pathogenicity in classic strain is more than El Tor strains (18). Furthermore, histogram and samples  $C_T$  results indicated that toxin production in classical strain is higher than El Tor strain (Fig 3). In conclusion, the results of our study suggest that other factors modulate the production of cholera toxin by regulating the CTX cassette, supporting the idea that cholera toxin production in *V. cholerae* classical and El Tor

strains is a multi-factorial phenomenon.

### Acknowledgment

We are grateful to Dr. Bitu Bakhshi for shipment of the *V. cholerae* O<sub>1</sub> classic ATCC 14035 and *V. cholerae* O<sub>1</sub> El Tor ATCC N16961 strains.

### References

1. Amin Marashi SM, Nasr Esfahani B, Tavakoli A, et al. Simultaneous detection of integrase and antibiotic resistance genes within SXT Constin in *Vibrio cholerae* O1 El Tor strains isolated from Iran using multiplex-PCR. Iran J Basic Med Sci 2012;15:885-9.
2. Taneja N, Mishra A, Sangar G, et al. Outbreaks caused by new variants of *Vibrio cholerae* O1 El Tor, India. Emerg Infect Dis 2009;15:352-4.
3. Baptista MA, Andrade JR, Vicente AC, et al. The Amazonia variant of *Vibrio cholerae*: molecular identification and study of virulence genes. Mem Inst Oswaldo Cruz 1998;93:601-7.
4. Bravo L, Monte RJ, Ramirez M, et al. Detection of toxigenic *Vibrio cholerae* O1 using polymerase chain reaction. Mem Inst Oswaldo Cruz 1992;87:443-4.
5. Bakhshi B, Pourshafie MR, Navabakbar F, et al. Genomic organisation of the CTX element among toxigenic *Vibrio cholerae* isolates. Clin Microbiol Infect 2008;14:562-8.
6. Sa LL, Fonseca EL, Pellegrini M, et al. Occurrence and composition of class 1 and class 2 integrons in clinical and environmental O1 and non-O1/non-O139 *Vibrio cholerae* strains from the Brazilian Amazon. Mem Inst Oswaldo Cruz; 105:229-32.
7. Withey JH, Dirita VJ. *Vibrio cholerae* ToxT independently activates the divergently transcribed *aldA* and *tagA* genes. J

### Determination of *ctxAB* Expression in *Vibrio cholerae*

Bacteriol 2005;187:7890-900.

8. Withey JH, DiRita VJ. The toxbox: specific DNA sequence requirements for activation of *Vibrio cholerae* virulence genes by ToxT. Mol

Microbiol 2006;59:1779-89.

9. Klose KE. Regulation of virulence in *Vibrio cholerae*. Int J Med Microbiol 2001;291:81-8.

10. Zhu J, Miller MB, Vance RE, et al. Quorum-sensing regulators control virulence gene expression in *Vibrio cholerae*. Proc Natl Acad Sci USA 2002;99:3129-34.

11. Matson JS, Withey JH, DiRita VJ. Regulatory networks controlling *Vibrio cholerae* virulence gene expression. Infect Immun 2007;75:5542-9.

12. Davis BM, Waldor MK. Filamentous phages linked to virulence of *Vibrio cholerae*. Curr Opin Microbiol 2003;6:35-42.

13. Bakhshi B, Pourshafie MR. Assessing clonality of *Vibrio cholerae* strains isolated during four consecutive years (2004 - 2007) in Iran. Scand J Infect Dis 2009;41:256-62.

14. Pourshafie MR, Bakhshi B, Ranjbar R, et al. Dissemination of a single *Vibrio cholerae* clone in cholera outbreaks during 2005 in Iran. J Med Microbiol 2007;56:1615-9.

15. Iwanaga M, Yamamoto K, Higa N, et al. Culture conditions for stimulating cholera toxin production by *Vibrio cholerae* O1 El Tor. Microbiol Immunol 1986;30:1075-83.

16. Amin Marashi M, Bakhshi B, Imani Fooladi AA, et al. Quantitative expression of cholera toxin mRNA in *Vibrio cholerae* isolates with different CTX cassette arrangements. J Med Microbiol.

17. DiRita VJ, Neely M, Taylor RK, et al. Differential expression of the ToxR regulon in classical and El Tor biotypes of *Vibrio cholerae* is due to biotype-specific control over toxT expression. Proc Natl Acad Sci U S A 1996;93:7991-5.

18. Gonzalez-Bonilla C, Gutierrez-Cogco L, Moguel-Pech L, et al. [Evaluation of the ELISA method for cholera toxin determination in *Vibrio cholerae* cultures]. Rev Latinoam Microbiol 1994;36:273-6.