

Heterogeneity of the Level of Activity of Lgr5+ Intestinal Stem Cells

Shirin Moossavi*

Digestive Oncology Research Center, Digestive Disease Research Institute, Tehran University of Medical Sciences, Tehran, Iran.

Submitted 4 August 2014; Accepted 15 September 2014; Published 22 September 2014

Intestinal stem cells (ISCs) are a group of rare cells located in the intestinal crypts which are responsible for the maintenance of the intestinal epithelial homeostasis and regeneration following injury or inflammation. Lineage tracing experiments in mice have proven that ISCs can repopulate the entire intestinal crypt. It is noteworthy that in such experiments, only a subset of intestinal crypts is marked by the specific marker. This is suggestive of different levels of activity of stem cells in different crypts i.e. intracryptal variation. Niche succession i.e. dominating the entire crypt by the progenies of one stem cell is also suggestive of the intercryptal stem cell heterogeneity. Regional differences in crypt size, proliferative index, and distribution of proliferative cells along the crypt axis have been reported. It is conceivable that ISCs are heterogeneous in terms of their levels of activity. Appreciation of such heterogeneity will significantly challenge the way in which ISCs are investigated. A better understanding of ISC biology will in turn improve our mechanistic understanding of major intestinal disease including inflammatory bowel disease and colorectal cancer.

Key words: Intestinal stem cells, heterogeneity, dormancy

The intestinal epithelial layer is organized into distinct anatomical features called villi which are protrusions in the lumen, and crypts of Lieberkühn which are invaginations of the epithelial layer into the lamina propria. Villi are absent in the colon. Intestinal stem cells (ISCs) are a group of rare cells located in the intestinal crypts which are responsible for the maintenance of the intestinal epithelial homeostasis and regeneration following injury. Stem cells are quiescent cells with infrequent proliferation; the result of which is usually a new stem cell and a transient-amplifying

cell (TAC). TACs differentiate into intestinal epithelial subtypes, i.e. enterocytes, goblet cells, neuroendocrine cells, and Paneth cells; which migrate upward along the crypt axis. Paneth cells are an exception to the migration pattern of mature epithelial cells as they reside at the base of the crypts. It is noteworthy that Paneth cells do not exist in the colon under physiologic conditions but appear in the chronically inflamed colon. ISCs are believed to reside in a functionally defined 'niche' at the base of the crypt. The stem cell niche maintains the stem cell properties and regulates the

* Corresponding author: Digestive Oncology Research Center, Digestive Disease Research Institute, Tehran University of Medical Sciences - Shariati Hospital, North Amirabad Ave., Tehran 14117, Iran. Email: shirin.moossavi@gmail.com

Intestinal Stem Cell Heterogeneity

quiescence /proliferation of the stem cell via stem cell regulatory pathways namely Wnt, Notch, BMP and Hedgehog pathways (1). ISCs are defined by their self-renewal capacity and multipotency. Multiple stem cells exist at the base of each crypt. However, only a single stem cell dominates the niche (niche succession); followed by cellular domination of the crypt (monoclonal conversion) (2, 3). This is suggestive of a level of heterogeneity in ISC activity. Our understanding of the ISCs has extensively expanded over the past few decades. It is already known that two types of fast-and slow-cycling stem cells reside in the lower crypts; which are different in terms of cell cycle activity (4). However, it is less clear whether the members of each group also have distinctive characteristics. Hereby, the evidence for the heterogeneity of Lgr5+ ISCs is summarized.

Intestinal stem cells

In the murine small intestine, two positions are suggested for the stem cells: at the base of the crypt and at the position 4 from the base of the crypt. The crypt base columnar (CBC) cells reside at the base of the crypt interspersed by the Paneth cells. They are positive for Ki67 proliferation marker and uptake BrdU as early as 6 hours post injection. Their cycling rate is estimated to be once every 24 hours. The CBC cells are Lgr 5 positive and are rapid cycling (5). The position 4 cells, on the other hand, are thought to be quiescent and slow cycling. They are capable of label-retention and are located at position 2-7 immediately above the Paneth cells. Slow cycling cells have been historically identified by their ability to retain BrdU which is diluted in the cycling cells of the crypt. Recently, they have been found to express Bmi1 and mTert (6, 7). Position 4 stem cells do not exist in the colon and their role as ISCs are controversial. For example, label-retaining position 4 cells have been observed to be secretory cell precursors (8). On the other hand, markers of the position 4 cells namely Bmi1, Tert, Lrig1, and Hopx have been found to be either

expressed in the Lgr5+ cells or uniformly along the crypt axis (9, 10). It has recently been demonstrated that Lgr5 and Bmi1 mark non- overlapping stem cell populations in mice. CBCs and position 4 stem cells seem to demonstrate distinct functional properties. In a systematic comparison of the two cell population in the murine small intestine, it was found that under steady state, Bmi1+ cells are more quiescent and less efficiently contribute to the epithelial turnover compared to Lgr5+ ISCs. In addition, modulation of the Wnt pathway by intravenous adenoviral delivery of either Wnt agonist, R-Spondin1 (Rspo1), or antagonist, Dickkopf-1 (Dkk1), does not affect Bmi1+ cells whilst Lgr5+ cells are highly sensitive to the level of Wnt pathway activity. Lgr5+ cells are completely lost after 12 Gy whole body γ -irradiation and reappear at a very low frequency 4.5 to 7 days post-irradiation. In contrast, Bmi1+ cells are radioresistant and do not quantitatively change. However, they demonstrate a significant increase in the proliferation rate and niche succession post-irradiation. It is suggested that Lgr5+ cells are the active ISCs under homeostatic condition whereas Bmi1+ cells are responsible for epithelial regeneration upon injury (4). In an attempt to investigate the consequence of loss of Lgr5+ CBCs, Tian et al. knocked in diphtheria toxin receptor in the Lgr5 locus. Upon long-term administration of the toxin, CBCs are ablated. However, the crypt anatomy is preserved. CBC-deficient crypts are also capable of generating organoids *in vitro* with comparable efficiency to the control. Upon removal of the toxin from the medium, Lgr5 expressing cells reappear in the organoids. In other words, the regenerational capacity of the crypt is preserved in spite of loss of Lgr5+ ISCs. Hence, Bmi1+ cells are suggested to be the quiescent ISCs which function upon injury. It was observed that Bmi1+ cells are expanded in Lgr5+ cells-depleted crypts in the proximal small intestine. It is concluded that Bmi1+ cells function as reserve stem cells upon

damage or loss of more rapidly cycling Lgr5⁺ cells (11).

Intestinal stem cell heterogeneity

Stem cell heterogeneity has been described for embryonic (12-16), muscle (17), hematopoietic (18, 19), neural (20), and induced pluripotent stem cells (21). Stem cell heterogeneity at the level of dormancy has been described in murine hair follicle. The hair follicle undergoes cycling phases of destruction, rest and rapid proliferation which are regulated by major signalling pathways including Wnt, TGF- β , and BMP. Janich et al. demonstrated that the bulge CD34⁺/ α 6 integrin^{high} stem cells have distinct levels of clock pathway activity at each stage of the hair cycling. Different state of the clock activity was further shown to affect the stem cell regulatory pathways such as Wnt and TGF- β at gene expression level (22); thus, conferring the stem cells with distinct levels of activity in terms of readiness to self-renew and fate decision.

Intestinal crypts are clonal units which are maintained by the continual division of the crypt stem cells. Before genetic manipulation and transgenic mice were available, ISCs were studied by analysis of somatic mutations. The rationale was that if the mutation occurs in the stem cell, the mutation will be fixed and the whole crypt will be composed of the mutated epithelium. The time which takes for the whole mutant crypt to appear i.e. the clonal stabilization time, has been the subject of studies in mouse and human intestine. Interestingly, differences have been observed in the cell kinetics of small vs. large intestine in mice. Campbell et al. investigated the mutation fixation in colectomy samples which had received radiotherapy prior to surgery. The proportion of somatic mutation fixation rose significantly one month after irradiation and reached the peak at 4-12 months. Subsequently, the proportion of partially mutated crypts decreased significantly with time i.e. at 4-12 months the majority of mutated stem cells have dominated the niche. Once the wholly mutated

crypts appear, they persist in the colon for considerable length of time. This is suggestive of the total replacement of the stem cells by one or more ancestral mutated stem cell. Interestingly, the partially mutated crypts are not persistent. They either switch to wholly mutated crypts if the mutation is in the stem cell, or they will become normal again if the mutation is in the TACs (23). This is in accordance with a recent finding that heterozygous APC mutation will initiate intestinal tumourigenesis if it is induced in the stem cells and not TACs (24).

One caveat of mutation analysis is that it might not reflect the normal behaviour of ISCs (25). Yatabe et al. have examined methylation status of CpG islands of myogenic factor 3 (MYOD1), cardiac-specific homeobox (CSX) and an X-chromosome CpG-rich region in biglycan (BGN) to assess the epigenetic distance between crypts and indirectly study the stem cell dynamic in human colon. The results are presented in a binary system of 0 and 1; where 0 represents an unmethylated state whereas 1 is a methylated state. Epigenetic distance was defined as the absolute number of differences in methylation status of the studied genes; maximum of which was 5, 8, and 9 for MYOD1, CSX and BGN, respectively. The intracryptal and intercryptal epigenetic distances were calculated as the average epigenetic distance of all possible pairs of molecules within a crypt and between crypts, respectively. Applying genetic phylogenetic analysis, they argue that if there are multiple immortal stem cells in a crypt, the crypt will be polyclonal with each clone having a separate last common ancestor. However, if a stem cell dominates the niche, eventually the crypt will become monoclonal; which in phylogenetic terms equate to the origination of all cells from one last common ancestor. Interestingly, multiple unique epigenetic tags were identified in each crypt which suggests that crypts are not monoclonal (26).

Intestinal Stem Cell Heterogeneity

Mathematical modelling and computational biology provide advantageous tools to assess biologic hypotheses which are not amenable to experimentation. Intestinal crypt homeostasis and intestinal tumorigenesis have been the topic of several mathematical models. Of relevance to the question of ISC heterogeneity, Lobachevsky and Radford have constructed a model based on the assumption that two tiers of stem cells exist in the intestinal crypt with distinct levels of activity and cycling patterns (27). This is in accordance with the experimental data in small intestine where CBC and position 4 cells appear to have non-overlapping functionality (4). However, experimental approach thus far has failed to produce the evidence for the existence of the slow-cycling stem cell in the colon. It is assumed that the cycling pattern of each stem cell type is the average of all its members. Therefore, it is conceivable that the stem cell pool is consistent of stem cells with distinct cycling rates. Hormoz suggests that stem cell heterogeneity in terms of the cell cycle rate results in an overall reduced rate of replicative aging in the tissue which is of importance in a rapidly proliferating tissue such as the intestine. It is further proposed that one single stem cell cycling rate is influenced by the microenvironmental signals. Nevertheless, the overall average is similar to that of the stem cell population (28). Monoclonal conversion of the intestinal crypt has been experimentally demonstrated in mouse duodenum. It was argued that niche succession occurs as a result of a neutral competition among stem cells which were assumed to be functionally homogeneous (29). Nevertheless, the possibility that some stem cells might outcompete their counterparts implies that the stem cell pool is conceivably heterogeneous, at least transiently.

The hurdles of directly investigating ISC behavior were overcome by the availability of transgenic mice. Lineage-tracing experiments in mice have proven that a subset of cells highly

expressing Lgr5, CD133, Bmi1, or mTert can repopulate the entire intestinal crypt (5-7, 30). It is noteworthy that in such experiments, only a subset of intestinal crypts is marked by the specific stem cell marker. In addition, distinct marking patterns have been observed in a lineage-tracing experiment where some crypts are marked continuously whereas others are either discontinuously marked or only a few marked cells form short v-like formations within a crypt (31). This is suggestive of different levels of activity of stem cells in different crypts i.e. intracryptal variation. Indeed, Lgr5⁺ cells are found to be distinct in terms of proliferation activity as evidenced by the different levels of Ki67 expression. Based on the expression of Lgr5 and Ki67 four groups of cells were identified: Lgr5^{high}Ki67⁺ dividing ISCs, Lgr5^{high}Ki67⁻ quiescent ISCs, Lgr5^{low}Ki67⁺ proliferating progenitors, and Lgr5^{low}Ki67⁻ quiescent progenitors (32). Niche succession i.e. dominating the entire crypt by the progenies of one stem cell is also suggestive of the intercryptal stem cell heterogeneity. Single cell imaging of hair follicle stem cell compartment revealed that stem cells have distinct behaviour depending on their position within the 'niche' in the hair follicle (33). It was previously shown that the proximity to Paneth cells determines the stemness of ISCs (34). Ritsma et al. have investigated spatial heterogeneity of Lgr5⁺ ISCs at single cell level using multiphoton intravital microscopy through an abdominal imaging window. It was observed that stem cells rearrange following a division event which could eventually lead to removal of the border ISCs from the 'niche'. In addition, it was observed that the position of the ISC within the crypt was a determining factor of its self-renewal capacity, i.e. cells located at positions 0 to+2 were more likely to retain their stemness whilst cells at positions+3 and+4 were more likely to divide into TACs (35).

Heterogeneity of ISCs can also be considered along the intestinal tract. Regional differences in

crypt size, proliferative index and the distribution of proliferative cells along the crypt axis have been reported. For example, the crypt length is found to be significantly shorter in the ascending colon compared to transverse and descending colon. Moreover, the total number of proliferative cells is significantly lower in ascending colon compared to other colon segments. Other suggested variations include the cell cycle time, duration of mitosis, and apoptotic index post-irradiation (36, 37). Furthermore, the total number of Lgr5⁺ stem cells significantly varies in different segments of the small intestine as well as between small intestine and colon in mouse and human. There is also a regional difference in the gene expression level of stem cell related genes and Wnt pathway modulators along the intestinal tract (38). In addition, not all crypts contains a non-Paneth label- retaining cell with only 10% having one and 0.1% having two such cells in the small intestine (39). Middendorp et al. have recently demonstrated that ISCs from different regions of the intestinal tract have distinct gene expression profile which is sustained following long *in vitro* organoid culture (40). Overall, it seems that stem cells vary along the intestinal tract.

Although, niche succession is a commonly accepted phenomenon in the intestinal crypt, the underlying mechanism is less well understood. The clock pathway is found to be required for ISC proliferative response following injury in *Drosophila*. Interestingly, clock-intact ISCs are found at different stages of the cell cycle, whereas ISCs are homogeneously either at G0 or S/ G2/ M phase irrespective of the time of the experiment upon RNAi clock knockdown (41). It is suggestive that the clock pathway maintains a degree of heterogeneity in the level of activity of ISCs. It was further shown that clock pathway regulates Wnt pathway components at gene expression level in the skin (22). This is not known in the intestinal epithelium and ISCs. Besides, the ISC hetero-

geneity in terms of the clock pathway activity has not been investigated. However, the clock pathway is required for intestinal homeostasis (42). Although, the clock pathway has been shown to be functional in ISCs in *Drosophila* and was required for an appropriate proliferative response following dextran sodium sulphate treatment; it was not shown whether stem cells differ in the level of clock pathway activity (41). Other potential mechanisms which contribute to ISC heterogeneity need to be explored.

Cancer stem cell heterogeneity

It is long known that tumor tissue is composed of heterogeneous cell populations; a small fraction of which, i.e. cancer stem cells (CSC) have the capacity of self-renewal and sustaining the tumor evolution. Colorectal CSC are traditionally characterized by their ability to reproduce the original tumor hierarchy in mouse transplant experiments (43-45). However, the role of the microenvironment and an induced wound healing process are confounding factors in such experiments (46, 47). This issue is overcome in lineage tracing experiment. Murine experiments have clearly shown that tumorigenic hits can only result in an intestinal tumor if they happen in the stem cell compartment and not the transient-amplifying region (48). In a recent paper, Schepers et al. have demonstrated that APC-driven adenoma originates either from a single or multiple APC-mutated Lgr5⁺ cells in mice (24). Moreover, constitutive activation of NF-κB and Wnt in intestinal epithelial cells results in the formation of aberrant crypt- like structures in the villi with high proliferation rate and high level of expression of stem cell markers. Interestingly, in the presence of constitutive NF-κB activity, a Wnt-activating mutation in the Lgr5 cell is sufficient to induce 'stemness' and initiate adenoma formation (49). If an ISC is affected by an oncogenic mutation, it could potentially result in the fixation of the mutation in a patch of cells which are histologically

Intestinal Stem Cell Heterogeneity

normal (field characterisation); providing that it retains its self-renewal capacity (50). Interestingly, ISC heterogeneity can be the result of an oncogenic mutation in the course of CRC tumorigenesis. For example, mosaic activated oncogenic K-ras^{G12D} in Lgr5⁺ISCs, results in growth advantage of mutated ISCs as evidenced by niche succession and clonal fixation by mutated ISCs (51).

It has recently been suggested that the CSC population is also heterogeneous with some cells more readily capable of tumor initiation and propagation than others (52). The CSCs heterogeneity has been demonstrated for breast and colorectal cancers (53, 54). Microenvironmental cues are suggested to fine-tune Wnt pathway activity in the course of intestinal tumorigenesis which in turn, can affect cancer stemness behavior of tumor cells. Despite the loss of APC in CRC, nuclear localization of β -catenin is not homogeneously observed in all tumor cells (55). Heterogeneous nuclear β -catenin staining is also observed in colonospheres. Gene expression comparison of colonosphere-derived cells with high vs. low Wnt pathway activity revealed that stem cell markers Lgr5 and Ascl2 are enriched in Wnt^{high} population whereas differentiation markers Muc2 and Krt20 are abundant in Wnt^{low} cells. Cells with high level of Wnt activity demonstrate higher level of clonogenicity and stemness features. Wnt^{high} colonosphere-derived cells are more efficient in reproducing the original tumor in immunocompromised mice (56). Colorectal cancer (CRC)-derived colonospheres were studied at single cell level. Each colonosphere was composed of cells with different proliferative capacity including postmitotic cells; cells which could undergo 1-3 divisions; and a fraction capable of self-renewal and sphere formation. The percentage of sphere forming cells varied in three tumor cells derived from primary tumors from different patients. Interestingly, the percentage of the sphere-forming cells remained stable in serial replating

experiments for each patient. Using a cell tracking method in xenotransplantation experiment confirmed that tumors were polyclonal. Further investigation revealed that three populations of cells reproduced the tumor in mice: the so-called long-term tumor-initiating cell and tumor transient-amplifying cells; which were present in the primary tumour as well as following xenotransplants; and delayed-contributing tumor-initiating cells; which did not contribute to the first tumour formation in mice but appeared in the following rounds of serial transplantation (54). It is concluded that CRC-derived tumour-initiating cells are functionally heterogeneous.

Both normal intestinal and CRC cells demonstrate plasticity i.e. differentiated cells can dedifferentiate into cells with 'stemness' potential. Cellular heterogeneity within a tumor poses challenges regarding diagnosis, subtyping and treatment. Moreover, heterogeneity within CSCs adds another layer of complexity. However, unravelling genetic and microenvironmental cues which regulate the level of 'stemness' of cells could conceivably be directly targeted in a therapeutic setting (57, 58).

Conclusion and Perspective

Lgr5⁺ISCs are often considered as collective homogeneous population of identical cells. However, different spatial positioning of the stem cells within a crypt plausibly results in a small yet effective difference in the level of the availability of stem cell regulatory molecules. This will in turn affect the transcriptomic landscape of each stem cell; which determines the stem cell behavior in terms of readiness to enter cell cycle as well as the choice between symmetric vs. asymmetric division. The fact that progenies of one stem cell dominate the crypt, suggests that some stem cells can outcompete others. In other words, stem cells are distinct in their capacity to populate the crypt. It is conceivable that a similar phenomenon to niche succession also exists in the course of CRC

tumorigenesis where some CSCs are more active than others. The existence of colorectal CSCs with distinct tumor-initiating potential is supporting evidence to this hypothesis. CSC hypothesis proposes that cancer originates from a last common ancestral cell which is most likely the normal tissue stem cell. However, stem cell heterogeneity means that only a subset of stem cells can potentially be the cell-of-origin of the tumour. Therefore, it is crucial to discover the factors which underlie ISC heterogeneity to better understand the CRC tumorigenesis. Understanding the stem cell heterogeneity could then be a target for more selective cancer treatment.

Conflict of interests

The author declared no conflict of interests.

References

1. Medema JP, Vermeulen L. Microenvironmental regulation of stem cells in intestinal homeostasis and cancer. *Nature* 2011;474:318-26.
2. van der Flier LG, Clevers H. Stem cells, self-renewal, and differentiation in the intestinal epithelium. *Annu Rev Physiol* 2009;71:241-60.
3. Umar S. Intestinal stem cells. *Curr Gastroenterol Rep* 2010;12:340-8.
4. Yan KS, Chia LA, Li X, et al. The intestinal stem cell markers *Bmi1* and *Lgr5* identify two functionally distinct populations. *Proc Natl Acad Sci U S A* 2012;109:466-71.
5. Barker N, van Es JH, Kuipers J, et al. Identification of stem cells in small intestine and colon by marker gene *Lgr5*. *Nature* 2007;449:1003-7.
6. Sangiorgi E, Capocchi MR. *Bmi1* is expressed *in vivo* in intestinal stem cells. *Nat Genet* 2008;40:915-20.
7. Montgomery RK, Carlone DL, Richmond CA, et al. Mouse telomerase reverse transcriptase (*mTert*) expression marks slowly cycling intestinal stem cells. *Proc Natl Acad Sci U S A* 2011;108:179-84.
8. Buczacki SJ, Zecchini HI, Nicholson AM, et al. Intestinal label-retaining cells are secretory precursors expressing *Lgr5*. *Nature* 2013;495:65-9.
9. Itzkovitz S, Lyubimova A, Blat IC, et al. Single-molecule transcript counting of stem-cell markers in the mouse intestine. *Nat Cell Biol* 2012;14:106-14.
10. Munoz J, Stange DE, Schepers AG, et al. The *Lgr5* intestinal stem cell signature: robust expression of proposed quiescent '+4' cell markers. *EMBO J* 2012;31:3079-91.
11. Tian H, Biehs B, Warming S, et al. A reserve stem cell population in small intestine renders *Lgr5*-positive cells dispensable. *Nature* 2011;478:255-9.
12. Canham MA, Sharov AA, Ko MS, et al. Functional heterogeneity of embryonic stem cells revealed through translational amplification of an early endodermal transcript. *PLoS Biol* 2010;8:e1000379.
13. Martinez Arias A, Brickman JM. Gene expression heterogeneities in embryonic stem cell populations: origin and function. *Curr Opin Cell Biol* 2011;23:650-6.
14. Galvin-Burgess KE, Travis ED, Pierson KE, et al. TGF-beta-superfamily signaling regulates embryonic stem cell heterogeneity: self-renewal as a dynamic and regulated equilibrium. *Stem Cells* 2013;31:48-58.
15. Turco MY, Furia L, Dietze A. Cellular heterogeneity during embryonic stem cell differentiation to epiblast stem cells is revealed by the *ShcD/RaLP* adaptor protein. *Stem Cells* 2012;30:2423-36.
16. MacArthur BD, Sevilla A, Lenz M, et al. Nanog-dependent feedback loops regulate murine embryonic stem cell heterogeneity. *Nat Cell Biol* 2012;14:1139-47.
17. Collins CA, Olsen I, Zammit PS, et al. Stem cell function, self-renewal, and behavioral heterogeneity of cells from the adult muscle satellite cell niche. *Cell* 2005;122:289-301.
18. Gibbs KD Jr, Gilbert PM, Sachs K, et al. Single-cell phospho-specific flow cytometric analysis demonstrates biochemical and functional heterogeneity in human hematopoietic stem and progenitor compartments. *Blood* 2011;117:4226-33.
19. Copley MR, Beer PA, Eaves CJ. Hematopoietic stem cell heterogeneity takes center stage. *Cell Stem Cell* 2012;10:690-7.
20. Lee da Y, Gianino SM, Gutmann DH. Innate neural stem cell heterogeneity determines the patterning of glioma formation in children. *Cancer Cell* 2012;22:131-8.
21. Narsinh KH, Sun N, Sanchez-Freire V, et al. Single cell transcriptional profiling reveals heterogeneity of human induced pluripotent stem cells. *J Clin Invest* 2011;121:1217-21.

Intestinal Stem Cell Heterogeneity

22. Janich P, Pascual G, Merlos-Suarez A, et al. The circadian molecular clock creates epidermal stem cell heterogeneity. *Nature* 2011;480:209-14.
23. Campbell F, Williams GT, Appleton MA, et al. Post-irradiation somatic mutation and clonal stabilisation time in the human colon. *Gut* 1996;39:569-73.
24. Schepers AG, Snippet HJ, Stange DE, et al. Lineage tracing reveals Lgr5+ stem cell activity in mouse intestinal adenomas. *Science* 2012;337:730-5.
25. Kim KM, Shibata D. Methylation reveals a niche: stem cell succession in human colon crypts. *Oncogene* 2002;21:5441-9.
26. Yatabe Y, Tavaré S, Shibata D. Investigating stem cells in human colon by using methylation patterns. *Proc Natl Acad Sci U S A* 2001;98:10839-44.
27. Lobachevsky PN, Radford IR. Intestinal crypt properties fit a model that incorporates replicative ageing and deep and proximate stem cells. *Cell Prolif* 2006;39:379-402.
28. Hormoz S. Stem cell population asymmetry can reduce rate of replicative aging. *J Theor Biol* 2013;331:19-27.
29. Snippet HJ, van der Flier LG, Sato T, et al. Intestinal crypt homeostasis results from neutral competition between symmetrically dividing Lgr5 stem cells. *Cell* 2010;143:134-44.
30. Zhu L, Gibson P, Currle DS, et al. Prominin 1 marks intestinal stem cells that are susceptible to neoplastic transformation. *Nature* 2009;457:603-7.
31. Pruitt SC, Freeland A, Kudla A. Cell cycle heterogeneity in the small intestinal crypt and maintenance of genome integrity. *Stem Cells* 2010;28:1250-9.
32. Basak O, van de Born M, Korving J, et al. Mapping early fate determination in Lgr5+ crypt stem cells using a novel Ki67-RFP allele. *EMBO J* 2014;33:2057-68.
33. Rompolas P, Mesa KR, Greco V. Spatial organization within a niche as a determinant of stem-cell fate. *Nature* 2013;502:513-8.
34. Lopez-Garcia C, Klein AM, Simons BD, et al. Intestinal stem cell replacement follows a pattern of neutral drift. *Science* 2010;330:822-5.
35. Ritsma L, Ellenbroek SI, Zomer A, et al. Intestinal crypt homeostasis revealed at single-stem-cell level by *in vivo* live imaging. *Nature* 2014;507:362-5.
36. Sunter JP, Appleton DR, De Rodriguez MS, et al. A comparison of cell proliferation at different sites within the large bowel of the mouse. *J Anat* 1979;129:833-42.
37. Gandara RM, Mahida YR, Potten CS. Regional differences in stem and transit cell proliferation and apoptosis in the terminal ileum and colon of mice after 12 Gy. *Int J Radiat Oncol Biol Phys* 2012;82:e521-8.
38. Leedham SJ, Rodenas-Cuadrado P, Howarth K, et al. A basal gradient of Wnt and stem-cell number influences regional tumour distribution in human and mouse intestinal tracts. *Gut* 2013;62:83-93.
39. Hughes KR, Gandara RM, Javkar T, et al. Heterogeneity in histone 2B-green fluorescent protein-retaining putative small intestinal stem cells at cell position 4 and their absence in the colon. *Am J Physiol Gastrointest Liver Physiol* 2012;303:G1188-201.
40. Middendorp S, Schneeberger K, Wiegerinck CL, et al. Adult stem cells in the small intestine are intrinsically programmed with their location-specific function. *Stem Cells* 2014;32:1083-91.
41. Karpowicz P, Zhang Y, Hogenesch JB, et al. The circadian clock gates the intestinal stem cell regenerative state. *Cell Rep* 2013;3:996-1004.
42. Mukherji A, Kobiita A, Ye T, et al. Homeostasis in intestinal epithelium is orchestrated by the circadian clock and microbiota cues transduced by TLRs. *Cell* 2013;153:812-27.
43. O'Brien CA, Pollett A, Gallinger S, et al. A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature* 2007;445:106-10.
44. Ricci-Vitiani L, Lombardi DG, Pilozzi E, et al. Identification and expansion of human colon-cancer-initiating cells. *Nature* 2007;445:111-5.
45. Dalerba P, Dylla SJ, Park IK, et al. Phenotypic characterization of human colorectal cancer stem cells. *Proc Natl Acad Sci U S A* 2007;104:10158-63.
46. Hill RP. Identifying cancer stem cells in solid tumors: case not proven. *Cancer Res* 2006;66:1891-5; discussion 1890.
47. Kelly PN, Dakic A, Adams JM, et al. Tumor growth need not be driven by rare cancer stem cells. *Science* 2007;317:337.
48. Barker N, Ridgway RA, van Es JH, et al. Crypt stem cells as the cells-of-origin of intestinal cancer. *Nature* 2009;457:608-11.
49. Schwitalla S, Fingerle AA, Cammareri P, et al. Intestinal tumorigenesis initiated by dedifferentiation and acquisition of stem-cell-like properties. *Cell* 2013;152:25-38.

50. Garcia SB, Park HS, Novelli M, et al. Field cancerization, clonality, and epithelial stem cells: the spread of mutated clones in epithelial sheets. *J Pathol* 1999;187:61-81.
51. Snippert HJ, Schepers AG, van Es JH, et al. Biased competition between Lgr5 intestinal stem cells driven by oncogenic mutation induces clonal expansion. *EMBO Rep* 2014;15:62-9.
52. Tang DG. Understanding cancer stem cell heterogeneity and plasticity. *Cell Res* 2012;22:457-72.
53. Leth-Larsen R, Terp MG, Christensen AG, et al. Functional heterogeneity within the CD44 high human breast cancer stem cell-like compartment reveals a gene signature predictive of distant metastasis. *Mol Med* 2012;18:1109-21.
54. Dieter SM, Ball CR, Hoffmann CM, et al. Distinct types of tumor-initiating cells form human colon cancer tumors and metastases. *Cell Stem Cell* 2011;9:357-65.
55. Fodde R, Brabletz T. Wnt/beta-catenin signaling in cancer stemness and malignant behavior. *Curr Opin Cell Biol* 2007;19:150-8.
56. Vermeulen L, De Sousa EMF, van der Heijden M, et al. Wnt activity defines colon cancer stem cells and is regulated by the microenvironment. *Nat Cell Biol* 2010;12:468-76.
57. Ong BA, Vega KJ, Houchen CW. Intestinal stem cells and the colorectal cancer microenvironment. *World J Gastroenterol* 2014;20:1898-909.
58. Buczacki SJ, Davies RJ. The confounding effects of tumour heterogeneity and cellular plasticity on personalized surgical management of colorectal cancer. *Colorectal Dis* 2014;16:329-31.