Association Between MTHFR Genetic Variants and Multiple Sclerosis in a Southern Iranian Population

Fakhraddin Naghibalhossaini^{1,2*}, Hesam Ehyakonandeh¹, Alireza Nikseresht², and Eskandar Kamali²

- 1. Department of Biochemistry, Shiraz University of Medical Sciences, School of Medicine, Shiraz, Iran.
- 2. Autoimmune Research Center, Shiraz University of Medical Sciences, School of Medicine, Shiraz, Iran.

Submmited 27 February 2015; Accepted 16 March 2015; Published 29 April 2015

Multiple sclerosis (MS) is a demyelinating neuro- inflammatory autoimmune disease of the central nervous system. Genetic predisposition has long been suspected in the etiology of this disease. The association between *MTHFR* polymorphisms and MS has been ivestigated in different ethnic groups. We investigated the association between *MTHFR C677T* and *A1298C* missense variants and MS in 180 patients and 231 age- and gendermatched healthy controls in a Southern Iranian population. The mutagenically separated PCR (MS-PCR) and PCR-RFLP methods were used to genotype *MTHFR* at position 677 and 1298, respectively. Compared with controls, we observed a strong association between two *MTHFR* variants and the risk of developing MS. Subjects carrying 677T allele (*CT* and *TT* genotypes) had increased susceptibility to MS as compared to those carrying *CC* genotype (odds ratio (OR) for *CT*= 2.9, 95% confidence interval (95% CI)= 1.88-4.49; OR for *TT*= 6.23, 95% CI= 3.08-12.59). The variant *1298AC* genotype also increased the risk for MS among our study population (OR= 2.14, 95% CI= 1.37-3.34). Combined genotype analysis for two *MTHFR* SNPs revealed that compared to the wild type genotypes (677CC/1298AA), 3 genotypes including *TT/AC*, *CT/AC*, and *TT/AA* were significantly at increased risk for MS development (OR= 13.9, 5.3, and 4.9, respectively). Our results suggest a possible gene dose- dependent association between *MTHFR* mutrant alleles and the risk of MS development.

Key words: Multiple sclerosis, MTHFR, SNP, genotype

ultiple sclerosis (MS) is a complex autoimmune disease of the central nervous system (CNS) resulting in CNS inflammation and demyelination of nerve axons (1, 2). A significant increase in MS incidence has been reported in Iran over the last decade, especially in females (3).

The MS etiology is not well-understood; however, several studies suggest that environmental and genetic factors might be involved in the etiology of this disease (4, 5). Age-dependant exposure to viral infection may also play a role in MS susceptibility (6).

Corresponding author: Department of Biochemistry, Shiraz University of Medical Sciences, School of Medicine, Zand Street, Shiraz, Iran. E-mail: fakhraddin.naghibalhossaini@mail.mcgill.ca.

Previous studies have indicated that MS patients have elevated plasma and cerebrospinal fluid (CSF) levels of neurotoxic amino acid homocysteine (7). Other studies have suggested that hyperhomocystenemia is a risk factor for dementia and Alzheimer's disease (8, 9) and might be associated with cognitive impairment in MS patients (10). Increased blood homocysteine was shown to be associated with the sensitization of neurons to oxidative stress that promotes apoptosis and hypersensitivity to excitotoxicity (11). Homocysteine may induce neurotoxicity through the oxidation of sulfhydryl groups resulting in generation of reactive oxygen species (11) and overstimulation of N-methyl-D-aspartate (NMDA) receptors resulting in neuronal damage due to excessive Ca²⁺ influx (12).

Methylenetetrahydrofolate reductase (MTHF-R) is a key folate metabolizing enzyme that functions at the junction between two critical pathways regulating one carbon metabolism, nucleotide synthesis and synthesizing the universal methyl donor S-adenosyl methionine (SAM). MTHFR gene is polymorphic and two common non-synonymous mutations, C677T (A222V; rs1801133) and A1298C (E429A; rs1801131), have been associated with decreased enzyme activity and the increased levels of plasma homocysteine (13-16). As a result, the MTHFR genotypes may play a role in MS susceptibility. Both of the abovementioned MTHFR polymorphisms have been extensively studied for associations with several diseases including neural tube defects (15, 17), and cardiovascular disease (18, 19). A few studies have also investigated the relationship between these polymorphisms and MS (20-22). The aim of this study was to investigate the association between functional polymorphisms of the MTHFR gene with MS among Southern Iranian population.

Materials and methods

Study population

This case- control study consisted of 180 unrelated patients and 231 healthy controls. The MS population was obtained from patients in university hospitals in Shiraz, Southern Iran, and the diagnosis was made by a neurologist according to the revised McDonald criteria (23). The associated MS population was comprised of three clinical subtypes: 128 relapsing-remitting MS (RR-MS; 71.1%), 43 secondary progressive MS (SP-MS; 23.9%), and 9 primary progressive MS (PP-MS; 5%). The control group was also obtained from healthy volunteers from the general population, which had been matched for age, gender, and ethnicity. Ethics approval for experimentation on humans was obtained from the Institutional Ethics Committee.

Genotype analysis

Genomic DNA was extracted from peripheral blood using a standard salting-out procedure (24). Genotyping of *MTHFR* at position 677 of DNA from healthy subjects (control) and multiple sclerotic patients was performed using the mutagenically separated PCR (MS-PCR) method, as previously described (25). The *A1298C* mutation of *MTHFR* was also examined by PCR-RFLP of DNA samples using the enzyme MboII (MBI Fermentas, Lithuania) as described previously (26).

Statistical analysis

All statistical analyzes were performed using the SPSS version 16 software package (SPSS Inc., Chicago, IL). Genotype and allele frequencies for the *MTHFR* genotype variants were investigated using standard Chisquare (χ^2) analysis. In addition, conditional multivariate logistic regression analysis for matched case- control groups was used to calculate odds ratio (OR) and 95% confidence intervals (95% CI). A p-value <0.050 was considered as statistically significant.

Results

We investigated the association between two common functional polymorphisms of MTHFR

(C677T and A1298C) and MS incidence among Iranian patients. The case-control populations consisted of 180 patients and 231 healthy controls. The cases were more likely to be females (74%) older than 26 years (54.4%). The mean age of patients was 26.0± 12.3 years. The groups of patients and controls did not significantly differ concerning gender or age.

MTHFR C677T and A1298C genotyping was performed by MS-PCR and PCR-RFLP methods, respectively. Illustrative examples of genotype analysis of the two MTHFR variant genotypes are shown in Fig. 1. The results of genotype frequencies and odds ratios for MTHFR genotypes and MS are presented in Table 1. There was no significant HardyWeinberg disequilibrium concerning the MTHFR 677 and 1298 genotypes in controls. The distribution of MTHFR 677 genotypes among patients also agreed with that expected from the Hardy-Weinberg equilibrium ($\chi^2 = 0.5$, P= 0.47). However, significant departures from Hardy-Weinberg equilibrium were observed for MTHFR 1298 genotypes among cases (P= 0.00).

In our study, the allele frequency distributions of MTHFRC 677T were significantly different between patient and control groups (41.9% versus 20.3%, P= 0.00). The frequencies of the MTHFR C677T genotypes in the patients (CC, 35%; CT, 46.1%; TT, 18.9%) were also significantly different from controls (CC, 65%; CT, 29.5%; TT, 5.5%) (P= 0.00; Table 1). The MS patients were presented with higher homozygous TT genotype than control group, as manifested by an odds ratio of 6.23 (95% CI= 3.08- 12.59). This could be translated into that people having the TT genotype are 6.23- fold more at risk of developing MS, in the multivariate logistic regression analysis (Table 1). Under the codominant model of inheritance, the CT genotype was also associated with an increased risk for MS with an odds ratio lower than recessive model (adjusted OR= 2.9, 95% CI= 1.88- 4.49). When we combined heterozygous and homozygous variant genotypes, the adjusted OR for the CT/TT genotypes was 3.4 (95% CI= 2.29- 5.17). In casecase comparisons, we observed no differences in frequencies of MTHFR C677T genotypes in patients stratified by the clinicopathologic variables, including age, sex, and disease type (data not shown).




Fig. 1. Representative examples of genotyping of *MTHFR* positions 677 and 1298. **A.** MS-PCR assay for the genotyping of *MTHFR C677T* polymorphism. The 677C alleles (168 base pair product) were separated from the 677T alleles (148-base pair product) by electrophoresis on a 2.5 % agarose gel. **B.** PCR-RFLP assay for genotyping of *MTHFR A1298C* polymorphisms. Digestion of the 163 bp PCR product of the 1298A allele yields five fragments of 56, 31, 30, 28, and 18 base pairs, whereas the 1298C allele results in four PCR bands of 84, 31, 30, and 18 base pairs. The digested PCR products were separated by electrophoresis on a 2.5% agarose gel. The three possible genotypes are discernible by detection of the 84and 56 bp fragments.

For the A1298C polymorphism, while the allele frequency distribution in MS patients was almost the same as that of controls (36.7% versus 34.7%), the genotype frequency distributions were significantly different (P = 0.001; Table 1). Analysis of the MTHFR A1298C frequency data obtained in this study showed that the heterozygotes (AC) were overrepresented in MS patients (67.8% versus 45.7%, P= 0.00) and a small trend for a higher frequency of the homozygous mutant genotype (CC) was observed in controls

(11.8% versus 2.8%, P= 0.04). According to the logistic regression model, in the entire group of patients, *MTHFR AC* and *AC+CC* genotypes were strongly associated with a higher risk of MS incidence (Table 1). The adjusted OR for *AC* and *AC+CC* genotypes were 2.14 (95% CI 1.37- 3.34) and 1.77 (95%CI 1.15- 2.73), respectively. In casecase comparison, no statistically significant differences in frequencies of the *MTHFR A1298C* genotypes were found in patients stratified by age, sex, and disease type (data not shown).

MTHFR	Genotypes & alleles	Patients N (%)	Controls	OR (95% CI)	P *
C(77T	& alleles	180	N (%)	(95% C1)	
C677T	CC (··· f)		231	1	
Total	CC (ref)	63 (35)	150 (65)	1	
	CT	83 (46.1)	68 (29.5)	2.9 (1.88- 4.49)	0.00
	TT	34 (18.9)	13 (5.5)	6.23 (3.08-12.59)	0.00
	CT+TT	117 (65)	81 (35.1)	3.44 (2.29-5.17)	0.00
	C	209 (58.1)	368 (79.7)	1	0.00
	T	151 (41.9)	94 (20.3)	2.83 (2.08-3.85)	
					0.00
		180	186		
A1298C	AA (ref)	53 (29.4)	79 (42.5)	1	
Total	AC	122 (67.8)	85 (45.7)	2.14 (1.37-3.34)	
	CC	5 (2.8)	22 (11.8)	0.34 (0.12-0.950)	0.001
	AC+CC	127 (70.6)	107 (57.5)	1.77 (1.15-2.73)	0.04
	A	228 (63.3)	243 (65.3)	1	0.010
	C	132 (36.7)	129 (34.7)	1.09 (0.81-1.48)	0.574
Combined genotypes		180	186		0.28
(677/1298)	CC/AA (ref)	22 (12.2)	46 (24.7)	1	0.13
Total	CC/AC	39 (21.7)	57 (30.8)	1.43 (0.75- 2.74)	0.50
	CC/CC	2 (1.1)	14 (7.5)	0.30 (0.06- 1.43)	0.00
	CT/AA	17 (9.4)	27 (14.5)	1.32 (0.60- 2.91)	0.77
	CT/AC	63 (35)	25 (13.4)	5.27 (2.65- 10.48)	0.004
	CT/CC	3 (1.7)	5 (2.7)	1.26 (0.28- 5.73)	0.00
	TT/AA	14 (7.8)	6 (3.2)	4.87 (1.65-14.41)	ND
	TT/AC	20 (11.1)	3 (1.6)	13.94 (3.74-	
	TT/CC	0(0.00)	3 (1.6)	51.95)	
	21,00	0(0.00)	0 (1.0)	ND	

^{*}For OR and 95% CI calculations, controls with the wild-type CC and AA MTHFR genotypes were used as reference category. ND: not determined.

For *MTHFR* 677 and 1298 combined genotypes, double heterozygotes (677 CT/1298 AC) had 5.3- fold (95% CI 2.65- 10.48) increased risk

compared with the wild- type (677 CC/1298 AA) genotypes of controls (Table 1). Individuals with the 677TT/1298AC and 677TT/1298AA genotypes

were also at higher risk of developing MS in comparison to controls. The adjusted OR for *TT/AC* and *TT/AA* combined genotypes were 13.94 (95% CI 3.74- 51.95) and 4.87 (95% CI 1.65- 14.41), respectively. Due to the small number of cases in the current study, it was not possible to perform the analyses of MS risk associated with double homozygous mutants of *677TT/1298CC* genotypes.

Discussion

Previous studies have suggested an influence of genetic factors in the aetiology of multiple sclerosis; however, the underlying molecular mechanisms of MS remain unidentified (27, 28). It has been reported that MS patients have elevated levels of plasma and cerebrospinal fluid homocysteine, a neurotoxic metabolite (7, 10, 29-31).

MTHFR deficiency is the most common genetic cause of hyperhomocysteinemia (13, 16). MTHFR enzyme reduces 5, 10-methylenetetrahydrofolate to 5-methyltetrahydrofolate, which is required for remethylation and conversion of homocysteine to methionine.

The results of previous studies with regard to the association between two polymorphisms of MTHFR with reduced enzyme activity (C677T and A1298C) and MS risk have been inconsistent. We examined the relationship between MTHFR C677T and A1298C polymorphisms and the risk of MS in a Southern Iranian population. According to the logistic regression model, in the entire group of patients, **MTHFR** C677T and A1298C polymorphisms were strongly associated with a higher risk of MS (Table 1). Compared with controls, the MTHFRC677T genotype showed a higher risk of MS incidence both in the recessive and codominant models (for TT versus CC: OR= 6.23, 95 % CI= 3.08-1 2.59 and CT versus CC: OR= 2.9, 95 % CI= 1.88- 4.49, respectively). We also found a higher risk associated with the MTHFR 1298AC genotypes when the MS patients were compared with controls and the wild- type AA genotype was used as a reference category (OR= 2.14, 95% CI= 1.37-3.34). However, under the recessive model of inheritance, the homozygous CC genotype was slightly associated with a decreased risk for MS (OR= 0.34, 95% CI= 0.12-0.950). Such result could be due to low statistical power because of the limited number of homozygous subjects (5 out of 180 cases). It has been previously reported that compared to the 677T allele, 1298 C allele has a minor effect on MTHFR activity. Neither the homozygous nor the heterozygous state of the MTHFR 1298C genotype is associated with higher plasma homocysteine or a lower plasma folate concentration—that was observed with allele. However, double homozygous 677T heterozygosity for both MTHFR mutations results in similar features as observed in homozygotes for the 677T allele (15). When we considered both MTHFR 1298CC and CT genotypes, we found the increased risk of MS incidence associated with the AC/CC genotypes in the entire group of patients (Table 1). Our finding is consistent with most previous reports in which an increased risk of MS was observed associated with the MTHFR A1298C genotype (20, 32). This is in contrast to another study (21) that found no association between the MTHFR A1298C polymorphism and MS in Australian population. In agreement with a previous report from Iran (33), we also found increased risk of MS associated with the MTHFR C677T genotype. A non- significant increased MS risk associated with the C677T variant genotype was also reported in a group of Australian population (22). However, some studies conducted on relatively small groups of cases, observed no association between MS and MTHFR C677T polymorphism (20, 22, 32). Both the C677T and A1298C variant genotypes of MTHFR have been associated with decreased enzyme activity, with the

C677T having a more severe effect than the A1298C variant. In vitro studies have shown that the 677TT and 1298CC variant genotypes have 60% and 30% reduced enzyme activity in comparison to the wild type MTHFR genotypes, respectively (16, 18, 34). Inconsistency of findings across studies could be explained with the differences in study designs, sample size, genotyping methods, racial, nutritional, and other environmental factors.

To examine the joint effect of MTHFR 677 and 1298 genotypes on MS risk, we analyzed the relationship between combined MTHFR SNPs at these loci and MS risk. Out of five major combined genotypes (CC/AC, CT/AA, CT/AC, TT/AA, and TT/AC) that constitute almost 80% of all genotypic diversity in our study group, the distribution of three genotypes was significantly different between cases and controls (Table 1). Based on our findings, 3 genotypes with the mutant alleles (TT/AC,CT/AC, and TT/AA) are high-risk genotypes for developing MS (13.9, 5.3, and 4.9-fold increased risk, respectively). According to a previous study, subjects with the combined 677CT/1298AC heterozygosity had significantly higher fasting serum homocysteine levels compared to those that were C677T heterozygous (16). Since both C677T and A1298C mutations can influence MTHFR activity and plasma homocysteine concentrations (15, 35), it is intriguing to beleive that mutated alleles of MTHFR increase the risk for developing MS in a gene dose-related manner.To our knowledge, the combined effect of MTHFR genotypes has not been previously analyzed in MS populations and further studies are necessary to understand the dose-dependent association of MTHFR alleles with MS risk.

Acknowledgments

This study was part of the dissertation of Hesam Ehyakonandeh, submitted in partial fulfillment of the requirements for the MSc in biochemistry. This work was supported by the office of the Vice Chancellor for Research, Shiraz University of Medical Sciences, and grant number 90-01-17-3031.

Conflict of interests

The authors declared no conflict of interests.

References

- 1. Hafler DA. Multiple sclerosis. J Clin Invest 2004;113:788-94.
- 2. Rosati G. The prevalence of multiple sclerosis in the world: an update. Neurol Sci 2001;22:117-39.
- 3. Etemadifar M, Sajjadi S, Nasr Z, et al. Epidemiology of multiple sclerosis in Iran: a systematic review. Eur Neurol 2013;70:356-63.
- 4. Dyment DA, Ebers GC, Sadovnick AD. Genetics of multiple sclerosis. Lancet Neurol 2004;3:104-10.
- Baranzini SE, Wang J, Gibson RA, et al. Genome-wide association analysis of susceptibility and clinical phenotype in multiple sclerosis. Hum Mol Genet 2009;18:767-78.
- Levin LI, Munger KL, Ruberstone MV. Temporal relationship between elevation of Epstein–Barr virus antibody titers and initial onset of neurological symptoms in multiple sclerosis JAMA 2005;293:2496–500.
- 7. Ramsaransing GS, Fokkema MR, Teelken A, et al. Plasma homocysteine levels in multiple sclerosis. J Neurol Neurosurg Psychiatry 2006;77:189-92.
- 8. Kujala P, Portin R, Ruutiainen J. The progress of cognitive decline in multiple sclerosis. A controlled 3-year follow-up. Brain 1997;120 (Pt 2):289-97.
- 9. Seshadri S, Beiser A, Selhub J, et al. Plasma homocysteine as a risk factor for dementia and Alzheimer's disease. N Engl J Med 2002;346: 476-83.
- Russo C, Morabito F, Luise F, et al. Hyperhomocysteinemia is associated with cognitive impairment in multiple sclerosis. J Neurol 2008;255:64-9.
- 11. Kruman, II, Culmsee C, Chan SL, et al. Homocysteine elicits a DNA damage response in neurons that promotes apoptosis and hypersensitivity to excitotoxicity. J Neurosci 2000;20:6920-6.
- Lipton SA, Kim WK, Choi YB, et al. Neurotoxicity associated with dual actions of homocysteine at the N-methyl-Daspartate receptor. Proc Natl Acad Sci U S A 1997;94:5923-8.
- 13. Rozen R. Genetic predisposition to hyperhomocysteinemia: deficiency of methylenetetrahydrofolate reductase (MTHFR).

MTHFR Genetic Variants and Multiple Sclerosis

Thromb Haemost 1997;78:523-6.

- 14. Bagley PJ, Selhub J. A common mutation in the methylenetetrahydrofolate reductase gene is associated with an accumulation of formylated tetrahydrofolates in red blood cells. Proc Natl Acad Sci U S A 1998;95:13217-20.
- 15. van der Put NM, Gabreels F, Stevens EM, et al. A second common mutation in the methylenetetrahydrofolate reductase gene: an additional risk factor for neural-tube defects? Am J Hum Genet 1998;62:1044-51.
- 16. Weisberg IS, Jacques PF, Selhub J, et al. The 1298A-->C polymorphism in methylenetetrahydrofolate reductase (MTHFR): in vitro expression and association with homocysteine. Atherosclerosis 2001;156:409-15.
- 17. de Franchis R, Botto LD, Sebastio G, et al. Spina bifida and folate-related genes: a study of gene-gene interactions. Genet Med 2002;4:126-30.
- 18. Frosst P, Blom HJ, Milos R, et al. A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. Nat Genet 1995;10:111-3.
- 19. Markan S, Sachdeva M, Sehrawat BS, et al. MTHFR 677 CT/MTHFR 1298 CC genotypes are associated with increased risk of hypertension in Indians. Mol Cell Biochem 2007;302:125-31.
- 20. Klotz L, Farkas M, Bain N, et al. The variant methylenetetrahydrofolate reductase c.1298A>C (p.E429A) is associated with multiple sclerosis in a German case-control study. Neurosci Lett 2010;468:183-5.
- 21. Szvetko AL, Fowdar J, Nelson J, et al. No association between MTHFR A1298C and MTRR A66G polymorphisms, and MS in an Australian cohort. J Neurol Sci 2007;252:49-52.
- 22. Tajouri L, Martin V, Gasparini C, et al. Genetic investigation of methylenetetrahydrofolate reductase (MTHFR) and catechol-O-methyl transferase (COMT) in multiple sclerosis. Brain Res Bull 2006;69:327-31.
- 23. Polman CH, Reingold SC, Edan G, et al. Diagnostic criteria for multiple sclerosis: 2005 revisions to the "McDonald Criteria". Ann Neurol 2005;58:840-6.
- 24. Lahiri DK, Bye S, Nurnberger JI, Jr., et al. A non-organic and non-enzymatic extraction method gives higher yields of genomic DNA from whole-blood samples than do nine other

- methods tested. J Biochem Biophys Methods 1992;25:193-205.
- 25. Mokarram P, Naghibalhossaini F, Saberi Firoozi M, et al. Methylenetetrahydrofolate reductase C677T genotype affects promoter methylation of tumor-specific genes in sporadic colorectal cancer through an interaction with folate/vitamin B12 status. World J Gastroenterol 2008;14:3662-71.
- 26. Naghibalhossaini F, Mokarram P, Khalili I. Easy detection of 5,10-methylenetetrahydrofolate reductase 1298A/C genotype by mutagenically separated PCR assay. Clin Chem Lab Med 2008;46:987-9.
- 27. Goertsches R, Villoslada P, Comabella M, et al. A genomic screen of Spanish multiple sclerosis patients reveals multiple loci associated with the disease. J Neuroimmunol 2003;143:124-8.
- 28. Jonasdottir A, Thorlacius T, Fossdal R, et al. A whole genome association study in Icelandic multiple sclerosis patients with 4804 markers. J Neuroimmunol 2003;143:88-92.
- 29. Besler HT, Comoglu S. Lipoprotein oxidation, plasma total antioxidant capacity and homocysteine level in patients with multiple sclerosis. Nutr Neurosci 2003;6:189-96.
- 30. Vrethem M, Mattsson E, Hebelka H, et al. Increased plasma homocysteine levels without signs of vitamin B12 deficiency in patients with multiple sclerosis assessed by blood and cerebrospinal fluid homocysteine and methylmalonic acid. Mult Scler 2003;9:239-45.
- 31. Aksungar FB, Topkaya AE, Yildiz Z, et al. Coagulation status and biochemical and inflammatory markers in multiple sclerosis. J Clin Neurosci 2008;15:393-7.
- 32. Fekih Mrissa N, Mrad M, Klai S, et al. Association of methylenetetrahydrofolate reductase A1298C polymorphism but not of C677T with multiple sclerosis in Tunisian patients. Clin Neurol Neurosurg 2013;115:1657-60.
- 33. Alatab S, Hossein-nezhad A, Mirzaei K, et al. Inflammatory profile, age of onset, and the MTHFR polymorphism in patients with multiple sclerosis. J Mol Neurosci 2011;44:6-11.
- 34. Rozen R. Molecular genetics of methylenetetrahydrofolate reductase deficiency. J Inherit Metab Dis 1996;19:589-94.
- 35. De Marco P, Calevo MG, Moroni A, et al. Study of MTHFR and MS polymorphisms as risk factors for NTD in the Italian population. J Hum Genet 2002;47:319-24.