Diversity of Variable Number Tandem Repeat Loci in Shigella Species Isolated from Pediatric Patients

Reza Ranjbar¹, Mojtaba Memariani^{1*}, Hamed Memariani^{2,3}

- 1. Molecular Biology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran.
- 2. Department of Molecular and Cell Biology, Faculty of Basic Sciences, University of Mazandaran, Babolsar, Iran.
- 3. Biotechnology Research Center, Pasteur Institute of Iran, Tehran, Iran.

Submmited 24 March 2015; Accepted 3 May 2015; Published 30 June 2015

Multilocus variable number tandem repeat (VNTR) analysis (MLVA) is a new typing method with several advantages compared to other methods. Dissemination of *Shigella* is highly significant in developing countries. Whilst *Shigella* is becoming increasingly important as an etiologic agent of pediatric shigellosis in Iran, little is known about the genetic diversity of the local strains. Therefore, the aim of this study was to describe the genetic diversity of *Shigella* species isolated from pediatric patients in Tehran, Iran. A total of 53 *Shigella* isolates were obtained from 1070 patients with diarrhea (less than 12 years of age). All isolates were identified by routine biochemical and serological tests. The confirmed *Shigella* isolates were further serogrouped (by the slide agglutination) using slide agglutination method. MLVA assay with the seven loci resolved 53 *Shigella* isolates into 36 different genotypes. Almost all the isolates were classified into five clonal complexes. Furthermore, our MLVA assay could effectively distinguish the four *Shigella* species. This study has provided valuable insights into the genetic heterogeneity of *Shigella* species in Tehran, Iran. Our findings can be helpful for further epidemiological surveillance of *Shigella* species in this country in the future.

Key words: Shigella, MLVA, genotyping

Diarrheal diseases are one of the most important causes of children mortality, accounting for around two million deaths annually. Bacillary dysentery, also known as shigellosis, causes a significant proportion of morbidity and mortality, especially in children with diarrhea in developing countries (1). Shigellosis is primarily a disease of poor, crowded communities which do not have adequate sanitation or clean water. The genus

Shigella includes four species: S. dysenteriae, S. flexneri, S. boydii, and S. sonnei. Of these species, shigellosis is predominantly caused by S. flexneri in developing countries especially in Asia, whereas S. sonnei is predominant in industrialized countries (2, 3).

Genotyping methods provide useful information for establishing the genetic relatedness among isolates for the purposes of epidemiological

^{**}Corresponding author: Molecular Biology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran. E-mail: memaryani@gmail.com

investigation and evolutionary studies. A variety of molecular typing methods have been developed for Shigella species, including pulsed- field gel electrophoresis (PFGE) (4-6), multilocus sequence typing (MLST) (7), multilocus variable-number tandem-repeat (VNTR) analysis (MLVA) (8-11), ribotyping (12-15), plasmid profiling (4, 16) and repetitive sequence-based PCR (Rep-PCR) (7, 17). Although PFGE is still a golden standard for genotyping and source tracking of food-borne pathogens, it is laborious, expensive, often difficult to interpret, and requires rigorous standardization. Furthermore, it needs experienced personnel in order to achieve reliable, consistent, and reproducible results. By contrast, MLVA is a PCRbased genotyping method which is rapid, relatively cheap and easy to perform. This method targets multiple VNTR loci and relies on the detection of different copy numbers inside each locus (18).

Dissemination of *Shigella* is highly significant in developing countries. Whilst *Shigella* is becoming increasingly important as an etiologic agent of pediatric shigellosis in Iran (2, 3, 19), little is known about the genetic diversity of the local strains. To the best of our knowledge, this is the first study on genotype of *Shigella* by MLVA in Iran. Therefore, the aim of this study was to describe the genetic diversity of *Shigella* species isolated from pediatric patients in Tehran, Iran.

Materials and methods

Bacterial strains

In this study, from June 2008 to October 2010, a total of 53 *Shigella* isolates were obtained from 1070 patients with diarrhea (less than 12 years of age) in Tehran, Iran. All of the isolates were identified by routine biochemical and serological tests. The confirmed *Shigella* isolates were further serogrouped by the slide agglutination test (Mast Diagnostic, Merseyside, UK). The verified isolates were preserved at -70°C in tripticase soy broth

(TSB) with 25% (v/v) glycerol for further analysis (3). All ethical issues were considered. Life, health, dignity, integrity, right to self-determination, privacy, and confidentiality of personal information of research subjects were protected in this study.

DNA extraction

Each *Shigella* isolate was plated on nutrient agar and incubated overnight at 37 $^{\circ}$ C. A single colony was removed from the plate, suspended in 200 μ l of sterile deionized water and boiled for 15 min. After centrifugation at 8,000x g for 6 min, the supernatant was transferred into a new tube for subsequent PCR analysis.

MLVA assay

The VNTR loci selected along with the primers used for MLVA genotyping were those of Gorge et al., which were seven loci; ms06, ms07, ms09, ms11, ms21, ms23, and ms32 (20). The list of primers used are shown in Table 1. For each loci, PCR was performed in 25 µl volume containing 1X PCR buffer (50 mmol/L KCL, 10 mmol/L Tris, pH 9), 2.5 mmol/L MgCl₂, 0.2 mmol/L of each primer with 0.5 U TaqDNA polymerase (CinnaGen Co., Iran), and 3 µl of crude DNA extract. Cycling conditions for all PCR reactions were 93 °C for 5 min, followed by 34 cycles of 93 °C for 30 s, 55 °C for 1 min, and 72 °C for 1 min. A final extension of 72 °C for 6 min was also employed. The PCR products were run on 1.5% agarose gels, stained with ethidium bromide, and visualized under UV transillumination.

Data analysis

The repeat units of each locus were determined by subtracting the size of flanking regions (offset size) from PCR amplicon size and then dividing the difference by the repeat size. The result was then rounded to the nearest integer value to simplify reporting of MLVA data. Repeat units were imported into Microsoft Excel. The minimum spanning tree (MST) was constructed with a categorical coefficient based on allelic profiles of

the *Shigella* strains using the trial version of Ridom MLVA compare software (Ridom® GmbH, Germany). MST is a convenient complementary tool to cluster multiple isolates and visualize the relative diversity within different lineage. A dendrogram of genetic relationships was also generated using the unweighted pair group method with arithmetic averages (UPGMA) method (21). *S.flexneri* Sf2457T (serotype 2a) was also included as an outgroup.

Results

Of the 53 Shigella isolates, S. sonnei (50.9%, n=27) was the most common, followed by S. flexneri (35.8%, n= 19), S. boydii (9.4%, n= 5), and S. dysenteriae (3.8%, n= 2). The age distribution data revealed that Shigella was isolated from 40 (75.5%) cases in the 2-5 year-old, 11 (20.7%) in the 6-12 year-old and 2 (3.8%) in \leq one year-old age Furthermore, this difference statistically significant (P< 0.001), suggesting that the peak age of shigellosis was 2-5 years in children. MLVA based on seven VNTR loci was carried out to characterize the Shigella isolates (Figure 1). Overall, Shigella isolates were discriminated into 36 different MLVA types (genotypes). Almost all of the strains were classified into five clonal complexes (CCs). CCs are defined as a group of allelic profiles in which

every profile shares at least 5 loci in common with at least an other member of the group. MST analysis showed that most of the S. sonnei strains fell into one CC (i.e. CC2), whereas most of the S. flexneri strains were categorized into three CCs (i.e. CC1, CC3, and CC5) (Figure 2). The UPGMA dendrogram with detail information including isolate code, species, and allelic profile is shown in Figure 3. All the Shigella isolates were categorized into two main clusters (i.e. cluster A and cluster B). Cluster A consisted of two subclusters, which represented most of our isolates (n=47). Interestingly, subcluster A1 contains only S. sonnei isolates, whereas subcluster A2 is relatively diverse. Cluster B only includes some strains of S. boydii and S. flexneri.

Discussion

In developing countries, poor hygiene, lack of clean water supply and overcrowded living conditions facilitate *Shigella* transmission, mainly through direct oral-fecal contamination. While *S. flexneri* is primarily endemic in developing world, *S. sonnei* is more commonly reported in industrialized countries. Based on a surveillance study in six countries across Asia, *S. flexneri* was the most frequently isolated *Shigella* species in Bangladesh, China, Indonesia, Pakistan, and Vietnam, while *S. sonnei* was the predominant species in Thailand.




Fig. 1. Polymorphism of ms07, ms09, ms11, and ms32 loci in different *Shigella* species. The number of repeats can be directly deduced by manual reading. The numbers above the PCR amplicons provide the repeat numbers of each VNTR locus. Lanes M, DNA markers.




Fig. 2. Minimum spanning tree (MST) for *Shigella* isolates. Each circle represents one strain with a unique genotype (MLVA profile). The size of the circles indicates the number of isolates. The number of loci which differ between two MLVA types is indicated on the lines connecting the MLVA types. Clonal Complexes (CCs) were indicated by grey halos. The color of the circles corresponds to *Shigella* species. *S. flexneri* Sf2457T [serotype 2a] was also used as an outgroup.

Table 1. PCR primers for specific VNTR loci and their corresponding tandem repeat (TR) and offset sizes in
reference strain S. flexneri 2457T chromosome.

VNTR locus	Primer sequence (5' to 3')	Tandem Repeat (TR) size, bp	Offset size, bp	Referenc es
ms06	F: AAA CGG GAG AGC CGG TTA TT R: TGT TGG TAC AAC GGC TCC TG	39	223	20
ms07(CVN001)	F: GTC AGT TCG CCC AGA CAC AG R: CGG TGT CAG CAA ATC CAG AG	39	353	20
ms09	F: GTG CCA TCG GGC AAA ATT AG R: CCG ATA AGG GAG CAG GCT AGT	179	177	20
ms11	F: GAA ACA GGC CCA GGC TAC AC R: CTG GCG CTG GTT ATG GGT AT	96	286	20
ms21	F: GCT GAT GGC GAA GGA GAA GA R: GGG AGT ATG CGG TCA AAA GC	141	82	20
ms23	F: GCT CCG CTG ATT GAC TCC TT R: CGG TTG CTC GAC CAC TAA CA	375	209	20
ms32	F: GAG ATT GCC GAA GTG TTG C R: AAC TGG CGG CGT TTA TCA AG	101	254	20

However, *S. boydii* is mainly restricted to the Indian subcontinent, being the second most abundant species isolated in Bangladesh (22). Similar to many previous studies from developing countries (22-24), *S. flexneri* is still predominant species in many parts of Iran (25, 26). However, in the present study, almost half of the isolates belonged to *S.*

sonnei which may suggest the possible succession of *S. flexneri* by *S. sonnei* in urban areas in Iran as the standard of living improved, as inferred from observations obtained from industrialized countries (2, 3).

MLVA is an effective molecular typing technique based on counting the number of VNTR




Fig. 3. UPGMA analysis of Shigella isolates based on VNTR profiles. The highlighted colors represent Shigella species. S. flexneri Sf2457T [serotype 2a] was also used as an outgroup.

repeats and has advantages such as rapidity, ease, and convenience of interpretation. Since VNTR loci have various degrees of variability, they are useful molecular markers for phylogenetic and evolutionary study of bacterial pathogens. MLVA has been applied to investigate the clonal relationships among isolates of E. coli and Shigella species (20). In this study, we used MLVA profiles to establish phylogenetic relationships among 53 Shigella isolates. Moreover, our MLVA assay could effectively distinguish the four Shigella species. In the current survey, S. flexneri isolates showed higher diversity compared to S. sonnei isolates. This is not a surprise because unlike S.sonnei, which has only one serotype, S. flexneri has eight serotypes with at least 12 subserotypes (27). In recent years, several MLVA assays have been developed for Shigella species. In a study conducted by Ou et al. (2012) in Beijing, China, 180 strains of S. flexneri and S. sonnei were divided into 84 MLVA subtypes based on 10 VNTR loci (28). In a study by Koh et al. 40 strains of S. sonnei strains were isolated during the years 1997-2000, and during 2007-2009 they were classified into different clones based on 7 VNTR loci. They demonstrated that S. sonnei isolates with no epidemiological linkage were clustered together which could be due to traveling within the country and/or person-to-person spread of a particular strain over a long period of time with minor genetic changes (8). Izumiya et al. analyzed S. sonnei strains isolated from cases associated with foreign travel in Japan by MLVA and they showed that these isolates could be grouped into several clusters based on their countries of origin (11). In another study, Chiou et al. used 26 VNTR loci for phylogenetic analysis of 916 S. sonnei isolates collected in Taiwan over 9 years (10). Wang et al. also developed and evaluated MLVA assay based on 36 VNTR loci for discriminating the different serotypes of S. flexneri (27). Generally, these

studies showed that MLVA has the potential to replace other genotyping methods as a standard method for *Shigella* typing. However, lack of standardization of the methodology and interpretive criteria is problematic and hinders comparison of data between laboratories (9).

On the other hand, it is important to note that most of these mentioned MLVA schemes require a high precision of DNA length measurement, for instance, by microcapillary electrophoresis and fluorescent markers because the selected VNTRs have very short repeat units. Moreover, developing countries have limited accessibility to such equipment and their expenditures would be a major obstacle for many laboratories (20, 29). In this study, we exploited VNTR markers which can be easily analyzed by eye on agarose gels. Hence, this assay can be performed in a laboratory equipped with simple equipment.

The present study provided valuable insights into the genetic heterogeneity of *Shigella* species in Tehran. We hope that our findings can be helpful for further epidemiological surveillance of *Shigella* species in our country in the future. Finally, this study showed that MLVA is a promising typing technique for epidemiological studies of *Shigella* species due to low cost, the speed of analysis, and the ability to produce numerical data which can be easily shared among laboratories.

Acknowledgement

The authors are grateful to the Molecular Biology Research Center, of Baqiyatallah University of Medical Sciences, Tehran, Iran.

Conflict of interests

The authors declared no conflict of interests.

Rferences

- 1. Boschi-Pinto C, Velebit L, Shibuya K. Estimating child mortality due to diarrhoea in developing countries. Bull World Health Organ 2008;86:710-7.
- 2. Farshad S, Ranjbar R, Hosseini M. Molecular Genotyping of

- Shigella sonnei Strains Isolated From Children With Bloody Diarrhea Using Pulsed Field Gel Electrophoresis on the Total Genome and PCR-RFLP of IpaH and IpaBCD Genes. Jundishapur J Microbiol 2015;8:e14004.
- 3. Ranjbar R, Ghazi FM, Farshad S, et al. The occurrence of extended-spectrum beta-lactamase producing Shigella spp. in Tehran, Iran J Microbiol 2013;5:108-12.
- Pichel M, Gonzalez Fraga S, Terragno R, et al. Short report: analysis of clonal relationship among Shigella sonnei isolates circulating in Argentina. Epidemiol Infect 2007;135:681-7.
- Ruekit S, Wangchuk S, Dorji T, et al. Molecular characterization and PCR-based replicon typing of multidrug resistant Shigella sonnei isolates from an outbreak in Thimphu, Bhutan. BMC Res Notes 2014;7:95.
- Jin YH, Oh YH, Jung JH, et al. Antimicrobial resistance patterns and characterization of integrons of Shigella sonnei isolates in Seoul, 1999-2008. J Microbiol 2010;48:236-42.
- 7. Cao Y, Wei D, Kamara IL, et al. Multi-Locus Sequence Typing (MLST) and Repetitive Extragenic Palindromic Polymerase Chain Reaction (REP-PCR), characterization of shigella spp. over two decades in Tianjin China. Int J Mol Epidemiol Genet 2012;3:321-32.
- 8. Koh XP, Chiou CS, Ajam N, et al. Characterization of Shigella sonnei in Malaysia, an increasingly prevalent etiologic agent of local shigellosis cases. BMC Infect Dis 2012;12:122.
- Chiou CS, Izumiya H, Thong KL, et al. A simple approach to obtain comparable Shigella sonnei MLVA results across laboratories. Int J Med Microbiol 2013;303:678-84.
- 10. Chiou CS, Watanabe H, Wang YW, et al. Utility of multilocus variable-number tandem-repeat analysis as a molecular tool for phylogenetic analysis of Shigella sonnei. J Clin Microbiol 2009;47:1149-54.
- 11. Izumiya H, Tada Y, Ito K, et al. Characterization of Shigella sonnei isolates from travel-associated cases in Japan. J Med Microbiol 2009;58:1486-91.
- 12. Cabrera R, Echeita A, Herrera S, et al. Antibiotic resistance, plasmid profile and ribotyping in Cuban Shigella sonnei strains. Rev Esp Quimioter 2006;19:76-8.
- 13. Hinojosa-Ahumada M, Swaminathan B, Hunter SB, et al. Restriction fragment length polymorphisms in rRNA operons for subtyping Shigella sonnei. J Clin Microbiol 1991;29:2380-4.

- Nastasi A, Pignato S, Mammina C, et al. rRNA gene restriction patterns and biotypes of Shigella sonnei. Epidemiol Infect 1993;110:23-30.
- 15. Lee TM, Chang LL, Chang CY, et al. Molecular analysis of Shigella sonnei isolated from three well-documented outbreaks in school children. J Med Microbiol 2000;49:355-60.
- 16. Farshad S, Sheikhi R, Japoni A, et al. Characterization of Shigella strains in Iran by plasmid profile analysis and PCR amplification of ipa genes. J Clin Microbiol 2006;44:2879-83.
- 17. Kosek M, Yori PP, Gilman RH, et al. Facilitated molecular typing of Shigella isolates using ERIC-PCR. Am J Trop Med Hyg 2012;86:1018-25.
- 18. Liang SY, Watanabe H, Terajima J, et al. Multilocus variable-number tandem-repeat analysis for molecular typing of Shigella sonnei. J Clin Microbiol 2007;45:3574-80.
- 19. Tajbakhsh M, Garcia Migura L, Rahbar M, et al. Antimicrobial-resistant Shigella infections from Iran: an overlooked problem? J Antimicrob Chemother 2012;67:1128-33.

 20. Gorge O, Lopez S, Hilaire V, et al. Selection and validation of a multilocus variable-number tandem-repeat analysis panel for typing Shigella spp. J Clin Microbiol 2008;46:1026-36.
- 21. Weniger T, Krawczyk J, Supply P, et al. MIRU-VNTRplus: a web tool for polyphasic genotyping of Mycobacterium tuberculosis complex bacteria. Nucleic Acids Res 2010;38:W326-31.
- 22. von Seidlein L, Kim DR, Ali M, et al. A multicentre study of Shigella diarrhoea in six Asian countries: disease burden, clinical manifestations, and microbiology. PLoS Med 2006;3:e353.
- 23. Angelini M, Stehling EG, Moretti ML, et al. Molecular epidemiology of Shigella spp strains isolated in two different metropolitam areas of southeast Brazil. Braz J Microbiol 2009;40:685-92.
- 24. Ghosh S, Pazhani GP, Chowdhury G, et al. Genetic characteristics and changing antimicrobial resistance among Shigella spp. isolated from hospitalized diarrhoeal patients in Kolkata, India, J Med Microbiol 2011;60:1460-6.
- 25. Jomezadeh N, Babamoradi S, Kalantar E, et al. Isolation and antibiotic susceptibility of Shigella species from stool samples among hospitalized children in Abadan, Iran. Gastroenterol Hepatol Bed Bench 2014;7:218-23.
- 26. MoezArdalan K, Zali MR, Dallal MM, et al. Prevalence and

Genotyping of Shigella spp. by MLVA

pattern of antimicrobial resistance of Shigella species among patients with acute diarrhoea in Karaj, Tehran, Iran. J Health Popul Nutr 2003;21:96-102.

- 27. Wang YW, Watanabe H, Phung DC, et al. Multilocus variable-number tandem repeat analysis for molecular typing and phylogenetic analysis of Shigella flexneri. BMC Microbiol 2009;9:278.
- 28. Qu M, Zhang X, Huang Y, et al. Multilocus variable-number
- tandem-repeat analysis for molecular subtypes of Shigella isolates in Beijing. Zhonghua Yu Fang Yi Xue Za Zhi 2012;46:329-33.
- 29. Pourcel C, Hormigos K, Onteniente L, et al. Improved multiple-locus variable-number tandem-repeat assay for Staphylococcus aureus genotyping, providing a highly informative technique together with strong phylogenetic value. J Clin Microbiol 2009;47:3121-8.