# Mesenchymal stem cells from umbilical cord tissue as potential therapeutics for cardiomyodegenerative diseases – a review

Trixi Hollweck, Christian Hagl, Günther Eissner

Department of Cardiac Surgery, Munich University Medical Center, Munich, Germany.

Heart failure is one of the leading causes of death worldwide. End stage disease often requires heart transplantation, which is hampered by donor organ shortage. Tissue engineering represents a promising alternative approach for cardiac repair. For the generation of artificial heart muscle tissue several cell types, scaffold materials and bioreactor designs are under investigation. In this review, the use of mesenchymal stem cells derived from human umbilical cord tissue (UCMSC) for cardiac tissue engineering will be discussed.

**Key words:** Umbilical cord tissue, mesenchymal stem cells, cardiac differentiation, non-degradable scaffolds, polytetrafluorethylene, titanium, bioreactor

In humans, the heart is one of the least regenerative organs in the body (1). The limited ability of the heart to regenerate damaged tissue after major cardiac injuries often leads to heart failure (2). Despite a wide range of therapeutic approaches, heart failure remains the leading cause of death in modern societies (3,4). Myocardial infarction is the major cause of heart failure. Ischemic conditions result in an irreversible loss of functional cardiomyocytes which are gradually replaced by fibroblasts, forming non-contractile scar tissue (5). Resident cardiac progenitor cells can be found in transplanted human hearts, and evidence of myocyte proliferation in the human heart exists. However, this proliferation does not compensate for up to 1 billion cardiomyocytes being lost after MI (6). In end stage heart failure allogeneic heart transplantation remains the last treatment option, but it is limited due to donor organ shortage. According to the Eurotransplant International Foundation, in 2011 the demand for donor hearts was covered only to 35 % in Germany (7). The generation of artificial heart muscle tissue using cardiac tissue engineering might be a reasonable alternative to heart transplantation.

## Cardiac tissue engineering

Cardiac tissue engineering is an interdisciplinary research area in regenerative medicine. Besides paracrine effects supporting angiogenesis, modulation of extracellular matrix components, and stimulating interactions with resident cardiac progenitor cells, the main aim of

Corresponding author: Department of Cardiac Surgery, Munich University Medical Center, Marchioninistr. 15, 83177 Munich, Germany. Email: Guenther.Eissner@med.uni-muenchen.de

## **Cell application**

The route of administration of autologous and allogeneic cells is one of the central questions in cardiac tissue engineering. Cellular cardiomyoplasty is performed by intracoronary injection or direct implantation of a single cell suspension into the myocardium (12). Animal studies demonstrate an increase in the pumping function of the heart. However, myocardial regeneration was not observed (13). Functional improvement could be explained by secretion and stimulation of angiogenic growth factors resulting in the lack of myogenesis stimulation and contractility improvement (14). Systemic application also carries the risk of pulmonary accumulation of cells. Experimental injection of cells into the infarcted region ensures the delivery to the damaged area but is hampered by significant cell loss (12, 15).

An alternative approach to injection of isolated cells into the heart is the use of artificially engineered tissues that are geometrically, structurally and functionally defined prior to transplantation. Scaffolds are populated *in vitro* with cells and subsequently implanted onto the infarcted zone to allow precise cell delivery and mechanical support (4, 16).

Resident cardiac stem cells can thus be stimulated to migrate into the area of regeneration induced by growth factors released from the implanted cells. Reconstitution of heart muscle tissue would also be possible by implanted cells themselves, differentiated in vivo into cardiomyocytes by local tissue-specific mechanism or differentiated in vitro prior to transplantation. In contrast to cell injection, using artificial heart tissue might results in less cell loss due to cell immobilization scaffolds on by adhesion molecules (17).

For myocardiac regeneration, cells from several cell sources like skeletal muscle (18) or neonatal rat heart (19) have been investigated already. Although some of these cell types integrate into damaged myocardium, application is restricted by limited availability and poor proliferation capacity (20). This has led to the search for alternative more efficient cell populations.

### **Cell sources**

Heart muscle regeneration requires cells with the capability for proliferation, plasticity and functional integration into cardiac tissue (21). Stem cells feature unique regenerative potential and are consequently qualified for this claim (22-24). Due to their origin, stem cells are categorized into embryonic, induced and adult stem cells (25). Embryonic stem cells (ESC) derived from early embryos are well expandable and able to differentiate into various tissues. This pluripotency qualifies them for therapeutic applications, however ethical and legal concerns about using embryos for stem cell isolation exist. Moreover, in animal studies teratocarcinomas are described after implantation of ESC (26, 27). Induced pluripotent stem cells (iPSC) are thought to be an alternative cell source without ethical concerns (28). Since the discovery of genetic reprogramming of adult fibroblasts into pluripotent stem cells (29, 30) extensive efforts aim at the clinical applicability of iPSC, including reprogramming of fibroblasts using recombinant proteins (protein-induced pluripotent stem cells (p-iPSC)) (31).

Potential cardiomyogenic differentiation of p-iPSC also offers an option for cardiac tissue engineering. Other sources include hematopoetic stem cells and mesenchymal stem cells, either obtained from newborn, children or adults. They are collectively termed adult stem cells or postnatal stem cells, particulary if they are derived from infantile organisms (32). Mesenchymal stem cells (MSC) have the capability for self-renewal and differentiation into various lineages of

mesenchymal origin, nerve and myogenic cells. Besides a comparable differentiation capacity, MSC seem to be more efficacious in tissue reconstitution than adult hematopoetic stem cells, due to strong pro-angiogenic properties necessary for a functional myocardium (33).

Moreover, MSC show a higher homing potential towards tissue defects resulting in the production of repairing growth factors (34, 35). Since they have the ability to differentiate into cardiomyocyte (36), MSC are a potential cellular source for cardiac stem cell-based therapy (35, 37). MSC have been already tested clinically and do not raise any ethical concerns (38). To date, human bone marrow (BM) represents the major source of MSC. However, aspirating BM from the patient is an invasive procedure and the number as well as the differentiation potential and the maximum life span of human BM-derived MSC (BMMSC) significantly decline with donor age (39, 40).

The umbilical cord tissue may be an attractive alternative source to BM (41). The two arteries and the single vein of the umbilical cord with a length up to 60 cm are surrounded with fetal connective tissue – the so called Wharton's jelly, protecting the vessels against compression, torsion and bending (42). In line with several publications, Weiss et al. described MSC derived from the perivascular and intervascular region of the umbilical cord tissues (fig. 1), collectively termed umbilical cord tissue derived mesenchymal stem cells (UCMSC) (43-46).

In contrast to BMMSC, UCMSC are easily attainable and can extensively be expanded and maintained in culture, even after cryopreservation (43, 47-49). With regard to future clinical trials, our group successfully managed to grow UCMSC under GMP-compliant culture conditions, while retaining their phenotypic and functional properties (50). Due to close relation to the fetal phase, it is assumed that UCMSC are less determined than adult stem cells, show less teratogenic potential and are free of viruses (43, 51). In addition, UCMSC qualify for an allogeneic use due to their immunological naivity (51) and weaker response to inflammatory stimuli. With regard to their multipotency (32, 41, 51-52) UCMSC can be differentiated into bone, cartilage, neural and muscle cells as well as cardiomyocyte-like cells, as they express cardiac troponin-I and *N*-cadherin (32, 41, 43, 51-53).



## **Cardiomyogenic differentiation**

Cardiomyogenic differentiation is presumed to be triggered by an increased expression of the embryonic transcription factor GATA-4 (54). GATA-4 proteins are not only important for heart development, but also constitute one of the earliest cardiac markers (55). In adult hearts, GATA-4 regulates the expressions of several sarcomeric proteins, which are used in combination with proteins of the troponin complex for the verification of induced cardiomyocytes (56). In addition, electric coupling by gap junctional connexins is essential for contraction (57). Contraction of cardiomyogenically differentiated stem cells of embryonic and adult origin has already been described, however with a percentage of contracting cells less than 10 % (58-60).

Cardiomyogenic differentiation of BM-MSC and a significant improvement of left ventricular

function after application of BM-MSC have also been published (61-63). However, the broad differentiation potential of MSC could also lead to undesirable effects. Since undifferentiated MSC tend to spontaneously differentiate into multiple lineages when transplanted in vivo, it is possible that such uncommitted stem cells undergo maldifferentiation within the infracted myocardium with potentially life-threatening consequences, e.g. osteogenic differentiation of BM-MSC within ischemic myocardium in a murine model (64). Although such phenomenons are not yet described for UCMSC (49) it was postulated that a certain cardiac differentiation of stem cells prior to transplantation would result in enhanced myocardial regeneration and recovery of heart function (65).

In this context, initiating the transformation of stem cells into a cardiomyogenic lineage is accomplished by defined culture conditions (66). Embryo-like aggregates (67), the DNA demethylating agent 5-azacytidine (53, 68-69), several growth factors and the oxytocic hormone (70) are used to induce myocyte differentiation of various stem cell types. Maltsev et al. demonstrated the expression of cardio-specific genes, proteins and action potentials in cells differentiated from murine embryonal stem cells by cultivation in hanging drops as "embryoid bodies" (67). Using this differentiation system, UCMSC form aggregates, but cellular outspread is not sufficient for performing extensive analyses. Failure of cellular outgrowth may be explainable due to the dependence of this method on the initial cell number present in the aggregates (71).

Based on a yet unknown mechanism, cytostatic 5-azacytidine results in cardiac differentiation of stem cells by DNA-demethylation (72). Cardiac differentiation of MSC induced by 5azacytidine is controversially discussed. Martin-Rendon and colleagues report that 5-azacytidin treated human MSC derived from umbilical cord and bone marrow do not generate cardiomyocytes in vitro at high frequencies (23).

In contrast, results of Antonitsis et al. and Pereira et al. indicate that adult human bone marrow MSC (73) and MSC from umbilical cord (35) can differentiate towards a cardiomyogenic lineage after 5-azacytidine treatment. These discrepancies might be explained by the variability in culture conditions (74) or by different specification criteria for what makes a cell a cardiomyocyte. For example, the use of cytokines and growth factors is a step forward in the development of a defined culture milieu for directing the cardiomyogenic differentiation. In this context, TGF- $\beta$  and bFGF are the most important growth factors in embryonic cardiac development affecting cell proliferation, migration and differentiation (75). Xu et al. stated that bFGF is necessary during the differentiation process because of its capability to develop the myogenic phenotype and promote the formation of myotubes (76).

However, our study showed that UCMSC exposed to 5-azacytidine convert into cells changing their morphology and expressing cardiac-specific proteins irrespective of the presence of bFGF (71).

UCMSC differentiated according to Wu et al. (69), using 5 µM 5-azacytidine for 24 h and bFGF containing culture medium, increase in size with striate pattern and express cardiac actin, cardiac actinin, sarcomeric actin, sarcomeric actinin, myosin heavy chain as well as connexin 43 after 5 weeks of culture. UCMSC treated with 3 µM 5-azacytidine for 24 h according to Wang et al. (53) and 10 µM 5-azacytidine for 72 h according to Matsuura protocol I (70) also change their morphology and express these cardiac specific proteins known for regulating contraction and gapjunctional communication without supplemented bFGF. TGF-β1 in combination with 5-azacytidine have been found to promote differentiation of human cardiomyocyte progenitor cells (68).

However, in our hands, the combination of 5-azacytidine and TGF- $\beta$ 1 stimulation of UCMSC leads to a flattened appearance and the expression of cardiac actin, cardiac actinin, sarcomeric actin, sarcomeric actinin as well as connexin 43 after 5 weeks of culture, but UCMSC do not express any troponins or myosins necessary for contraction. In addition, during the differentiation process, cell numbers decreased to levels insufficient for immunocytochemical analyses (71).

Cardiac differentiation of embryonic P19 carcinoma cells and adult Sca-1+ cells of murine heart is also described after exposure to oxytocin, the mechanism of action, however, is unknown (70-77). Oxytocin, a female reproductive hormone, is necessary for uterine contractions during ovulation and parturition. The expression levels of oxytocin are higher in developing hearts than in adult hearts suggesting that oxytocin may be involved in cardiomyocyte differentiation (78). Data from Matsuura *et al.* indicate that oxytocin is a more potent inducer of cardiac differentiation of Sca-1+ adult murine heart cells than 5-azacytidine (70). This is supported by our results figure 2, demonstrating that human UCMSC exposed to 10nM oxytocin for 72 h express the cardiomyocyteassociated proteins including cardiac actin (fig. 2a), sarcomeric actin (fig. 2b), cardiac troponin T (fig. 2d), connexin 43 (fig. 2e) cardiac actinin (fig. 2f), sarcomeric actinin (fig. 2g), and myosin heavy chain (fig. 2h) in significantly higher frequencies than after 5-azacytidine treatment (71). This analysis revealed that UCMSC can be differentiated into cardiomyocyte-like cells, however, functional analyses of oxytocin-differentiated UCMSC, such as to monitor action potentials, have yet to be performed.

#### Scaffolds

It is known that isolated cells are generally not able to form new tissue autonomously (79). For generating tissue *in vitro*, cells have to be colonized onto natural or artificial scaffolds. However, development of functional tissue requires an optimal interaction of cells and scaffolds. A scaffold should provide chemical stability and physical properties matching the surrounding tissue to provide cytocompatibility, support adhesion, proliferation, and mechanical strength (80). Additionally, scaffolds are required to retain cell phenotype and ensure protein synthesis (81).



**Fig 2.** Immunocytochemical analysis of UCMSC differentiated according to Matsuura et al. using oxytocin (70). Cardiac differentiated UCMSC express the contractile proteins cardiac actin (a), sarcomeric actin (b), cardiac troponin T (c), cardiac actinin (d), sarcomeric actinin (e), myosin heavy chain (f) as well as the gap junctional protein connexin 43 (g) (a-g; all in green fluorescence) for electrical cell-to-cell coupling. Cell nuclei were stained by DAPI (a-g; blue) (71). Scale bars:  $a-f = 50 \mu m$ ,  $g = 25 \mu m$ .

In order to avoid implant rejection and inflammatory response, scaffolds should be biocompatible and sterilizable. For generating functional heart muscle, scaffolds should be made of flexible and tear-resistant material to allow contraction (80, 82). Regardless of which biocompatible material scaffolds are made of, the microarchitecture including porosity, pore geometry and the surface micro-texture considerably influence cell function (83). Both, scaffolds of biological origin and those made of synthetic material as well as some type of hybrids are currently under investigation for tissue engineering applications (84-86).

## **Biomaterials**

## Decellularised tissues

These tissues of allogeneic or xenogeneic origin are derived from enzymatic or detergent decellularisation (87). Cell-free tissues consist of natural extracellular matrix (ECM), degrade after implantation and are replaced by ECM-proteins of transplanted cells after re-seeding or by in-growing cells (88). Due to excellent mechanical properties, decellularised tissues are used for the development of viable heart valve prostheses (89). However, decellularisation can damage scaffold tissue, resulting in a decreased tensile strength and elasticity. Xenogenic decelluarised tissues undergo aneurysms and lead to infections and thrombosis. In addition, decellularisation might affect seeding efficiency due to residual antigenic components inducing humoral responses (87, 90, 91).

# Biopolymers

Natural polymers include fibrin, collagen, chitin, hyaluronic acid, and alginates. Besides enzymatic degradation, biopolymers show low inflammatory activity and toxicity (81). In addition, they support cell growth on implants due to their high protein content and accelerate healing because of strong adhesion to recipient organs (58). Fibrin is part of the blood clotting system and plays a central role in wound healing. As an alternative to conventional surgical sutures, fibrin glue is clinically used for wound closure (90). Collagen is another example of biopolymers in clinical practice. It is the predominant protein in the human body and the main component of ECM (81, 90). In cardiovascular surgery, collagen is used for heart valve replacement and blood vessel substitutes as well as for bone repair and burn and ulcer treatment (82). Naturally occurring biomaterials may most closely simulate the native cellular milieu, but large batch-to-batch variations upon isolation from biological tissues and poor mechanical strength are the main limitations for a clinical application. In addition, biopolymers are often denatured in a way no longer enabling tissue formation and often require chemical modifications, which can lead to toxicity (79, 81).

## Synthetic materials

## Degradable Polymers

Numerous synthetic and degradable polymers like poly( $\alpha$ -hydroxy ester), particulary polyglycolic acid (PGA) or polylactid acid (PLA), polyanhydrides, polyorthoester and polyphosphazanes have been developed to overcome the limitations of natural materials mentioned above (79). Synthetic degradable polymers undergo degradation during cell culture or after implantation upon formation of tissue specific neo-ECM. Most of these polymers are resistant to enzymatic digestion, they are rather chemically hydrolyzed resulting in consistent and patientindependent degradation (79, 90).

In order to allow tissue generation and remodeling processes, microstructure, mechanical properties and resorption rate can be regulated by porosity and pore size, for example. After cell seeding, synthetic degradable polymers initially retain the cellular compound and ensure mechanical function of implants until an ECM is formed by colonized cells (90). However, if polymers degrade

faster than the development of an ECM occurs, seeded cells lose their connectivity resulting in final cell loss and an inhibition of therapeutic effects. In addition, polyesters release degradation products which affect biocompatibility. Moreover, polyesters are stiff materials suitable for load-bearing implants, but the minimal flexibility precludes their use for soft tissues like heart muscle.

# Non-degradable polymers

Synthetic, non-degradable polymeres are characterized by structural resistance, a special topography and a three-dimensional form with defined pore sizes. Polyethylen terepthalate (PET, Dacron®), polyurethane (PU), and expanded polytetrafluorethylene (ePTFE) dominate the graft market (92) because of their anti-thrombotic properties. PET is a semi-crystalline aromatic polyester built of woven or knitted multiple fibres. While woven PET grafts feature small pores, knitted implants exhibit larger pores supporting tissue ingrowth. PET is mainly used for artificial blood vessels, tendon substitutes and surgical sutures (82, 86).

PU is a polymer with a characteristic urethane group. Within the monomeric unit, moieties could be substituted by different groups, resulting in versatile properties. Fabrication of hydrolytic stable PU led to the development of different implants like vascular grafts, artificial heart valves and catheters (82, 86). PTFE is an unbranched linear polymere built of fluorine and carbon. Expanded PTFE (ePTFE) features nodes in fibrillar structure with longitudinal internodal distances of 17-90  $\mu$ m. Due to the symmetrical design of the monomeric unit, crystallinity of ePTFE come up to 94 % preventing degradation. Besides easy availability, ePTFE offer nonimmunogenic and anti-thrombotic properties (86).

ePTFE is clinically used for cardiac, groin and vascular grafts. However, the use of synthetic non-degradable polymers as scaffolds for tissue engineering is often limited by the poor retention of cells to these hydrophobic biomaterials (82, 93-95, 96). Yu et al. report, that endothelial cells – in contrast to smooth muscle cells - adhere poorly to ePTFE. In our experiments, the investigation of adherence, viability, proliferation and morphology of UCMSC on uncoated ePTFE scaffolds showed poor results (97). This is in line with results from Neuss et al. (80), demonstrating that bone-marrow derived MSC (BMMSC) display a round, spherical morphology on ePTFE. Furthermore, ePTFE does not allow BMMSC proliferation, indicating that cells need an underlying matrix providing them with sufficient binding sites.

### Hybrids

Hybrids combine advantages of different materials in one composite. Surface modification of non-degradable polymers increases the wettability leading to an improved seeding efficiency. The knowledge that positively charged surfaces are more conductive to cell adhesion and morphological maturation, led to the examination of various adhesive coatings of synthetic materials (98). The matrix molecules of these coating substances, such as albumin, collagen, fibrin, gelatine, fibronectin, laminin and fibrin glue, bind directly to specific domains on the cell membrane (95, 99). For example, Kaehler et al. and Feugier et al. pre coated ePTFE vascular prostheses with fibronectin-treated Type I/III collagen and reported a higher cell adherence and spreading on these grafts (100, 101). Although coating with specific proteins improves cell adhesion, the integrity of the coating is compromised by mechanical stress (94). Furthermore, if surfaces are not completely endothelialized or endothelial cells are lost upon exposure to mechanical loadings, these coatings attract platelets. In these cases, the technique leads to a more thrombogenic surface, which defeats the purpose of cell seeding (102).

To overcome the limitation of biological coatings, a hydrophilic titanium-coated surface can

be obtained by plasma-assisted chemical vapour deposition (PACVD): The resulting covalent bonding can only be separated by destroying the synthetic structure itself. The titanium layer is extremely thin and has the same flexibility as the synthetic material. Titanium-coated synthetics feature outstanding wettability enabling them to adapt to the anatomical environment and to enhance cell adhesion (92). Moreover, titanium-coated synthetic implants provide excellent biocompatibility because of the oxide layer which forms under atmospheric conditions. Therefore titanium and its alloys are widely used as biomaterials in association with tissue, bone, and blood (103-105). Our findings, that titanium-coated ePTFE figure 3a, scaffolds are superior to uncoated ePTFE scaffolds figure 3b, in UCMSC adherence, viability and proliferation (97) are in line with results of previous studies, demonstrating a support of MSC adhesion and proliferation on titanium dishes (104, 106).

#### **Bioreactors**

For the use of tissue engineered constructs *in vivo*, it is essential to examine their functionality and mechanical integrity prior to implantation (107). In addition, forces acting directly or indirectly on cells, e.g. via scaffolds, can affect cellular differentiation (82). *In vivo*, cells are stimulated continuously by mechanical, electrical and chemical signals influencing their phenotype,

morphology and proliferation. If these signals are inappropriate or absent, cells lose their ability to form organized tissues (108). Thus, bioreactors simulating physiological conditions, such as mechanical shear stress, play a crucial role in the development of tissue engineered constructs (107).

The development of an effective bioreactor requires the consideration of various parameters. Ideally, bioreactors allow the regulation of physical parameters such as temperature, pH, pO<sub>2</sub>, pCO<sub>2</sub>, allow nutrient supply and removal of toxic metabolites as well as mechanical stimuli. Moreover, the material must be compatible with the manufacturing process, sterilization technique and the cultured cell type (109). Bioreactors can be applied for cell seeding, cultivation of colonized scaffolds and for conditioning of functional tissue engineered prostheses (110-112).

In heart valve fabrication, bioreactors for tissue formation under dynamic culture conditions were demonstrated several times (107, 113, 114). Bioreactors also support tissue formation of heart muscle in vitro (115, 116). An effective approach to improve the contractile properties of artificial heart muscle constructs is electrical field stimulation or mechanical stimulation by unidirectional or auxotonic stretching (117). Accompanied by an improvement of contractile function, some studies extracellular demonstrated matrix formation increased cell proliferation and uniform cell distribution of strained constructs (118,119).



**Fig 3.** Morphology of UCMSC seeded on uncoated and titanium-coated ePTFE. UCMSC display their characteristic spindle-shaped morphology in a homogenous coverage on titanium-coated ePTFE (a, arrow) in contrast to a spherical morphology seeded on uncoated ePTFE (b, arrow) (97). Scale bars:  $a = 10 \mu m$ ,  $e = 50 \mu m$ .

In this context, Zimmermann et al. reported from highly differentiated cardiac tissue constructs after cyclic mechanostimulation in a stretch device (58). Sodian et al. developed a closed-looped perfused bioreactor by combining pulsatile perfusion and periodically stretching of tissueengineered patch constructs (120). Birla et al. described a bioreactor system that applies electromechanical stretch to bioengineered heart muscle constructs with no evidence of physical damage (121). In order to repopulate ischemic myocardium with cells that might restore contractility, we analyzed the stability of the cellular coating upon mechanical stress in a newly developed bioreactor figure 4, mimicking myocardial contraction (122).

Three fluid compartments enable the comparison of different media, cells and scaffolds at defined mechanical loadings. Manufacturing of the core unit from acrylic glass provides optical transparency for macroscopical observation of processes within the unit. Elements of acrylic glass, stainless steel, Teflon® and silicone are robust and can be gas sterilized. Fixing of seeded scaffolds by clip-systems allows an easy assembling, reliable fixing and facilitates sterile handling. The speed controlled gear motor provides frequencies at 1-65

Hz, offering gradually increasing mechanical loadings of the tissue-engineered scaffolds. In addition, the bioreactor was designed in a dimension that allows its operation in a standard incubator. Preliminary experiments with UCMSCseeded ePTFE scaffolds show the mechanical integrity of the cellular coating after friction stress generated in the pulsatile bioreactor. Viability and ultrastructural morphology of the stem cells are also maintained upon mechanical stress (122).

#### Conclusion

Cardiac tissue engineering using UCMSC represents a promising approach for the repair of the injured heart, however, clinical relevance of tissue engineered constructs have to be evaluated in vivo. Functional regeneration of heart tissue after cardiomyodegenerative diseases should he demonstrated by the integration of UCMSC-seeded implants and/or their interaction with resident cardiac stem cells. In addition, survival of implanted constructs, tissue-specific differentiation and vascularization have to be verified. Moreover, electric integration resulting in functional reconstitution of the injured muscle tissue is a key step in the evaluation of safety and efficiency of UCMSC-seeded implants.



Fig 4. Pulsatile bioreactor. Comparison of different media, cells and scaffolds are enabled by three media compartments at defined mechanical loadings. Core unit manufacturing from acrylic glass provides optical transparency for macroscopical observation of processes within the unit (122). Scale bar = 20 mm.

#### Acknowledgement

The research of the authors was supported by funds from the Else Kröner-Fresenius-Foundation and the Elisabeth und Rudolf Hirsch-Foundation for Medical Research. The authors wish to thank Dr.-Ing. Bassil Akra from our institution for helpful discussions.

### References

1. Capi O, Gepstein L. Myocardial regeneration strategies using human embryonic stem cell-derived cardiomyocytes. J Control Release 2006;116:211-8.

2. Wu KH, Zhou B, Yu CT, et al. Therapeutic potential of human umbilical cord derived stem cells in a rat myocardial infarction model. Ann Thorac Surg 2007;83:1491-500.

3. Tousoulis D, Briasoulis A, Antoniades C, et al. Heart regeneration: what cells to use and how? Curr Opin Pharmacol 2008;8:211-8.

4. Zimmermann WH, Eschenhagen T. Cardiac tissue engineering for replacement therapy. Heart Fail Rev 2003;8:259-69.

5. Kadivar M, Khatami S, Mortazavi Y, et al. In vitro cardiomyogenic potential of human umbilical vein-derived mesenchymal stem cells. Biochem Bioph Res Co 2006;340: 639-47.

6. Collins SD, Baffour R, Waksman R. Cell therapy in myocardial infarction. Cardiovasc Revasc Med 2007;8:43-51.

7. Eurotransplant International Foundation. Available at: http://www.eurotransplant org/cms/index php?page=yearlystats; Accessed February 29, 2012.

8. Marelli D, Desrosiers C, el-Alfy M, et al. Cell transplantation for myocardial repair: an experimental approach. Cell Transplant 1992;1:383-90.

9. Reinecke H, Zhang M, Bartosek T, et al. Survival, integration, and differentiation of cardiomyocyte grafts: a study in normal and injured rat hearts. Circulation 1999;100:193-202.

10. Stamm C, Westphal B, Kleine HD, et al. Autologous bonemarrow stem-cell transplantation for myocardial regeneration. Lancet 2003;361:45-6.

11. Stamm C, Nasseri B, Choi YH, et al. Cell Therapy for Heart Disease: Great Expectations, As Yet Unmet. Heart Lung Circ 2009;18:245-56. 12. Jawad H, Ali NN, Lyon AR, et al. Myocardial tissue engineering: a review. J Tissue Eng Regen Med 2007;1:327-42.

13. Orlic D, Kajstura J, Chimenti S, et al. Bone marrow stem cells regenerate infarcted myocardium. Pediatr Transplant 2003;7 Suppl 3:86-8.

14. Caplan AI, Dennis JE. Mesenchymal stem cells as trophic mediators. J Cell Biochem 2006;98:1076-84.

15. Blocklet D, Toungouz M, Berkenboom G, et al. Myocardial homing of nonmobilized peripheral-blood CD34(+) cells after intracoronary injection. Stem Cells 2006;24:333-6.

 Christman KL, Lee RJ. Biomaterials for the treatment of myocardial infarction. J Am Coll Cardiol 2006;48:907-13.

17. Jawad H, Lyon AR, Harding SE, et al. Myocardial tissue engineering. Brit Med Bull 2008;87:31-47.

 Taylor DA, Atkins BZ, Hungspreugs P, et al. Regenerating functional myocardium: Improved performance after skeletal myoblast transplantation. Nat Med 1998;4:929-33.

19. Zimmermann WH, Fink C, Kralisch D, et al. Threedimensional engineered heart tissue from neonatal rat cardiac myocytes. Biotechnol Bioeng 2000;68:106-14.

20. Kofidis T, Akhyari P, Wachsmann B, et al. A novel bioartificial myocardial tissue and its prospective use in cardiac surgery (Retracted article. See vol. 40, pg. 1554, 2011). Eur J Cardio-Thorac 2002;22:238-43.

 Zimmermann WH, Didie M, Doker S, et al. Heart muscle engineering: An update on cardiac muscle replacement therapy. Cardiovasc Res 2006;71:419-29.

22. Alison MR, Poulsom R, Forbes S, et al. An introduction to stem cells. J Pathol 2002;197:419-23.

23. Martin-Rendon E, Sweeney D, Lu F, et al. 5-Azacytidinetreated human mesenchymal stem/progenitor cells derived from umbilical cord, cord blood and bone marrow do not generate cardiomyocytes in vitro at high frequencies. Vox Sang 2008;95:137-48.

24. Spangrude GJ. When is a stem cell really a stem cell? Bone Marrow Transplant 2003;32 Suppl 1:S7-11.

 Wormer E. Stammzellen - Mehr Wissen? Köln: Lingen Verlag GmbH & Co.KG; 2003.

26. Dawson E, Mapili G, Erickson K, et al. Biomaterials for stem cell differentiation. Adv Drug Deliv Rev 2008;60:215-28.

27. Tondeur S, Assou S, Nadal L, et al. [Biology and potential of human embryonic stem cells]. Ann Biol Clin (Paris) 2008;66:241-7.

28. Feng B, Jiang JM, Kraus P, et al. Reprogramming of fibroblasts into induced pluripotent stem cells with orphan nuclear receptor Esrrb. Nat Cell Biol 2009;11:197-203.

29. Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. Cell 2006;126:663-76.

30. Takahashi K, Tanabe K, Ohnuki M, et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. Cell 2007;131:861-72.

31. Zhou HY, Wu SL, Joo JY, et al. Generation of Induced Pluripotent Stem Cells Using Recombinant Proteins. Cell Stem Cell 2009;4:381-4.

32. Aigner J, Eblenkamp M, Wintermantel E. Techniques for functional tissue and organ replacement using postnatal stem cells. Chirurg 2005;76:435-44.

33. Pittenger MF, Martin BJ. Mesenchymal stem cells and their potential as cardiac therapeutics. Circ Res 2004;95:9-20.

34. Kinnaird T, Stabile E, Burnett MS, et al. Local delivery of marrow-derived stromal cells augments collateral perfusion through paracrine mechanisms. Circulation 2004;109:1543-9.

35. Pereira WC, Khushnooma I, Madkaikar M, et al. Reproducible methodology for the isolation of mesenchymal stem cells from human umbilical cord and its potential for cardiomyocyte generation. J Tissue Eng Regen M 2008;2:394-9.

36. Shake JG, Gruber PJ, Baumgartner WA, et al. Mesenchymal stem cell implantation in a swine myocardial infarct model: Engraftment and functional effects. Ann Thorac Surg 2002;73:1919-25.

37. Lakshmipathy U, Verfaillie C. Stem cell plasticity. Blood Rev 2005;19:29-38.

38. Hida N, Nishiyama N, Miyoshi S, et al. Novel cardiac precursor-like cells from human menstrual blood-derived mesenchymal cells. Stem Cells 2008;26:1695-704.

39. Sethe S, Scutt A, Stolzing A. Aging of mesenchymal stem cells. Ageing Res Rev 2006;5:91-116.

40. Wang M, Yang Y, Yang DM, et al. The immunomodulatory activity of human umbilical cord blood-derived mesenchymal stem cells in vitro. Immunology 2009;126:220-32.

41. Eblenkamp M, Aigner J, Hintermair J, et al. [Umbilical cord stromal cells (UCSC). Cells featuring osteogenic differentiation potential]. Orthopade 2004;33:1338-45.

42. Nanaev AK, Kohnen G, Milovanov AP, et al. Stromal differentiation and architecture of the human umbilical cord. Placenta 1997;18:53-64.

43. Mitchell KE, Weiss ML, Mitchell BM, et al. Matrix cells from Wharton's jelly form neurons and glia. Stem Cells 2003;21:50-60.

44. Romanov YA, Svintsitskaya VA, Smirnov VN. Searching for alternative sources of postnatal human mesenchymal stem cells: Candidate MSC-like cells from umbilical cord. Stem Cells 2003;21:105-10.

45. Weiss ML, Troyer DL. Stem cells in the umbilical cord. Stem Cell Rev 2006;2:155-62.

46. Weiss ML, Medicetty S, Bledsoe AR, et al. Human umbilical cord matrix stem cells: Preliminary characterization and effect of transplantation in a rodent model of Parkinson's disease. Stem Cells 2006;24:781-92.

47. Fan CG, Zhang QJ, Zhou JR. Therapeutic Potentials of Mesenchymal Stem Cells Derived from Human Umbilical Cord. Stem Cell Rev Rep 2011;7:195-207.

 Seshareddy K, Troyer D, Weiss ML. Method to isolate mesenchymal-like cells from Wharton's Jelly of umbilical cord. Methods Cell Biol 2008;86:101-19.

49. Troyer DL, Weiss ML. Wharton's jelly-derived cells are a primitive stromal cell population. Stem Cells 2008;26:591-9.

50. Hartmann I, Hollweck T, Haffner S, et al. Umbilical cord tissue-derived mesenchymal stem cells grow best under GMP-compliant culture conditions and maintain their phenotypic and functional properties. J Immunol Methods 2010;363:80-9.

51. Weiss ML, Anderson C, Medicetty S, et al. Immune Properties of Human Umbilical Cord Wharton's Jelly-Derived Cells. Stem Cells 2008;26:2865-74.

52. Kestendjieva S, Kyurkchiev D, Tsvetkova G, et al. Characterization of mesenchymal stem cells isolated from the human umbilical cord. Cell Biol Int 2008;32:724-32.

53. Wang HS, Hung SC, Peng ST, et al. Mesenchymal stem cells in the Wharton's jelly of the human umbilical cord. Stem Cells 2004;22:1330-7. 54. Thurisch B. Untersuchung der Funktion des Transkriptionsfaktors GATA-4 durch eine Mausmutante mit einem induzierbaren RNA-Interferenz System. Dissertation: Humboldt-Universität Berlin; 1979.

55. Xu CH, Police S, Rao N, et al. Characterization and enrichment of cardiomyocytes derived from human embryonic stem cells. Circ Res 2002;91:501-8.

56. Charron F, Tsimiklis G, Areand M, et al. Tissue-specific GATA factors are transcriptional effectors of the small GTPase RhoA. Gene Dev 2001;15:2702-19.

57. Stahlhofen S. Lokalisation, Orientierung und Funktion von Connexin 43 in Kardiomyozyten-Mitochondrien. Dissertation: Universität Duisburg-Essen; 2010.

58. Zimmermann WH, Schneiderbanger K, Schubert P, et al. Tissue engineering of a differentiated cardiac muscle construct. Circ Res 2002;90:223-30.

59. Boheler KR, Czyz J, Tweedie D, et al. Differentiation of pluripotent embryonic stem cells into cardiomyocytes. Circ Res 2002;91:189-201.

60. Yankelson L, Feld Y, Bressler-Stramer T, et al. Cell therapy for modification of the myocardial electrophysiological substrate. Circulation 2008;117:720-31.

61. Fraser JK, Schreiber RE, Zuk PA, et al. Adult stem cell therapy for the heart. Int J Biochem Cell B 2004;36:658-66.

62. Schachinger V, Erbs S, Elsasser A, et al. Intracoronary bone marrow-derived progenitor cells in acute myocardial infarction. New Engl J Med 2006;355:1210-21.

63. Lunde K, Solheim S, Aakhus S, et al. Intracoronary injection of mononuclear bone marrow cells in acute myocardial infarction. New Engl J Med 2006;355:1199-209.

64. Breitbach M, Bostani T, Roell W, et al. Potential risks of bone marrow cell transplantation into infarcted hearts. Blood 2007;110:1362-9.

65. Mastitskaya S, Denecke B. Human spongiosa mesenchymal stem cells fail to generate cardiomyocytes in vitro. J Negat Results Biomed 2009;8:11.

66. Kocaefe C, Balci D, Hayta BB, et al. Reprogramming of Human Umbilical Cord Stromal Mesenchymal Stem Cells for Myogenic Differentiation and Muscle Repair. Stem Cell Rev Rep 2010;6:512-22.

67. Maltsev VA, Rohwedel J, Hescheler J, et al. Embryonic Stem-Cells Differentiate in-Vitro into Cardiomyocytes Representing Sinusnodal, Atrial and Ventricular Cell-Types. Mech Develop 1993;44:41-50.

68. Goumans MJ, de Boer TP, Smits AM, et al. TGF-beta1 induces efficient differentiation of human cardiomyocyte progenitor cells into functional cardiomyocytes in vitro. Stem Cell Res 2007;1:138-49.

69. Wu KH, Mo XM, Zhou B, et al. Cardiac potential of stem cells from whole human umbilical cord tissue. J Cell Biochem 2009;107:926-32.

70. Matsuura K, Nagai T, Nishigaki N, et al. Adult cardiac Sca-1-positive cells differentiate into beating cardiomyocytes. J Biol Chem 2004;279:11384-91.

71. Hollweck T, Hartmann I, Eblenkamp M, et al. Cardiac Differentiation of Human Wharton's Jelly Stem Cells – Experimental Comparison of Protocols Open Tissue Eng Regen Med J 2011;4:95-102.

72. Fukuda K. Development of regenerative cardiomyocytes from mesenchymal stem cells for cardiovascular tissue engineering. Artif Organs 2001;25:187-93.

73. Antonitsis P, Ioannidou-Papagiannaki E, Kaidoglou A, et al. In vitro cardiomyogenic differentiation of adult human bone marrow mesenchymal stem cells. The role of 5-azacytidine. Interact Cardiovasc Thorac Surg 2007;6:593-7.

 Bruinink A, Wintermantel E. Grooves affect primary bone marrow but not osteoblastic MC3T3-E1 cell cultures. Biomaterials 2001;22:2465-73.

75. Heng BC, Haider HK, Sim EKW, et al. Strategies for directing the differentiation of stem cells into the cardiomyogenic lineage in vitro. Cardiovasc Res 2004;62:34-42.
76. Xu WR, Zhang XR, Qian H, et al. Mesenchymal stem cells

from adult human bone marrow differentiate into a cardiomyocyte phenotype in vitro. Exp Biol Med 2004;229: 623-31.

77. Paquin J, Danalache BA, Jankowski M, et al. Oxytocin induces differentiation of P19 embryonic stem cells to cardiomyocytes. Proc Natl Acad Sci U S A 2002;99:9550-5.

78. Fathi F, Murasawa S, Hasegawa S, et al. Cardiac differentiation of P19CL6 cells by oxytocin. Int J Cardiol 2009;134:75-81.

79. Yang SF, Leong KF, Du ZH, et al. The design of scaffolds for use in tissue engineering. Part II. Rapid prototyping techniques. Tissue Eng 2002;8:1-11.

80. Neuss S, Apel C, Buttler P, et al. Assessment of stem cell/biomaterial combinations for stem cell-based tissue engineering. Biomaterials 2008;29:302-13.

81. Vats A, Tolley NS, Polak JM, et al. Scaffolds and biomaterials for tissue engineering: a review of clinical applications. Clin Otolaryngol Allied Sci 2003;28:165-72.

82. Wintermantel E, Ha SW. Medizintechnik Life Science Engineering. Berlin, Heidelberg, New York: Springer Verlag GmbH; 2008.

83. Mata A, Kim EJ, Boehm CA, et al. A three-dimensional scaffold with precise micro-architecture and surface micro-textures. Biomaterials 2009;30:4610-7.

 Ahsan T, Nerem RM. Bioengineered tissues: the science, the technology, and the industry. Orthod Craniofac Res 2005;8:134-40.

 Minuth WW, Strehl R. Zukunftstechnologie Tissue Engineering Weinheim, Berlin, Darmstadt Zürich: Wiley-VCH Verlag GmbH & Co.KGaA; 2003.

86. Wintermantel E, Ha SW. Medizintechnik mit biokompatiblen Werkstoffen und Verfahren

Berlin, Heidelberg, New York: Springer Verlag GmbH; 2002.

87. Stock UA, Vacanti JP. Tissue engineering: current state and prospects. Annu Rev Med 2001;52:443-51.

 Atala A. Engineering organs. Curr Opin Biotechnol 2009;20:575-92.

89. Stamm C, Steinhoff G. [Cardiac tissue engineering]. Herz 2002;27:445-52.

90. Isenberg BC, Williams C, Tranquillo RT. Small-diameter artificial arteries engineered in vitro. Circ Res 2006;98:25-35.

91. Borschel GH, Huang YC, Calve S, et al. Tissue engineering of recellularized small-diameter vascular grafts. Tissue Eng 2005;11:778-86.

92. Perea H. Magnetic vascular engineering: development and validation. Aachen: Shaker Verlag GmbH; 2008.

93. Gulbins H, Dauner M, Petzold R, et al. Development of an artificial vessel lined with human vascular cells. J Thorac Cardiovasc Surg 2004;128:372-7.

94. Kannan RY, Salacinski HJ, Butler PE, et al. Current status of prosthetic bypass grafts: a review. J Biomed Mater Res B Appl Biomater 2005;74:570-81.

95. Vara DS, Salacinski HJ, Kannan RY, et al. Cardiovascular tissue engineering: state of the art. Pathol Biol (Paris) 2005;53:599-612.

96. Yu H, Dai W, Yang Z, et al. Smooth muscle cells improve endothelial cell retention on polytetrafluoroethylene grafts in vivo. J Vasc Surg 2003;38:557-63.

97. Hollweck T, Marschmann M, Hartmann I, et al. Comparative analysis of adherence, viability, proliferation and morphology of umbilical cord tissue-derived mesenchymal stem cells seeded on different titanium-coated expanded polytetrafluoroethylene scaffolds. Biomed Mater 2010;5:065004.

98. Tiwari A, Salacinski HJ, Hamilton G, et al. Tissue engineering of vascular bypass grafts: role of endothelial cell extraction. Eur J Vasc Endovasc Surg 2001;21:193-201.

99. Carnagey J, Hern-Anderson D, Ranieri J, et al. Rapid endothelialization of PhotoFix natural biomaterial vascular grafts. J Biomed Mater Res B Appl Biomater 2003;65:171-9.

100. Kaehler J, Zilla P, Fasol R, et al. Precoating substrate and surface configuration determine adherence and spreading of seeded endothelial cells on polytetrafluoroethylene grafts. J Vasc Surg 1989;9:535-41.

101. Feugier P, Black RA, Hunt JA, et al. Attachment, morphology and adherence of human endothelial cells to vascular prosthesis materials under the action of shear stress. Biomaterials 2005;26:1457-66.

102. Bowlin GL, Rittgers SE. Electrostatic endothelial cell seeding technique for small-diameter (< 6 mm) vascular prostheses: Feasibility testing. Cell Transplantation 1997;6:623-9.

103. Schug-Pass C, Tamme C, Tannapfel A, et al. A lightweight polypropylene mesh (TiMesh) for laparoscopic intraperitoneal repair of abdominal wall hernias. Surg Endosc 2006;20:402-9.

104. Stiehler M, Lind M, Mygind T, et al. Morphology, proliferation, and osteogenic differentiation of mesenchymal stem cells cultured on titanium, tantalum, and chromium surfaces. Journal of Biomedical Materials Research Part A 2008;86:448-58.

105. Wu SL, Liu XM, Hu T, et al. A Biomimetic Hierarchical Scaffold: Natural Growth of Nanotitanates on Three-Dimensional Microporous Ti-Based Metals. Nano Lett 2008;8:3803-8. 106. Maeda M, Hirose M, Ohgushi H, et al. In vitro mineralization by mesenchymal stem cells cultured on titanium scaffolds. J Biochem 2007;141:729-36.

107. Morsi YS, Yang WW, Owida A, et al. Development of a novel pulsatile bioreactor for tissue culture. J Artif Organs 2007;10:109-14.

108. Barron V, Lyons E, Stenson-Cox C, et al. Bioreactors for cardiovascular cell and tissue growth: a review. Ann Biomed Eng 2003;31:1017-30.

109. Shachar M, Cohen S. Cardiac tissue engineering, ex-vivo: Design principles in biomaterials and bioreactors. Heart Fail Rev 2003;8:271-6.

110. Bilodeau K, Mantovani D. Bioreactors for tissue engineering: Focus on mechanical constraints. A comparative review. Tissue Eng 2006;12:2367-83.

111. Chen HC, Hu YC. Bioreactors for tissue engineering. Biotechnol Lett 2006;28:1415-23.

112. Goldstein AS, Christ G. Functional Tissue Engineering Requires Bioreactor Strategies Preface. Tissue Eng Pt A 2009;15:739-40.

113. Dumont K, Yperman J, Verbeken E, et al. Design of a new pulsatile bioreactor for tissue engineered aortic heart valve formation. Artif Organs 2002;26:710-4.

114. Engelmayr GC, Jr., Hildebrand DK, Sutherland FW, et al. A novel bioreactor for the dynamic flexural stimulation of tissue engineered heart valve biomaterials. Biomaterials 2003;24:2523-32.

115. Akins RE, Boyce RA, Madonna ML, et al. Cardiac organogenesis in vitro: Reestablishment of three-dimensional tissue architecture by dissociated neonatal rat ventricular cells. Tissue Eng 1999;5:103-18.

116. Freed LE, Vunjak-Novakovic G. Microgravity tissue engineering. In Vitro Cell Dev Biol Anim 1997;33:381-5.

117. Brown MA, Iyer RK, Radisic M. Pulsatile perfusion bioreactor for cardiac tissue engineering. Biotechnol Prog 2008;24:907-20.

118. Akhyari P, Fedak PW, Weisel RD, et al. Mechanical stretch regimen enhances the formation of bioengineered autologous cardiac muscle grafts. Circulation 2002;106:I137-42.

119. Fink C, Ergun S, Kralisch D, et al. Chronic stretch of engineered heart tissue induces hypertrophy and functional improvement. Faseb J 2000;14:669-79.

120. Sodian R, Lemke T, Loebe M, et al. New pulsatile bioreactor for fabrication of tissue-engineered patches. Journal of Biomedical Materials Research 2001;58:401-5.

121. Birla RK, Huang YC, Dennis RG. Development of a novel bioreactor for the mechanical loading of tissue-engineered heart muscle. Tissue Eng 2007;13:2239-48.

122. Hollweck T, Akra B, Häussler S. A Novel Pulsatile Bioreactor for Mechanical Stimulation of

Tissue Engineered Cardiac Constructs. J Funct Biomater 2011; 2:107-18.