

The Biological Significance of Long noncoding RNAs Dysregulation and their Mechanism of Regulating Signaling Pathways in Cervical Cancer

Maryame Lamsisi¹, Lahcen Wakrim², Amal Bouziyane^{1,3}, Mustapha Benhessou^{1, 4}, Mounia Oudghiri⁵, Abdelilah Laraoui⁶, Mohamed Elkarroumi⁴, Mohammed Ennachit⁴, Mohammed El Mzibri⁷, Moulay Mustapha Ennaji*

1. Team of Virology, Oncology and Medical Biotechnologies, Laboratory of Virology, Microbiology, Quality, and Biotechnologies/ ETB. Faculty of Science and Techniques Mohammedia, Hassan II University of Casablanca, Morocco.
2. Laboratory of Virology, Pasteur Institute of Morocco. Casablanca, Morocco.
3. University Mohammed VI of Health Science, Casablanca, Morocco.
4. School of Medicine and Pharmacy, University Hassan II of Casablanca, Morocco.
5. Immunology and Biodiversity laboratory, Faculty of Sciences Ain chock, Hassan II University of Casablanca, Morocco.
6. Research and Biosafety Laboratory, Mohammed V Military Hospital, University Mohammed V of Rabat, Morocco.
7. Biology and Medical Research Unit, CNESTEN, Rabat, Morocco.

Submitted 24 December 2020; Accepted 1 August 2021; Published 1 September 2021

Despite the remarkable decrease in cervical cancer incidence due to the availability of the HPV vaccine and implementation of screening programs for early detection in developed countries, this cancer remains a major health problem globally, especially in developing countries where most of the cases and mortality occur. Therefore, more understanding of molecular mechanisms of cervical cancer development might lead to the discovery of more effective diagnosis and treatment options. Research on long noncoding RNAs (lncRNAs) demonstrates the important roles of these molecules in many physiological processes and diseases, especially cancer. In the present review, we discussed the significance of lncRNAs altered expression in cervical cancer, highlighting their roles in regulating highly conserved signaling pathways, such as mitogen-activated protein kinase (MAPK), Wnt/ β -catenin, Notch, and phosphatidylinositol 3-kinase/protein kinase B (PI3K/AKT) pathways and their association with the progression of cervical cancer in order to bring more insight and understanding of this disease and their potential implications in cancer diagnosis and therapy.

Key words: Cervical cancer, human papillomavirus, long noncoding RNA, signaling pathways, gene regulation

Worldwide, cervical cancer (CC) is a major public health issue, ranking the fourth most

diagnosed cancer, and the second leading cause of cancer-related deaths in women (1). Clinical and

*Corresponding author: Faculty of Science and Techniques Mohammedia, University Hassan II of Casablanca, Morocco.
E-mail: m.ennaji@yahoo.fr

This work is published as an open access article distributed under the terms of the Creative Commons Attribution 4.0 License (<http://creativecommons.org/licenses/by-nc/4>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

epidemiological shreds of evidence reported that the occurrence of CC requires a prior persistent infection with human papillomavirus (HPV) (2). However, HPV infection alone is not sufficient and other cofactors including host genetic alterations and epigenetic modifications are needed for the progression from benign lesions to malignant tumors (3). The lack of accurate understanding of host factors and genetic background of this disease might explain the failure of current treatment options leading to high mortality rates.

The sequencing of the complete human genome by the human genome project in 2003 promised to offer more insights to our understanding of human physiology and resolving human genetic diseases including cancer (4). Thus, the encyclopedia of DNA elements (ENCODE) project that took over after human genome project completion, deciphered the obtained sequences and provided more in-depth data and analyzed the regulatory elements within the genome (5). Among the biggest discoveries of ENCODE is that the non-coding part of the genome which was described as junk DNA is mostly transcribed into functional RNA molecules, named non-coding RNAs (ncRNAs) (6). This part of the genome is not fully characterized despite the numerous studies on ncRNAs, providing an enormous field of genomics that is yet to be explored.

Several hypotheses are suggested regarding the role of ncRNAs, but their role in gene regulation is well discussed as they influence gene expression without DNA sequence alterations (7). ncRNAs are divided into 2 subclasses according to the length of the RNA molecule: small ncRNA (sncRNA) (20–200 nucleotides) and long ncRNA (lncRNA) (more than 200 nucleotides).

Emerging findings report that lncRNAs, with tissue-specific expression, are involved in diverse cellular and physiological pathways including cell differentiation, maintaining cellular homeostasis, regulation of the immune response to disease,

differentiation, and DNA damage repair (8,9). During malignancy, aberrant expression of lncRNAs is reported in many cancers, suggesting their role in the modulation of the physiological and molecular changes occurring in the transformed cells (9). Evidence from previous researches indicates that lncRNAs mainly interact with proteins, RNA, and DNA and function at transcriptional, translational, and post-translational levels (10). Moreover, Khalil et al. have reported that more than 20% of lncRNAs bind to the polycomb repressive complex 2 (PRC2) and other chromatin modifiers suggesting that chromatin modification might be a common mechanism of lncRNAs action (11).

In cervical cancer, an increasing number of functional studies have reported that dysregulation of the expression of diverse lncRNAs is involved in the regulation of malignant progression. In fact, the abnormal expression patterns of lncRNAs often correlate with the development and progression of cancer and play a crucial role in cell proliferation, invasion, and metastasis (12–14). LncRNAs exert their functions in CC mainly through the regulation of gene expression, which appears to be mediated by different processes such as chromatin state modulation and RNA processing (15). In CC, a number of lncRNAs showed abnormal expressions, such as HOX antisense intergenic RNA (*HOTAIR*), plasmacytoma variant translocation 1 (*PVT1*), and growth arrest specific 5 (*GAS5*), which are associated with disease progression and poor prognosis (16–18). On another hand, growing interest is given to the role of lncRNAs in viral replication and pathogenesis supporting their involvement in the host-pathogen interaction and suggesting the initiation and promotion of associated diseases (19,20). In the present review, we discuss the significance of lncRNAs altered expression in CC, highlighting their roles in regulating highly conserved signaling pathways, such as mitogen-activated protein kinase (MAPK),

Wnt/ β -catenin, Notch, and phosphatidylinositol 3-kinase/protein kinase B (PI3K/AKT) pathways and their association with the progression of CC.

The implication of lncRNAs in cancer progression

Up to now, many lncRNAs have been reported in CC and are involved in cell proliferation, cell cycle, apoptosis, epithelial to mesenchymal transition (EMT), migration, and/or invasion, such as *GAS5*, *HOTAIR*, metastasis associated lung adenocarcinoma transcript 1 (*MALAT1*), small nucleolar RNA host gene 8 (*SNHG8*), long intergenic non-protein coding RNA 511 (*LINC00511*) and *MAGI2* antisense RNA 3 (*MAGI2-AS3*) that were also widely considered as specific biomarkers for early diagnosis (21–26). Studies on the different mechanisms and interactions of lncRNAs with other genes and proteins that confirm the involvement of lncRNAs in CC development and progression are summarized in Table 1.

Almost all of the lncRNAs studied in CC interfere with cell proliferation through direct or indirect interaction with cell cycle proteins and apoptosis pathways. Highly expressed *C5orf66* antisense (*C5orf66-AS1*) in CC was reported to decrease the number of cells in the G1/G0 phase while increasing cell numbers in the G2/S phase. Moreover, overexpression of *C5orf66AS1* promoted the proliferation and affected apoptosis and cell cycle through adsorbing the regulator miR-637 (13). LncRNA *NCK1*-antisense 1 (*NCK1-AS1*) has been reported to promote cell proliferation and to induce cell cycle progression in CC by interacting with miR-6857 and affecting the cyclin-dependent kinase 1 pathway. *NCK1-AS1* induced elevated expression of cyclin dependent kinase (*CDK1/6*) by antagonizing miR-6857 and led to the control of the G1-S transition in CC cell lines (43,44).

Initially, *MAGI2-AS3* was reported to have tumor suppressive activities. However, Liu et al. have found that *MAGI2-AS3* up-regulated *CDK6*

and enhanced cell proliferation in CC (24). This oncogenic treat has been reported in other recent studies confirming that *MAGI2-AS3* promotes other cancers types such as colorectal and gastric cancers (45,46). On the other hand, *GAS5* was reported as a tumor suppressor lncRNA. Its ectopic overexpression induced cell cycle arrest at G2/M checkpoint which is mediated by the inhibition of cyclin B1 and *CDK1* expression by *GAS5*. Elevated expression of *BAX* and suppression of *BCL-2* is also a consequence of *GAS5* overexpression, which ultimately induces apoptosis (22).

LncRNA *MALAT1* was previously reported to be highly expressed in CC cells, and was correlated with cancer progression and metastasis (47). *MALAT1* is over-expressed in CC, and regulates the expression of apoptosis-related genes such as caspase-3 and 8, *BAX*, *BCL-2*, and *BCLxL* (48). Recent data suggest that HPV E6/E7 and IL-6/STAT3 signaling pathways work synergistically to up-regulate the transcription of *MALAT1* in CC HeLa cells, suggesting the cooperation of the virus oncoproteins with cellular inflammatory signaling in CC development (49).

In vitro studies on CC cell lines, showed that *HOTAIR* plays a role in apoptosis as its knockdown decreased protein levels of anti-apoptotic BCL-2, while it increased protein levels of pro-apoptotic BAX, apoptotic protease activating factor (APAF), caspase-3, caspase-9, and poly ADP-ribose polymerase (PARP) (26). *SNHG8*, another oncogenic lncRNA, promotes cell proliferation and inhibits apoptosis by recruiting enhancer of zeste homolog 2 (EZH2) to induce the trimethylation of reversion inducing cysteine rich protein with kazal motifs (*RECK*) promoter and thus inhibiting its expression (23). In addition, *LINC00511* recruits transcription factor retinoid X receptor alpha (RXRA) to upregulate the expression of phospholipase D1 (*PLD1*), and its knockdown promotes autophagy and apoptosis (21).

Table 1. LncRNAs interactions and roles in cervical cancer.

Lnc RNA	Expressi on level	Interaction with	Mechanism	Biological process	Ref.
<i>MALA T1</i>	Up	EMT genes	MALAT1 up-regulated Transcription factor snail and levels of β -catenin and Vimentin while downregulated E-cadherin and ZO-1	Invasion and Migration	(25)
<i>MAGI2-AS3</i>	Up	CDK6	MAGI2-AS3 up-regulated CDK6	Cell proliferation and cell cycle	(24)
<i>HAND2-AS1</i>	Down	ROCK1	HAND2-AS1 inhibited the expression of ROCK1	Cell proliferation and migration and invasion	(27)
	Down	C16orf74	HAND2-AS1 recruited transcription factor E2F4 to C16orf74 promoter and suppressed its expression	Cell proliferation, migration and invasion	(28)
<i>SOX20 T</i>	Depends on variants	SOX2	SOX20T modulated CC progression via the regulation of SOX2	Cell proliferation and migration and invasion	(29)
<i>SNHG1 6</i>	Up	SPI1/ PARP9 Axis	SNHG16 recruited SPI1 protein to promote transcription of PARP9.	Cell Proliferation, invasion and Cell Metastasis	(30)
<i>TUG1</i>	Up	PUM2	TUG1 enhanced the progression of CC by its interaction with PUM2.	Cell proliferation and migration	(31)
<i>MEG3</i>	Down	P-STAT3	MEG3 bound directly to P-STAT3 protein and induced its ubiquitination and degradation.	Cell proliferation, apoptosis	(32)
<i>LINC0 0511</i>	Up	RXRA/ PLD1	LINC00511 enriched RXRA to the promoter region of PLD1 and promoted its expression.	Cell proliferation, Apoptosis and tumor growth.	(21)
<i>MIR20 5HG</i>	Up	- SRSF1 - KRT17	Lnc-RNA MIR205HG regulated CC progression through KRT17 by binding with SRSF1	Cell proliferation, apoptosis and Migration	(33)
<i>lncOG FRP1</i>	Up	EMT and Apoptosis proteins	The depletion of lncOGFRP1 inhibited the expression of β -catenin, Vimentin, N-cadherin, SNAIL, Bcl-2, cyclinA1, CDK2, and PCNA, and promoted the expression of E-cadherin, Bax, p53, and caspase3	Cell proliferation, Cell cycle apoptosis and migration	(34)
<i>GPC3-AS1</i>	Up	GPC3	ELK1 acts as the transcription activator of GPC3- AS1 and GPC3	Cell proliferation and migration	(35)
<i>CRND E</i>	Up	PUMA	CRNDE binds to PUMA to inhibit its expression.	Cell proliferation, apoptosis and Tumor growth	(36)
<i>LINC0 0052</i>	Down	STAT3	The mRNA and protein expression of STAT3 was downregulated after overexpressing LINC00052.	Cell proliferation, tumor growth, invasion and migratin	(37)
<i>GAS5</i>	Down	Cyclin B1 and CDK1	GAS5 induced Cell cycle arrest by reducing the expression of Cyclin B1 and CDK1	Cell proliferation, Cell cycle, Apoptosis, tumor growth, Invasion and migration	(22)
<i>SNHG8</i>	Up	EZH2 / RECK	SNHG8 bound to EZH2 and epigenetically inhibited RECK transcription in CC.	Cell proliferation and migration	(23)
<i>SNHG1</i>	Up	ERK/Slug	SNHG12 is modulated by human	Tumor growth and	(38)

2			papillomavirus 16 E6/E7 and migration promoted CC progression via ERK/Slug pathway	
<i>Lnc-CC3</i>	Up	Slug	Lnc-CC3 increased Slug expression, and reduced the expression of E-cadherin.	Migration and invasion (39)
<i>ARAP1-AS1</i>	Up	c-Myc	ARAP1-AS1 might interact with PSF to release PTB, which accelerated IRES-driven translation of proto-oncogene c-Myc	Cell proliferation and migration (40)
<i>LINP1</i>	Up	KLF2 and PRSS8	LINP1 scaffolded EZH2, LSD1 and DNMT1 to suppress KLF2 and PRSS8	Cell proliferation and apoptosis (41)
<i>LINC02535</i>		PCBP2	LINC02535 interacted with PCBP2 to regulate DNA damage repair by stabilizing RRM1 mRNA in CC	Cell proliferation, migration and invasion, (42)

ARAP1-AS1: ARAP1 Antisense RNA 1; C16orf74: Chromosome 16 Open Reading Frame 74; CC: Cervical cancer; CDK1: Cyclin Dependent Kinase 1; CDK6: Cyclin Dependent Kinase 6; ROCK1: Rho Associated Coiled-Coil Containing Protein Kinase 1; CRNDE: Colorectal Neoplasia Differentially Expressed; DNMT1: DNA Methyltransferase 1; E2F4: E2F Transcription Factor 4; EMT: Epithelial to Mesenchymal Transition; ERK: Extracellular-signal-regulated kinase; EZH2: Enhancer Of Zeste 2 Polycomb Repressive Complex 2 Subunit; GAS5: Growth Arrest Specific 5; GPC3: Glypican 3; GPC3-AS1: GPC3 Antisense RNA 1; HAND2-AS1: HAND2 Antisense RNA 1; KLF2: Kruppel Like Factor 2; KRT17: Keratin 17; LINC00052: Long Intergenic Non-Protein Coding RNA 52; LINC00511: Long Intergenic Non-Protein Coding RNA 511; LINC02535: Long Intergenic Non-Protein Coding RNA 2535; LINP1: LncRNA In Non-Homologous End Joining Pathway 1; lncOGFRP1: Long noncoding RNA OGFRP1; LSD1: Lysine-specific demethylase 1; MAGI2-AS3: MAGI2 Antisense RNA 3; MALAT1: Metastasis Associated Lung Adenocarcinoma Transcript 1; MEG3: Maternally Expressed 3; MIR205HG: MIR205 Host Gene; PARP9: Poly(ADP-Ribose) Polymerase Family Member 9; PCBP2: Poly(RC) Binding Protein 2; PLD1: Phospholipase D1; PRSS8: Serine Protease 8; PSF: PTB-associated Splicing Factor; PTB: Polypyrimidine tract-binding protein; PUM2: Pumilio RNA Binding Family Member 2; PUMA: P53 upregulated modulator of apoptosis; RECK: Reversion Inducing Cysteine Rich Protein with Kazal Motifs; RRM1: Ribonucleotide Reductase Catalytic Subunit M1; RXRA: Retinoid X Receptor Alpha; SNHG12: Small Nucleolar RNA Host Gene 12; SNHG16: Small Nucleolar RNA Host Gene 16; SNHG8: Small Nucleolar RNA Host Gene 8; SOX2: SRY-Box Transcription Factor 2; SOX2OT: SOX2 Overlapping Transcript; SPI1: Spi-1 Proto-Oncogene; SRSF1: Serine and arginine Rich Splicing Factor 1; STAT3: Signal Transducer and Activator of Transcription 3; TUG1: Taurine Up-Regulated 1; ZO-1: Zonula occludens-1

The role of lncRNAs in epithelial to mesenchymal transition, invasion, and migration

EMT is a cellular biological program that drives the transition of cells between adherent epithelial state to mesenchymal phenotypes. Epithelial cells undergo series of changes to acquire the characteristics of mesenchymal cells such as stemness, motility, invasiveness, and resistance to therapy, leading to an increased ability of transformation and migration to distant organs (50). Several studies have indicated that EMT, invasion, and migration are in part regulated by some lncRNAs. For instance, lncRNA SRY-box transcription factor 2 (SOX2) overlapping transcript (SOX2OT) contributed to cell proliferation, migration and invasion of CC cells via the regulation of -box SOX2 (29). HOTAIR interacted with key genes that regulate cell invasion and

metastasis such as STAT3, β -catenin, vascular endothelial growth factor (VEGF), E-cadherin, matrix metalloproteinases (MMP-9), vimentin, snail, and twist, all of which are involved in EMT, invasion, and migration (51). Consistently, Lee et al. (2016) have investigated the expression levels of EMT related genes *in vivo* and found that β -catenin, N-cadherin, vimentin, snail, and twist were highly expressed in tumors overexpressing HOTAIR in comparison with the controls (26).

LncRNA-CTS may contribute to EMT, migration and invasion in CC cells through TGF- β 1. In fact, lncRNA-CTS regulates TGF- β 1 via sponging miR-505, which in turn is responsible for the regulation of zinc finger E-box binding homeobox 2 (ZEB2) mRNA (52). ZNF667-AS1 is a tumor suppressor lncRNA that also employs sponging microRNA mechanism to reduce tumor

invasion and metastasis in CC by competitive binding to miR-93-3p, and thus upregulating *PEG3* (53).

GAS5-ASI is another tumor suppressor that inhibits cell proliferation and metastasis of CC both *in vitro* and *in vivo* through increasing the expression of another tumor suppressor lncRNA, *GAS5*. *GAS5-ASI* appear to enhance the stability of *GAS5*, and thus increasing its expression, by reducing its N6-methyladenosine (m6A) modification (54).

Involvement of lncRNAs in signaling pathways

Deregulated expression of lncRNAs is involved in the initiation and promotion of CC development, invasion, and metastasis through their interactions with several signaling pathways. Numerous lncRNAs, comprising among others *HOTAIR*, *MALAT1*, *GAS5*, *EMT*, and maternally expressed gene 3 (*MEG3*) are involved in conserved signaling pathways such as Wnt, MAPK, NOTCH, and PI3K/AKT pathways (Table 2). Altogether, they have been shown to be associated with various pathogenic processes such as tumor progression, invasion as well as therapeutic resistance, and have emerged as new diagnostic and prognostic biomarkers in CC (55).

lncRNAs interfere with the Wnt signaling pathway in CC

Wnt/ β -catenin is a highly conserved signaling pathway that plays key roles in the development of cancer through modulating cell growth, cell regulation, and cell differentiation. Abnormal activation of the Wnt signaling pathway, which is the result of aberrant genetic and epigenetic regulation of its components, is linked to the progression of various types of cancers, including CC (84). As for every signaling pathway, Wnt pathway requires spatiotemporal regulation to maintain appropriate biological response and to prevent disease.

Several studies indicate that lncRNAs induce malignant behavior in CC by playing important

roles in this regulation. For instance, lncRNA colon cancer associated transcript 1 (*CCAT-1*) promotes cell proliferation through inhibiting apoptosis in CC cells and *RP11-480I12.5* induces the EMT of CC through the Wnt/ β -catenin pathway (56,60). In addition, lncRNA *ASB16* antisense RNA 1 (*ASB16-ASI*) acts as a sponge of miR-1305 to prevent its inhibitory effect on Wnt2 and enhance cell proliferation, migration, and invasion (65).

HOTAIR is one of the most studied lncRNAs that is overexpressed in several cancers including CC, and is known by its role in modulating chromatin state by scaffolding the three components of the chromatin-modifying complex PRC2: EZH2, SUZ12, and embryonic ectoderm development (EED) and directs them to distant targeted loci, which consequently induces the H3K27 tri-methylation on promoters of specific genes (16)(85). Through a similar mechanism, *HOTAIR* appears to regulate the Wnt/ β -catenin pathway as well. In fact, *HOTAIR* was found to recruit tet methylcytosine dioxygenase 1 (*TET1*) to induce methylation in the promoters of negative regulators of the Wnt/ β -catenin pathway such as protocadherin 10 (*PCDH10*), *SOX17*, adherens junctions associated protein 1 (*AJAPI*), and *MAGI2*, to decrease their expression in HeLa cells (66) (Figure 1).

In vitro downregulation of lncRNA cancer susceptibility 11 (*CASC11*) in HeLa cells, inhibits the activity of Wnt/ β -catenin signaling pathway while overexpression of *CASC11* in CaSki cells significantly up-regulated the signaling activity, suggesting that *CASC11* was involved in the activation of Wnt/ β -catenin signaling pathway (63). *CALML3* antisense RNA 1 (*CALML3-ASI*) is another overexpressed lncRNA in CC. The levels of the Wnt/ β -catenin pathway-related proteins such as β -catenin, cyclin D1, and c-MYC were observed to be down-regulated due to *CALML3-ASI* knockdown in CC cells, suggesting that the activity of Wnt/ β -catenin pathway is promoted by

CALML3-AS1, which might be the mechanism by

When Wnt is not expressed, cytoplasmic β -

Table 2. LncRNAs involved in regulating signaling pathways.

Pathway	LncRNAs involved		Ref.
	Activators	Inhibitors	
Wnt/β-catenin pathway	<i>CCAT1</i> ; <i>DANCR</i> ; <i>BLACAT1</i> ; <i>CALML3-AS1</i> ; <i>RP11-480I12.5</i> ; <i>SNHG7</i> ; <i>PCAT6</i> ; <i>CASC11</i> ; <i>NNT-AS1</i> ; <i>ASB16-AS1</i>	<i>DGCR5</i>	(56,57,66,67,58–65)
PI3K/AKT/mTOR pathway	<i>CRNDE</i> ; <i>RP1-93H18.6</i> ; <i>ANRIL</i> ; <i>CCAT1</i> ; <i>MFI2</i> ; <i>NEAT1</i> ; <i>MIAT</i>	<i>LINC00037</i> (<i>DGCR5</i>)	(68–74)(75)
NOTCH pathway	<i>HOTAIR</i> ; <i>SRA</i>	-	(26,76)
NF-κB Pathway	<i>PVT1</i> ; <i>NEAT1</i>	-	(77,78)
MAPK Pathway	<i>CASC2</i> ; <i>MXN1-AS1</i> ; <i>TUG1</i> ; <i>TDRG1</i>	-	(16,79–82)
JAK/STAT3	<i>LINC00518</i>	-	(83)

ANRIL: antisense non-coding RNA in the INK4 locus; ASB16-AS1: ASB16 antisense RNA 1; BLACAT1: bladder cancer associated transcript 1; CALML3-AS1: CALML3 antisense RNA 1; CASC11: cancer susceptibility 11; CASC2: cancer susceptibility 2; CCAT1: colon cancer associated transcript 1; CRNDE: colorectal neoplasia differentially expressed; DANCR: differentiation antagonizing non-protein coding RNA; DGCR5: DiGeorge syndrome critical region gene 5; HOTAIR : HOX transcript antisense RNA ; JAK: janus kinase; LINC00037: long intergenic non-protein coding RNA 37; LINC00518: long intergenic non-protein coding RNA 518; MAPK: mitogen-activated protein kinase 1; MFI2: melanotransferrin 2; MIAT: myocardial infarction associated transcript; MNX1-AS1: MNX1 antisense RNA 1; NEAT1: nuclear paraspeckle assembly transcript 1; NF- κ B: nuclear factor kappa B subunit 1; NNT-AS1: NNT antisense RNA 1; PCAT6: prostate cancer associated transcript 6; PVT1: plasmacytoma variant translocation 1; SNHG7: small nucleolar RNA host gene 7; SRA: steroid receptor RNA activator; STAT3: signal transducer and activator of transcription 3; TDRG1: testis development related 1; TUG1: taurine up-regulated 1.

which *CALML3-AS1* promotes CC (59) (Figure 2).

catenin is degraded by a protein complex composed




Fig. 1. Regulation of the Wnt/ β -catenin signaling pathway and MAPK pathway by lncRNAs and the crosstalk between the pathways. *DANCR* recruit *FRAT1* and *FRAT2* to negatively regulate GSK-3, which inhibits the accumulation of β -catenin and its translocation to the nucleus. LncRNA *GHET* positively regulates PI3K/AKT/mTOR pathway, which in turn targets GSK-3 and regulates Wnt/ β -catenin pathway.




Fig. 2. Involvement of lncRNAs in the regulation of conserved signaling pathways.

of axin protein, adenomatous polyposis coli (APC), the E3-ubiquitin ligase β -TrCP, CK1, ser/thr kinases GSK-3 protein phosphatase 2A (PP2A), and glycogen synthase kinase 3 (GSK3). β -catenin degradation prevents its transfer to the nucleus, and thus repressing the transcription of Wnt targeted genes (86).

Differentiation antagonizing non-protein coding RNA (*DANCR*) activates Wnt/ β -catenin signaling pathway through positively regulating frequently rearranged in advanced T-cell lymphomas 1 (*FRAT1*) and *FRAT2* expressions which belong to the GSK-3-binding proteins family that inhibit GSK-3-mediated β -catenin phosphorylation and degradation, which allows β -catenin to reach the nucleus to regulate targeted genes expression (57) (Figure 1). Consistently, the findings of this study indicated that induced overexpression of *DANCR* enhanced the mRNA and protein expression levels of *c-MYC* and cyclin D1, which are targeted genes of the Wnt/ β -catenin signaling pathway while knockdown of *DANCR* exhibited the opposite effect (57) (Figure 1).

lncRNAs regulation of phosphatidylinositol 3-kinase (PI3K)/protein kinase B (AKT) pathway in CC

PI3K is a member of the lipid kinases family. In the normal state of the cell, various extracellular factors, such as hormones, growth factors, and cytokines send signals to activate PI3K through the interaction with a phosphorylated tyrosine receptor. PI3K downstream cascade generates signals received by its targets, the most important one being the protein kinase B (AKT) that dominates the signal transduction of the PI3K pathway (87). Activation of AKT is a common phenomenon in human cancers leading to the promotion of cell proliferation (88). The entire PI3K/AKT signaling pathway plays key roles in regulating cell physiology and pathology, including apoptosis, cell proliferation, invasion, and metastasis (88). This pathway is abnormally activated in different tumors including CC (89).

Among the many regulators of this pathway, lncRNAs are also involved, adding more complexity to these processes. Decreased

expression of the antisense non-coding RNA in the *INK4* locus (*ANRIL*) inhibits cell proliferation, migration, and invasion in CC. After the inhibition of *ANRIL*, the PI3K/AKT pathway was found to be inactivated in CC cells, which indicates that *ANRIL* might regulate CC progression through the PI3K/AKT pathway (70). In addition, overexpressed *RPI-93H18.6* is an oncogenic lncRNA, its down-regulation resulted in the inhibition of cell proliferation and EMT in HeLa cells while promoting cell apoptosis via blocking the PI3K/AKT/mTOR signaling pathway (69). Moreover, lncRNA myocardial infarction associated transcript (*MIAT*) promotes CC and up-regulates PI3K, AKT, and mTOR levels, indicating its ability to activate PI3K/AKT/mTOR signaling pathway (74) (Figure 1).

LncRNA gastric carcinoma proliferation enhancing transcript 1 (*GHET1*) was found to regulate CC progression through modulating AKT/mTOR and its cross-talk with Wnt/ β -catenin pathways (Figure 1) (90). mTOR is one of the downstream targets of the PI3K/AKT axis. The mTOR axis is up-regulated in CC, and is suggested as a therapeutic target for anti-CC drug development. Blocking mTOR has shown a significant effect in treating HPV-related oral cancer and CCs (91). The crosstalk between AKT/mTOR and Wnt/ β -catenin has been demonstrated in many studies. In fact, p-AKT could induce the phosphorylation of Wnt protein receptor GSK3 β , which as mentioned above, induces the accumulation and nuclear migration of β -catenin, leading to the activation of Wnt/ β -catenin pathway (92) (Figure 2). However, the exact mechanism of action of lncRNAs in regulating this crosstalk in CC is not fully elucidated.

LncRNAs and Notch signaling pathway in CC

Notch signaling pathway plays an important role in different cellular processes such as cell proliferation and apoptosis. NOTCH signaling pathway has two main groups of ligands such as

delta-like 1, 3, and 4 and jagged 1 and 2. The binding of these ligands to NOTCH receptors, such as NOTCH 1, 2, 3, and 4 induces the activation of the pathway (93). The activation of the pathways triggers NOTCH cleavage and release of activated NOTCH intracellular domain (NICD). NICD is then translocated into the nucleus, where it activates the transcription of its targeted genes, mainly hairy and enhancer of split-1 (*HES1*), cyclin D1, and *c-MYC*. Otherwise, NOTCH can initiate the activation of other signaling pathways such as PI3K-AKT (94).

The Crosstalk between lncRNAs and Notch pathway was found in several solid cancers. For instance, lncRNA *MIR22HG* inhibits gastric cancer development and progression through its negative interaction with NOTCH2 signaling (95). LncRNA *SNHG12* promotes the progression of osteosarcoma by sponging miR-195-5p, thereby up regulating *NOTCH2* (96). *GHET1* promotes prostate cancer progression through targeting KLF2 which activates the HIF-1 α /NOTCH-1 pathway, and *MACC1-ASI* drives pancreatic cancer progression through activating PAX8/NOTCH1 signaling (97,98).

In CC, the NOTCH signaling pathway has a controversial role in alternating pro-oncogenic and tumor-suppressive roles (56). *In vitro* and *in vivo* studies showed that high levels of *HOTAIR* induce higher expression of *NOTCH1*, *HES1*, and *p300* in CC (26). Steroid receptor RNA activator (*SRA*) is a type of lncRNA which coordinates the functions of various transcription factors. *SRA* is related to the EMT and NOTCH signaling pathways, through which it induces *in vitro* tumor proliferation, migration and invasion (76) (Figure 2). These findings suggest that lncRNAs might promote CC through the NOTCH signaling pathway, representing an interesting way to deeply understand the complex role of this pathway in CC and its relation to lncRNAs.

The role of lncRNAs in mitogen-activated protein kinase (MAPK) pathways in

In its activated state, the MAPK phosphorylates its downstream targets in the nucleus and cytosol to regulate gene expression. There are three families of MAP kinases: JNKs (Jun amino-terminal kinases), ERKs (extracellular-signal-regulated kinases), and p38/SAPKs (stress-activated protein kinases). Numerous studies have shown that MAPK pathways play pivotal roles in CC (99), and numerous lncRNAs have been identified as regulators of the MAPK pathways in CC, through which they modulate cell proliferation, EMT, migration, and response to treatment (80–82).

lncRNA *CASC2* is reported to be down-regulated in CC, and acts as a tumor suppressor by inhibiting cell proliferation and migration. Overexpression of *CASC2* significantly inhibited the level of proteins of the MAPK pathway such as p-JNK and p-ERK1 *in vitro*, suggesting that *CASC2* might inhibit CC progression via negatively regulating the MAPK pathway (81). Jiang *et al.* demonstrated that testis development related gene 1 (*TDRG1*) sponged miR-326 to activate MAPK1, also known as ERK2, and thus suggested the miR-326/MAPK1 as a modulator of CC cell proliferation, migration, and invasion (79) (Figure 2).

In another study, lncRNA taurine up-regulated 1 (*TUG1*) controlled CC sensitivity to cisplatin through the MAPK pathway. *TUG1* knockdown inhibited the proliferative rate but accelerated the apoptosis of cisplatin-induced CC cells (82). Both mRNA and protein levels of regulatory factor X7 (*RFX7*) were down-regulated by the *TUG1* knockdown. Indeed, knockdown of *RFX7* could inhibit the proliferative rate and colony formation ability of CC cells. After cisplatin induction in CC cells, phosphorylated levels of *p38* and *JNK* increased, whereas *ERK1/2* expression decreased (82). *TUG1* knockdown could inhibit the proliferative rate and accelerate the apoptosis of CC cells by activating the MAPK pathway (82) (Figure

2). Zhang *et al.* analyzed the interaction between *HOTAIR* and STAT3. They identified a binding site for STAT3 in the promoter region of *HOTAIR* which is a GAS element. The genes containing GAS elements are regulated by STAT3, therefore, *HOTAIR* might be regulated by STAT3 as well. Moreover, they showed that *HOTAIR* and STAT3 affect synergically the aggressiveness of CC (100).

Competing endogenous pathway of lncRNAs in CC

It is widely accepted that gene regulation is more complex than previously expected, involving various regulators, enhancers, and/or transcription factors, acting in *cis* or in *trans*. Moreover, several studies have demonstrated that gene regulation is also mediated by microRNAs through complex mechanisms by which they interact with multiple networks. Since then, a growing interest was given to these microRNAs and their role in disease development, including cancer, which has been widely discussed and documented (101–103). Recently, several studies have reported that both coding and non-coding RNA molecules can regulate gene expression in *cis* and in *trans* by acting as sponges of microRNAs. These molecules, called competing endogenous RNAs (ceRNAs), represent a major group of gene regulators (104).

Intriguing relation is reported between lncRNAs and microRNAs; lncRNAs often act as molecular sponges or decoys to microRNAs and inactivate them. In turn, microRNAs have the ability to degrade lncRNAs. Together, lncRNAs and microRNAs can compete for binding sites on mRNAs (12,105) (Figure 3). Through this crosstalk between different RNA classes, lncRNAs regulate cancer progression and contribute to the regulation of cell proliferation, invasion, and migration in various cancer cells, including CC (12, 72, 73). Table 3 summarizes the main lncRNAs involved in CC development, their targeted microRNAs, and corresponding downstream dysregulated genes (12,105,106).

Of particular interest, most lncRNAs are up-regulated to sponge microRNAs and control cancer-

Table 3. Recent studies on ceRNA mechanism of lncRNAs in cervical cancer and downstream-targeted genes.

LncRNA	Expression level	Targeted miRNA	Downstream genes	Reference
<i>SNHG16</i>	Up-regulated	miR-216-5p	<i>ZEB1</i>	(107)
	Up-regulated	miR-128	<i>GSPT1</i> and <i>WNT3A</i>	(108)
<i>SNHG12</i>	Up-regulated	miR-125b	<i>STAT3</i>	(109)
<i>NEAT1</i>	Up-regulated	miR-133a	<i>SOX4</i>	(110)
	Up-regulated	miR-124	<i>NF-κB</i>	(78)
<i>MEG3</i>	Down-regulated	miR-7-5p	<i>STC1</i>	(111)
<i>MACC1-AS1</i>	Up-regulated	miR-34a	<i>CDK6</i>	(112)
<i>C5orf66-AS1</i>	Up-regulated	miR-637	<i>RING1</i>	(13)
<i>Linc00483</i>	Up-regulated	miR-508-3p	<i>RGS17</i>	(113)
<i>LINC01133</i>	Up-regulated	miR-4784	<i>AHDC1</i>	(114)
<i>LINC00152</i>	Up-regulated	miR-216b-5p	<i>HOXA1</i>	(115)
<i>lncRNA799</i>	Up-regulated	miR-454-3P	<i>TBL1XR1</i>	(116)
<i>LINC01503</i>	Up-regulated	miR-342-3p	<i>FXYD3</i>	(117)
<i>ZFPM2-AS1</i>	Up-regulated	miR-511-3p	<i>FGFR2</i>	(118)
<i>MAGI2-AS3</i>	Down-regulated	miRNA-233	<i>EPB41L3</i>	(119)
<i>MIR210HG</i>	Up-regulated	miR-503-5p	<i>TRAF4</i>	(120)
<i>LINC00173</i>	Down-regulated	miR-182-5p	<i>FBXW7</i>	(121)
<i>FENDRR</i>	Down-regulated	MiR-15a-5p/miR-15b-5p	<i>TUBA1A</i>	(122)
<i>CDKN2B-AS1</i>	Up-regulated	miR-181a-5p	<i>TGFβ1</i>	(123)
<i>LINC01128</i>	Up-regulated	miR-383-5p	<i>SFN</i>	(124)
<i>NNT-AS1</i>	Up-regulated	miR-186	<i>HMGB1</i>	(125)
<i>PITPNA-AS1</i>	Up-regulated	miR-876-5p	<i>c-MET</i>	(126)
<i>ZNF667-AS1</i>	Down-regulated	miR-93-3p	<i>PEG3</i>	(53)
<i>TTN-AS1</i>	Up-regulated	miR-573	<i>E2F3</i>	(127)
<i>FOXP4-AS1</i>	Up-regulated	miR-136-5p	<i>CBX4</i>	(128)
<i>CASC9</i>	Up-regulated	miR-215	<i>TWIST2</i>	(129)
<i>LINC00473</i>	Up-regulated	miR- 34a	<i>ILF2</i>	(130)
<i>TP73-AS1</i>	Up-regulated	microRNA-607	<i>Cyclin D2</i>	(131)
	Up-regulated	microRNA-329-3p	<i>SMAD2</i>	(132)
	Up-regulated	miR- 329- 3p	<i>ARF1</i>	(133)
<i>TUG1</i>	Up-regulated	miR-138-5p	<i>SIRT1</i>	(134)
<i>DDN-AS1</i>	Up-regulated	miR-15a/ miR-16	<i>TCF3</i>	(135)
<i>EWSAT1</i>	Up-regulated	miR-330-5p	<i>CPEB4</i>	(136)
<i>LINC00467</i>	Up-regulated	miR-107	<i>KIF23</i>	(137)
<i>SNHG20</i>	Up-regulated	miR-140-5p	<i>ADAM10</i>	(138)
<i>ATB</i>	Up-regulated	miR-144	<i>ITGA6</i>	(139)
<i>PCGEM1</i>	Up-regulated	miR-182	<i>FBXW11</i>	(140)
<i>CAR10</i>	Up-regulated	miR-125b-5p	<i>PDPK1</i>	(141)
<i>RP11-552M11.4</i>	Up-regulated	miR-3941	<i>ATF1</i>	(142)
<i>GAS5</i>	Down-regulated	miR-21	<i>STAT3</i>	(143)
<i>WT1-AS</i>	Down-regulated	miR-330-5p	<i>p53</i>	(144,145)
	Down-regulated	miR-203a-5p/	<i>FOXN2</i>	(146)
<i>HOTAIR</i>	Up-regulated	miR-148a	<i>HLA-G</i>	(147)
	Up-regulated	miR-23b	<i>MAPK1</i>	(16)

	Up-regulated	miR-143-3p	<i>BCL2</i>	(148)
	Up-regulated	miR206	<i>MKL1</i>	(149)
H19	Up-regulated	miR-138-5p	<i>SIRT1</i>	(150)
DSCAM-AS1	Up-regulated	mir-877-5p	<i>ATXN7L3</i>	(151)
RHPN1-AS1	Up-regulated	miR-299-3p	<i>FGF2</i>	(152)
SBF2-AS1	Up-regulated	miR-361-5p	<i>FOXMI</i>	(153)
NR2F2-AS1	Up-regulated	miR-4429	<i>MBD1</i>	(154)
POU3F3	Up-regulated	miR-127-5p	<i>FOXD1</i>	(155)
DLG1-AS1	Up-regulated	miR-107	<i>ZHX1</i>	(156)
HCP5	Up-regulated	microRNA-15a	<i>MACC1</i>	(157)
TCONS_00026907	Up-regulated	miR-143-5p	<i>ELK1</i>	(158)
CRNDE	Up-regulated	miR-183	<i>CCNB1</i>	(159)
SPRY4-IT1	Up-regulated	mir-101-3p	<i>ZEB1</i>	(160)
LINC01783	Up-regulated	mir-199b-5p	<i>GBP1</i>	(161)
miR503HG	Down-regulated	miR-155	<i>Caspase-3</i>	(162)
RHPN1-AS1	Up-regulated	miR-299-3p	<i>FGF2</i>	(152)
PVT1	Up-regulated	miR-140-5p	<i>SMAD3</i>	(17)
OIP5-AS1	Up-regulated	miR-143-3p	<i>SMAD3</i>	(163)
	Up-regulated	miR-143-3p	<i>ITGA6</i>	(164)
SNHG14	Up-regulated	miR-206	<i>YWHAZ</i>	(165)
UCA1	Up-regulated	miR-493-5p	<i>HK2</i>	(166)
STXBP5-AS1	Down-regulated	miR-96-5p	<i>PTEN</i>	(167)
CASC2	Down-regulated	miR-21	<i>PTEN</i>	(168)
TUSC8	Down-regulated	miR-641	<i>PTEN</i>	(169)
PTENP1	Down-regulated	miR-106b	<i>PTEN</i>	(170)
	Down-regulated	miR-19b	<i>MTUS1</i>	(171)
SOX21- AS1	Up-regulated	miR- 7	<i>VDAC1</i>	(172)
TMPO-AS1	Up-regulated	miR-577	<i>RAB14</i>	(173)
	Up-regulated	miR-143-3p	<i>ZEB1</i>	(174)
TP73-AS1	Up-regulated	miR-329-3p	<i>SMAD2</i>	(132)
	Up-regulated	miR-329-3p	<i>ARF1</i>	(133)
LINC01535	Up-regulated	miR- 214	<i>EZH2</i>	(175)
Linc00483	Up-regulated	miR-508-3p	<i>RGS17</i>	(113)
FOXD2-AS1	Up-regulated	miR-760	<i>HDGF</i>	(176)
PCAT6	Up-regulated	miR-543	<i>ZEB1</i>	(177)
MIR205HG	Up-regulated	miR-122e5p	<i>FOXP2</i>	(178)
SNHG7	Up-regulated	miR-485	<i>PAK4</i>	(179)
	Up-regulated	miR-485	<i>JUND</i>	(180)
CCAT1	Up-regulated	miR-181a-5p	<i>MMP14</i> and <i>HB-EGF</i>	(181)
HULC	Up-regulated	miR-218	<i>TPD52</i>	(182)
SOX21-AS1	Up-regulated	miR-7	<i>VDAC1</i>	(172)
BBOX1-AS1	Up-regulated	miR-361-3p	<i>HOXC6</i>	(183)
LncRNATP73- AS1	Up-regulated	miR- 329- 3p	<i>ARF1</i>	(133)
DLEU1	Up-regulated	miR-381	<i>HOXA13</i>	(184)
NOC2L- 4.1	Up-regulated	miR- 630	<i>YAP1</i>	(185)
LINC00319	Up-regulated	miR-3127-5p	<i>RPP25</i>	(186)
DLX6-AS1	Up-regulated	miR-16-5p	<i>ARPP19</i>	(187)

<i>XIST</i>	Up-regulated	miR-200a	<i>FUS</i>	(188)
	Up-regulated	miR-889-3p	<i>SIX1</i>	(189)
	Up-regulated	MiR-140-5p	<i>ORC1</i>	(190)
<i>OIP5-AS1</i>	Up-regulated	miR-143-3p	<i>ROCK1</i>	(191)
<i>LINC00958</i>	Up-regulated	miR- 625- 5p	<i>LRRC8E</i>	(192)
	Up-regulated	miR- 5095	<i>RRM2</i>	(193)
<i>SNHG4</i>	Up-regulated	miR-148a-3p	<i>c-MET</i>	(194)
<i>RUSC1-AS1</i>	Up-regulated	miR-744	<i>BCL-2</i>	(195)
<i>NOC2L-4.1</i>	Up-regulated	miR-630	<i>YAP1</i>	(185)
<i>TINCR</i>	Up-regulated	miR-302	<i>Cyclin D1</i>	(196)
<i>TDRG1</i>	Up-regulated	miR-326	<i>MAPK1</i>	(79)
	Up-regulated	miR-330-5p	<i>ELK1</i>	(197)
	Up-regulated	miR-214-5p	<i>SOX4</i>	(198)

ADAM10: A disintegrin and metalloproteinase 10; AHDC1: AT-hook DNA binding motif containing 1; ARF1: ADP ribosylation factor 1; ARPP19: cAMP regulated phosphoprotein 19; ATB: activated by transforming growth factor- β ; ATF1: activating transcription factor 1; ATXN7L3: ataxin 7 like 3; BBOX1-AS1: BBOX1 antisense RNA 1; BCL2: B-cell lymphoma 2; C5orf66-AS1: C5orf66 antisense RNA 1; CAR10: caspase recruitment domain family member 10; CASC2: cancer susceptibility 2; CASC9: cancer susceptibility 9; CBX4: chromobox 4; CCAT1: colon cancer associated transcript 1; CCNB1: cyclin B1; CDK6: cyclin dependent kinase 6; CDKN2B-AS1: CDKN2B antisense RNA 1; CPEB4: cytoplasmic polyadenylation element binding protein 4; CRNDE: colorectal neoplasia differentially expressed; DDN-AS1: DDN and PRKAG1 antisense RNA 1; DLEU1: deleted in lymphocytic leukemia 1; DLG1-AS1: DLG1 antisense RNA 1; DLX6-AS1: DLX6 antisense RNA 1; DSCAM-AS1: DSCAM antisense RNA 1; E2F3: E2F transcription factor 3; ELK1: ETS transcription factor ELK1; EPB41L3: erythrocyte membrane protein band 4.1 like 3; EWSAT1: Ewing sarcoma associated transcript 1; EZH2: enhancer of zeste homolog 2; FBXW11: F-box and WD repeat domain containing 11; FBXW7: F-box and WD repeat domain containing 7; FENDRR: FOXF1 adjacent non-coding developmental regulatory RNA; FGF2: fibroblast growth factor 2; FGFR2: fibroblast growth factor receptor 2; FOXD1: forkhead box D1; FOXD2-AS1: FOXD2 adjacent opposite strand RNA 1; FOXM1: forkhead box M1; FOXN2: forkhead box N2; FOXP2: forkhead box P2; FOXP4-AS1: FOXP4 antisense RNA 1; FUS: fused in sarcoma; FXYP3: FXYP domain containing ion transport regulator 3; GAS5: growth arrest specific 5; GBP1: guanylate binding protein 1; GSPT1: G1 to S phase transition 1; WNT3A: Wnt family member 3A; H19: H19 imprinted maternally expressed transcript; HB-EGF: heparin binding EGF like growth factor; HCP5: HLA complex P5; HDGF: heparin binding growth factor; HK2: hexokinase 2; HLA-G: major histocompatibility complex class I, G; HMGB1: high mobility group box 1; HOTAIR: HOX transcript antisense RNA; HOXA1: Homeobox A1; HOXA13: homeobox A13; HOXC6: homeobox C6; HULC: hepatocellular carcinoma up-regulated long non-coding RNA; ILF2: interleukin enhancer binding factor 2; ITGA6: integrin subunit alpha 6; JUND: jun D proto-oncogene subunit; KIF23: kinesin family member 23; LINC: long intergenic non-coding RNA; TP73-AS1: TP73 antisense RNA 1; LRRC8E: leucine rich repeat containing 8 VRAC subunit E; MACC1: MET transcriptional regulator MACC1; MACC1-AS1: MACC1 antisense RNA 1; MAGI2-AS3: MAGI2 antisense RNA 3; MAPK1: mitogen-activated protein kinase 1; MBD1: methyl-CpG binding domain protein 1; MEG3: maternally expressed gene 3; MIR205HG: MIR205 host gene; MKL1: megakaryoblastic leukemia 1; MMP14: matrix metalloproteinase 14; MTUS1: microtubule associated scaffold protein 1; NEAT1: nuclear paraspeckle assembly transcript 1; NF- κ B: nuclear factor kappa B subunit 1; NNT-AS1: NNT antisense RNA 1; NOC2L-4.1: long noncoding RNA NOC2L-4.1; NR2F2-AS1: NR2F2 antisense RNA 1; OIP5-AS1: OIP5 antisense RNA 1; ORC1: origin recognition complex subunit 1; PAK4: P21 activated kinase 4; PCAT6: prostate cancer associated transcript 6; PCGEM1: prostate-specific transcript; PDPK1: 3-phosphoinositide dependent Protein kinase 1; PEG3: paternally expressed 3; PIPNA-AS1: PIPNA antisense RNA 1; POU3F3: POU class 3 homeobox 3; PTEN: phosphatase and tensin homolog; PTENP1: phosphatase and tensin homolog pseudogene 1; PVT1: plasmacytoma variant translocation 1; RAB14: member RAS oncogene family; RGS17: regulator of G protein signaling 17; RHPN1-AS1: RHPN1 antisense RNA 1; RING1: ring finger protein 1; ROCK1: rho associated coiled-coil containing protein kinase 1; RPP25: ribonuclease P and MRP subunit P25; RRM2: ribonucleotide reductase regulatory subunit M2; RUSC1-AS1: RUSC1 antisense RNA 1; SBF2-AS1: SBF2 antisense RNA 1; STC1: stanniocalcin 1. SFN: stratifin; SIRT1: sirtuin 1; SIX1: sine oculis homeobox 1; SMAD2: SMAD family member 2; SMAD3: SMAD family member 3; SNHG: small nucleolar RNA host gene; SOX21-AS1: SOX21 antisense RNA 1; SOX4: SRY-box transcription factor 4; SPRY4-IT1: SPRY4 intronic transcript 1; STAT3: signal transducer and activator of transcription 3; STXP5-AS1: STXP5 antisense RNA 1; TBL1XR1: TBL1X receptor 1; TCF3: transcription factor 3; TDRG1: testis development related 1; TGF β 1: transforming growth factor beta 1; TINCR: TINCR ubiquitin domain containing; TMPO-AS1: TMPO antisense RNA 1; TP73-AS1: TP73 antisense RNA 1; TPD52: tumor protein D52; TRAF4: TNF receptor associated factor 4; TTN-AS1: TTN antisense RNA 1; TUBA1A: tubulin alpha 1a; TUG1: taurine up-regulated 1; TUSC8: tumor suppressor candidate 8; TWIST2: twist family BHLH transcription factor 2; UCA1: urothelial cancer associated 1; VDAC1: voltage dependent anion channel 1; WT1-AS: WT1 antisense RNA 1; XIST: X inactive specific transcript; YAP1: Yes1 associated transcriptional regulator; YWHAZ: tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein zeta; ZEB1: zinc finger E-box binding homeobox 1; ZFPM2-AS1: ZFPM2 antisense RNA 1; ZHX1: zinc fingers and homeoboxes 1; ZNF667-AS1: ZNF667 antisense RNA 1




Fig. 3. Competing endogenous RNAs interaction. There is a reciprocal negative regulation between LncRNAs and microRNAs, to both compete for mRNA binding sites. This competition leads eventually to gene expression and functional regulation.

development and progression. However, some of them are down-regulated and act as tumor suppressors. These include lncRNAs *STXBP5-AS1*, *TUSC8*, phosphatase and tensin homolog pseudogene 1 (*PTENP1*), and *CASC2* binding to miR-96-5p, miR-641, miR-106b, and miR-21 respectively, to regulate the expression of *PTEN* (167–170). And lncRNA *miR503HG*, *WT1-AS*, *GAS5*, *FENDRR*, *LINC00173*, *MAGI2-AS3*, *MEG3*, and *ZNF667-AS1* that bind to miR-155, miR-203a-5p, miR-330-5p, miR-21, miR-15a-5p/miR-15b-5p, miR-182-5p, miR-233, miR-7-5p, and miR-93-3p to regulate the expression of caspase-3, forkhead box N2 (*FOXN2*), P53, tubulin alpha 1a (*TUBA1A*), F-box/WD repeat-containing 7 (*FBXW7*), erythrocyte membrane protein band 4.1 like 3 (*EPB41L3*), *SCT1*, and paternally expressed gene (*PEG3*) that inhibit cell proliferation and induce apoptosis (53, 111, 119, 121, 122, 143, 146, 162).

The regulation of microRNAs by lncRNAs was also investigated for a better understanding of the treatment outcome in patients with CC. For instance, Feng et al. have shown that TNF- α treatment induced overexpression of lncRNA *LOC105374902*, which acts as a ceRNA for miR-1285-3p to promote the expression of ribosomal

protein L14 (*RPL14*), and thereby promoting the migration, invasion, and EMT of CC cells (199). Overexpression of lncRNA prostate cancer associated transcript 6 (*PCAT6*) down-regulated the expression of miR-543 in CC cells, thereby enhanced the level of zinc finger E-box-binding homeobox 1 (*ZEB1*), playing a key role in chemoresistance of CC cells to cisplatin, and consequently promoting cell proliferation and metastasis (177).

LncRNAs interaction with HPV in CC

HPV infection is a key event prior to CC development. Since HPV infection interferes with cellular mechanisms to induce aberrant cell proliferation, it was hypothesized that HPV interacts with lncRNAs in CC as well. Several studies demonstrated that lncRNAs are dysregulated in HPV positive cells and tissues (38, 200–203). This dysregulation is mainly mediated by HPV viral oncoproteins E6 and/or E7 (Figure 4).

Yang et al. reported significant change in lncRNAs expression patterns in HPV positive CC cell lines in comparison with HPV negative cells. They also found that these altered lncRNAs interacted with mRNAs that appear to play key roles in key cellular processes such as DNA repair, cell death, response to stimuli among others, all of




Fig. 4. LncRNAs and their interaction with HPV viral proteins in cervical cancer. A) viral oncoproteins E6 and E7 recruit transcriptional factor c-MYC to induce the expression of lncRNA *SNHG12*; B) E6 enhances the expression of lncRNA *FAM83H-AS1* through a mechanism involving P300; C) E6 and E7 form a regulatory feedback loop with lncRNA *TMPOP2*, where E6 and E7 inhibit P53 and its inhibitory effect on *TMPOP2* expression, and *TMPOP2* induces the expression of E6 and E7, to promote CC; D: HPV viral proteins E6 and/or E7 mediate the overexpression of lncRNAs *MALAT1*, *CCEPR*, *LINC01101* and *LINC00277* in CC.

which being involved in HPV related oncogenesis (203). In Barr et al. study, RNA high-throughput sequencing (RNA-seq) analysis indicated that the expression of host lncRNAs was altered in primary human foreskin keratinocytes cells (HEK) after infection with HPV16 *E6* oncogene. The study showed that 151 lncRNAs were up-regulated and 100 were down-regulated. In addition, altered expression of some lncRNAs was observed between pre-malignant and cancerous cervical cells (200). Of particular interest, they further evaluated the expression of *FAM83H-AS1* lncRNA in primary human cervical keratinocytes (HCK) infected with HPV 16 whole-genome and they found higher expression levels of *FAM83H-AS1* in comparison with controls (200). *FAM83H-AS1* expression was also increased in HPV 16 positive cervical cell lines

(CaSki, W12/201402, W12/20863), and decreased in HPV negative CC cell line (C-33A) in comparison with HCK cells (200). They demonstrated in the same study that *FAM83H-AS1* upregulation by HPV 16 is mediated specifically by E6 in a mechanism that does not involve its major downstream target p53. Instead, E6 regulates *FAM83H-AS1* through p300 (200).

In another study, HPV16 *E6* oncogene-induced lncRNA cervical carcinoma expressed PCNA regulatory (*CCEPR*) expression. Both HFK cells expressing HPV *E6/E7* and HPV positive CC cells (CaSki) expressed higher levels of *CCEPR*, suggesting the involvement of HPV in increasing *CCEPR* levels in CC. Moreover, *CCEPR* overexpression induced by HPV16 *E6* was reported to occur in a p53 independent manner (204).

Microarray analysis showed that 3626 lncRNAs were aberrantly expressed in HPV positive cervical squamous cell carcinoma samples versus HPV negative normal controls. Among them, 2077 lncRNAs were upregulated and 1549 lncRNAs were downregulated. Further qPCR analysis confirmed the overexpression of *SNHG 12*, *MALAT1*, *HCG11*, colorectal neoplasia differentially expressed (*CRNDE*), and *PVT1* (38). Lai et al. showed also that *SNHG12* expression is closely linked to the expression of HPV16 *E6* or *E7*; *SNHG12* expression was down-regulated in cells not expressing HPV16 *E6* or *E7* and up-regulated in cells overexpressing HPV16 *E6* or *E7*, suggesting that HPV16 oncoproteins E6 and E7 might regulate the expression of *SNHG12* lncRNA through the modulation of *c-MYC* (38).

In *E7*-siRNA transfected HeLa cells, microarray analysis showed that the expression of 15387 RNA species was modified in comparison with controls; among them were 731 lncRNAs and 203 lincRNAs indicating that HPV18 *E7* is involved in dysregulating of the expression of RNAs. Among the most dysregulated lincRNAs following *E7* depletion, *LINC01101* and *LINC00277* were particularly increased, which was further confirmed by qPCR analysis. In clinical samples of HPV positive CC patients, *LINC01101* and *LINC00277* expression was decreased in precancerous and cancerous lesions and their reduced expression correlated with high- risk HPV infections including HPV16 and HPV18 (205).

He et al. found that HPV16/18 proteins E6 and E7 promoted the expression of lncRNA thymopoietin pseudogene 2 (*TMPOP2*) in CC cells in a mechanism involving p53. Precisely, they found that p53 represses the expression of *TMPOP2* by direct binding to its promoter. *TMPOP2* in turn regulates the expression of HPV16/18 *E6/E7* and enhances their mRNA and protein level at a post-transcriptional level, suggesting that HPV16/18 *E6/E7* along with

lncRNA *TMPOP2* form a positive regulatory loop to regulate gene expression in CC in a synergic manner (206).

MALAT1 was significantly overexpressed in high-risk HPV positive CC cells and tissues in comparison with normal controls and promoted cell proliferation and invasion. In addition, knockdown of HPV *E6/E7* inhibited *MALAT1* expression in CasKi cells. In clinical samples, *MALAT1* was expressed in 30% of HPV-positive normal cervical cells and 60% of HPV-positive cervical lesions, while no expression of *MALAT1* was identified in HPV-negative normal cervical squamous cells (47).

Controversially, cells transfected with HPV16 *E7* expressed lower levels of *HOTAIR*, which was described in many studies cited above as an oncogene. Lower expressions of neuropilin 2 (*NRP2*) and P53 as well as a higher level of miR331-3p were also reported in cells transfected with HPV16 *E7*, which induced cell growth and inhibited apoptosis. Consistently with these findings, normal HPV positive cervical tissues also showed a reduced level of *HOTAIR* and *NRP2* in comparison with HPV negative normal cervical tissues (202). The interaction of lncRNAs with HPV infection has also diagnosis and therapeutic significance. For instance, lncRNA oncogene-induced senescence 1 (*OIS1*) was down-regulated in tissues and sera from HPV-positive patients with cervical squamous cell carcinoma and no significant differences were observed between HPV-negative patients and healthy controls. Consistently, *OIS1* expression levels were lower in HPV-positive cancer cell lines in comparison with that in HPV-negative cancer cell lines, while no significant differences were found between HPV-positive and HPV-negative normal cell lines. In addition, ROC curve analysis demonstrated that *OIS1* could potentially be used as a diagnostic marker for HPV positive but not for HPV negative cervical squamous cell carcinoma (207). Interestingly, it was found that damage induced

noncoding (DINO) lncRNA could restore the function of *TP53* in CC. The reactivation of *TP53* by *DINO* increases the vulnerability of CC to standard chemotherapeutics as well as biguanide compounds that cause metabolic stress, which suggests that this lncRNA could be used as a therapeutic alternative to the existing unsuccessful approaches (201).

Conclusion

The field of research on lncRNAs is growing each day with newly discovered molecules and new roles and mechanisms of already characterized ones; which provides a large variety of potential clinical applications. lncRNAs function either by direct interaction and inhibition of targeted signaling molecules or indirectly by binding other intermediate molecules such as mRNAs, proteins and microRNAs to alter their regulatory functions.

In CC, a number of lncRNAs such as *HOTAIR*, *PVT1*, *MALAT1*, and *GAS5*, which are associated with disease progression and prognosis, showed abnormal expressions. They are also involved in the regulation of conserved signaling pathways, such as the Wnt/ β -catenin, NOTCH, PI3k/AKT and MAPK pathways. In addition, most lncRNAs are up-regulated to sponge microRNAs and promote cancer development and progression, while, some of them are down-regulated and act as tumor suppressors; these include lncRNAs *STXBP5-AS1*, *TUSC8*, *PTEENP1*, and *CASC2*.

Giving the unavailability of effective treatments for most advanced CCs, lncRNAs diversity in terms of roles and mechanisms provides another set of opportunities. However, lncRNAs occupy several cellular localizations and exert their regulatory functions in a wide range of cellular and pathological contexts. A single lncRNA might also possess different binding sites, and can function through different mechanisms depending on the cellular context. Therefore, more thorough studies are needed to identify key binding sites and to uncover their exact mechanism of action in HPV

infection and CC progression to provide precise and targeted options for clinical applications. In addition, tissue specificity and the correlation of lncRNA expression to malignant phenotypes and also to viral infection provides a large field of biomarker research. Thus, more studies on the clinical applications of lncRNAs are required for new targeted therapy approaches and biomarker discoveries.

Acknowledgment

Authors would like to thank Hassan II University of Casablanca, Faculty of Science and techniques, team members of Virology, Oncology and Medical Biotechnology, and Virology, Microbiology, Quality and Biotechnologies / Ecotoxicology and Biodiversity laboratory; for all their efforts to support us during the writing and editing this review.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012;62:10-29.
2. Bosch FX, Lorincz A, Munoz N, et al. The causal relation between human papillomavirus and cervical cancer. *J Clin Pathol* 2002;55:244-65.
3. Moore EE, Wark JD, Hopper JL, et al. The roles of genetic and environmental factors on risk of cervical cancer: a review of classical twin studies. *Twin Res Hum Genet* 2012;15:79-86.
4. Green ED, Watson JD, Collins FS. Human Genome Project: Twenty-five years of big biology. *Nature* 2015;526:29-31.
5. Kellis M, Wold B, Snyder MP, et al. Defining functional DNA elements in the human genome. *Proc Natl Acad Sci U S A* 2014;111:6131-8.
6. Pennisi E. Genomics. ENCODE project writes eulogy for junk DNA. *Science* 2012;337:1159, 61.
7. Morlando M, Fatica A. Alteration of Epigenetic Regulation by Long Noncoding RNAs in Cancer. *Int J Mol Sci* 2018;19.
8. Chen YG, Satpathy AT, Chang HY. Gene regulation in the immune system by long noncoding RNAs. *Nat Immunol* 2017;18:962-72.

9. Schmitz SU, Grote P, Herrmann BG. Mechanisms of long noncoding RNA function in development and disease. *Cell Mol Life Sci* 2016;73:2491-509.
10. Dykes IM, Emanuelli C. Transcriptional and Post-transcriptional Gene Regulation by Long Non-coding RNA. *Genomics Proteomics Bioinformatics* 2017;15:177-86.
11. Khalil AM, Guttman M, Huarte M, et al. Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc Natl Acad Sci U S A* 2009;106:11667-72.
12. Tong R, Zhang J, Wang C, et al. LncRNA PTCSC3 inhibits the proliferation, invasion and migration of cervical cancer cells via sponging miR-574-5p. *Clin Exp Pharmacol Physiol* 2020;47:439-48.
13. Rui X, Xu Y, Jiang X, et al. Long non-coding RNA C5orf66-AS1 promotes cell proliferation in cervical cancer by targeting miR-637/RING1 axis. *Cell Death Dis* 2018;9:1175.
14. Zhang S, Zhang G, Liu J. Long noncoding RNA PVT1 promotes cervical cancer progression through epigenetically silencing miR-200b. *APMIS* 2016;124:649-58.
15. Kopp F, Mendell JT. Functional Classification and Experimental Dissection of Long Noncoding RNAs. *Cell* 2018;172:393-407.
16. Li Q, Feng Y, Chao X, et al. HOTAIR contributes to cell proliferation and metastasis of cervical cancer via targetting miR-23b/MAPK1 axis. *Biosci Rep* 2018;38.
17. Chang QQ, Chen CY, Chen Z, et al. LncRNA PVT1 promotes proliferation and invasion through enhancing Smad3 expression by sponging miR-140-5p in cervical cancer. *Radiol Oncol* 2019;53:443-52.
18. Yang W, Xu X, Hong L, et al. Upregulation of lncRNA GAS5 inhibits the growth and metastasis of cervical cancer cells. *J Cell Physiol* 2019;234:23571-80.
19. Lamsisi M, Ennaji MM. Involvement and roles of long noncoding RNAs in the molecular mechanisms of emerging and reemerging viral infections. In: Ennaji MM, editor. *Emerging and Reemerging Viral Pathogens*: Elsevier; 2020. p. 71-92.
20. Chen L, Zhou Y, Li H. LncRNA, miRNA and lncRNA-miRNA interaction in viral infection. *Virus Res* 2018;257:25-32.
21. Shi Y, Liu M, Huang Y, et al. Promotion of cell autophagy and apoptosis in cervical cancer by inhibition of long noncoding RNA LINC00511 via transcription factor RXRA-regulated PLD1. *J Cell Physiol* 2020;235:6592-604.
22. Yan Z, Ruoyu L, Xing L, et al. Long non-coding RNA GAS5 regulates the growth and metastasis of human cervical cancer cells via induction of apoptosis and cell cycle arrest. *Arch Biochem Biophys* 2020;684:108320.
23. Qu X, Li Y, Wang L, et al. LncRNA SNHG8 accelerates proliferation and inhibits apoptosis in HPV-induced cervical cancer through recruiting EZH2 to epigenetically silence RECK expression. *J Cell Biochem* 2020;121:4120-9.
24. Liu Q, Liu S, Wang X, et al. LncRNA MAGI2-AS3 is involved in cervical squamous cell carcinoma development through CDK6 up-regulation. *Infect Agent Cancer* 2019;14:37.
25. Sun R, Qin C, Jiang B, et al. Down-regulation of MALAT1 inhibits cervical cancer cell invasion and metastasis by inhibition of epithelial-mesenchymal transition. *Mol Biosyst* 2016;12:952-62.
26. Lee M, Kim HJ, Kim SW, et al. The long non-coding RNA HOTAIR increases tumour growth and invasion in cervical cancer by targeting the Notch pathway. *Oncotarget* 2016;7:44558-71.
27. Jin L, Ji J, Shi L, et al. LncRNA HAND2-AS1 inhibits cancer cell proliferation, migration and invasion by downregulating ROCK1 in HPV-positive and negative cervical squamous cell carcinoma. *Exp Ther Med* 2019;18:2512-8.
28. Gong J, Fan H, Deng J, et al. LncRNA HAND2-AS1 represses cervical cancer progression by interaction with transcription factor E2F4 at the promoter of C16orf74. *J Cell Mol Med* 2020;24:6015-27.
29. Chang X, Zhang H, Yang Q, et al. LncRNA SOX2OT affects cervical cancer cell growth, migration and invasion by regulating SOX2. *Cell Cycle* 2020;19:1391-403.
30. Tao L, Wang X, Zhou Q. Long noncoding RNA SNHG16 promotes the tumorigenicity of cervical cancer cells by recruiting transcriptional factor SPI1 to upregulate PARP9. *Cell Biol Int* 2020;44:773-84.
31. Duan W, Nian L, Qiao J, et al. LncRNA TUG1 aggravates the progression of cervical cancer by binding PUM2. *Eur Rev Med Pharmacol Sci* 2019;23:8211-8.
32. Zhang J, Gao Y. Long non-coding RNA MEG 3 inhibits cervical cancer cell growth by promoting degradation of P-STAT3 protein via ubiquitination. *Cancer Cell Int* 2019;19:175.
33. Dong M, Dong Z, Zhu X, et al. Long non-coding RNA MIR205HG regulates KRT17 and tumor processes in cervical

cancer via interaction with SRSF1. *Exp Mol Pathol* 2019;111:104322.

34. Zou K, Yu H, Chen X, et al. Silencing long noncoding RNA OGFRP1 inhibits the proliferation and migration of cervical carcinoma cells. *Cell Biochem Funct* 2019;37:591-7.

35. Hu R, Zhu Z. ELK1-activated GPC3-AS1/GPC3 axis promotes the proliferation and migration of cervical cancer cells. *J Gene Med* 2019;21:e3099.

36. Zhang JJ, Fan LP. Long non-coding RNA CRNDE enhances cervical cancer progression by suppressing PUMA expression. *Biomed Pharmacother* 2019;117:108726.

37. Lin J, Nong LL, Li MQ, et al. LINC00052 inhibits tumor growth, invasion and metastasis by repressing STAT3 in cervical carcinoma. *Eur Rev Med Pharmacol Sci* 2019;23:4673-9.

38. Lai SY, Guan HM, Liu J, et al. Long noncoding RNA SNHG12 modulated by human papillomavirus 16 E6/E7 promotes cervical cancer progression via ERK/Slug pathway. *J Cell Physiol* 2020;235:7911-22.

39. Jiang B, Sun R, Fang S, et al. Lnc-CC3 increases metastasis in cervical cancer by increasing Slug expression. *Oncotarget* 2016;7:41650-61.

40. Zhang Y, Wu D, Wang D. Long non-coding RNA ARAP1-AS1 promotes tumorigenesis and metastasis through facilitating proto-oncogene c-Myc translation via dissociating PSF/PTB dimer in cervical cancer. *Cancer Med* 2020;9:1855-66.

41. Wu L, Gong Y, Yan T, et al. LINP1 promotes the progression of cervical cancer by scaffolding EZH2, LSD1, and DNMT1 to inhibit the expression of KLF2 and PRSS8. *Biochem Cell Biol* 2020;98:591-9.

42. Wen D, Huang Z, Li Z, et al. LINC02535 co-functions with PCBP2 to regulate DNA damage repair in cervical cancer by stabilizing RRM1 mRNA. *J Cell Physiol* 2020;235:7592-603.

43. Huang L, Gan X, He L, et al. Silencing of long non-coding RNA NCK1-AS1 inhibits cell proliferation and migration via inhibition of microRNA-134 in cervical cancer. *Exp Ther Med* 2019;18:2314-22.

44. Li H, Jia Y, Cheng J, et al. LncRNA NCK1-AS1 promotes proliferation and induces cell cycle progression by crosstalk NCK1-AS1/miR-6857/CDK1 pathway. *Cell Death Dis* 2018;9:198.

45. Ren H, Li Z, Tang Z, et al. Long noncoding MAGI2-AS3 promotes colorectal cancer progression through regulating miR-3163/TMEM106B axis. *J Cell Physiol* 2020;235:4824-33.

46. Li D, Wang J, Zhang M, et al. LncRNA MAGI2-AS3 Is Regulated by BRD4 and Promotes Gastric Cancer Progression via Maintaining ZEB1 Overexpression by Sponging miR-141/200a. *Mol Ther Nucleic Acids* 2020;19:109-23.

47. Jiang Y, Li Y, Fang S, et al. The role of MALAT1 correlates with HPV in cervical cancer. *Oncol Lett* 2014;7:2135-41.

48. Guo F, Li Y, Liu Y, et al. Inhibition of metastasis-associated lung adenocarcinoma transcript 1 in CaSki human cervical cancer cells suppresses cell proliferation and invasion. *Acta Biochim Biophys Sin (Shanghai)* 2010;42:224-9.

49. Hao Y, Yan Z, Zhang A, et al. IL-6/STAT3 mediates the HPV18 E6/E7 stimulated upregulation of MALAT1 gene in cervical cancer HeLa cells. *Virus Res* 2020;281:197907.

50. Dongre A, Weinberg RA. New insights into the mechanisms of epithelial-mesenchymal transition and implications for cancer. *Nat Rev Mol Cell Biol* 2019;20:69-84.

51. Kim HJ, Lee DW, Yim GW, et al. Long non-coding RNA HOTAIR is associated with human cervical cancer progression. *Int J Oncol* 2015;46:521-30.

52. Feng S, Liu W, Bai X, et al. LncRNA-CTS promotes metastasis and epithelial-to-mesenchymal transition through regulating miR-505/ZEB2 axis in cervical cancer. *Cancer Lett* 2019;465:105-17.

53. Li YJ, Yang Z, Wang YY, et al. Long noncoding RNA ZNF667-AS1 reduces tumor invasion and metastasis in cervical cancer by counteracting microRNA-93-3p-dependent PEG3 downregulation. *Mol Oncol* 2019;13:2375-92.

54. Wang X, Zhang J, Wang Y. Long noncoding RNA GAS5-AS1 suppresses growth and metastasis of cervical cancer by increasing GAS5 stability. *Am J Transl Res* 2019;11:4909.

55. Tornosello ML, Faraonio R, Buonaguro L, et al. The Role of microRNAs, Long Non-coding RNAs, and Circular RNAs in Cervical Cancer. *Front Oncol* 2020;10:150.

56. Zhang J, Gao Y. CCAT-1 promotes proliferation and inhibits apoptosis of cervical cancer cells via the Wnt signaling pathway. *Oncotarget* 2017;8:68059-70.

57. Tian W, Lei N, Guo R, et al. Long non-coding RNA

- DANCR promotes cervical cancer growth via activation of the Wnt/beta-catenin signaling pathway. *Cancer Cell Int* 2020; 20:61.
58. Wang CH, Li YH, Tian HL, et al. Long non-coding RNA BLACAT1 promotes cell proliferation, migration and invasion in cervical cancer through activation of Wnt/beta-catenin signaling pathway. *Eur Rev Med Pharmacol Sci* 2018;22: 3002-9.
59. Liu CN, Zhang HY, Liu CL, et al. Upregulation of lncRNA CALML3-AS1 promotes cell proliferation and metastasis in cervical cancer via activation of the Wnt/beta-catenin pathway. *Eur Rev Med Pharmacol Sci* 2019;23:5611-20.
60. Zhang L, Li Y, Sona L. Long non-coding RNA RP11-480I12.5 promotes cervical carcinoma progression by regulating the Wnt/beta-catenin signaling pathway. *Oncol Lett* 2020;19:469-75.
61. Chi C, Li M, Hou W, et al. Long Noncoding RNA SNHG7 Activates Wnt/beta-Catenin Signaling Pathway in Cervical Cancer Cells by Epigenetically Silencing DKK1. *Cancer Biother Radiopharm* 2020;35:329-37.
62. Lv XJ, Tang Q, Tu YQ, et al. Long noncoding RNA PCAT6 regulates cell growth and metastasis via Wnt/beta-catenin pathway and is a prognosis marker in cervical cancer. *Eur Rev Med Pharmacol Sci* 2019;23:1947-56.
63. Hsu W, Liu L, Chen X, et al. LncRNA CASC11 promotes the cervical cancer progression by activating Wnt/beta-catenin signaling pathway. *Biol Res* 2019;52:33.
64. Hua F, Liu S, Zhu L, et al. Highly expressed long non-coding RNA NNT-AS1 promotes cell proliferation and invasion through Wnt/beta-catenin signaling pathway in cervical cancer. *Biomed Pharmacother* 2017;92:1128-34.
65. Liu W, Zhuang R, Feng S, et al. Long non-coding RNA ASB16-AS1 enhances cell proliferation, migration and invasion via functioning as a ceRNA through miR-1305/Wnt/beta-catenin axis in cervical cancer. *Biomed Pharmacother* 2020;125:109965.
66. Salmeron-Barcenas EG, Illades-Aguilar B, Del Moral-Hernandez O, et al. HOTAIR Knockdown Decreased the Activity Wnt/beta-Catenin Signaling Pathway and Increased the mRNA Levels of Its Negative Regulators in Hela Cells. *Cell Physiol Biochem* 2019;53:948-60.
67. Liu Y, Chang Y, Lu S, et al. Downregulation of long noncoding RNA DGCR5 contributes to the proliferation, migration, and invasion of cervical cancer by activating Wnt signaling pathway. *J Cell Physiol* 2019;234:11662-9.
68. Yang HY, Huang CP, Cao MM, et al. Long non-coding RNA CRNDE may be associated with poor prognosis by promoting proliferation and inhibiting apoptosis of cervical cancer cells through targeting PI3K/AKT. *Neoplasia* 2018;65:872-80.
69. Wang Q, Yan SP, Chu DX, et al. Silencing of Long Non-coding RNA RP1-93H18.6 Acts as a Tumor Suppressor in Cervical Cancer through the Blockade of the PI3K/Akt Axis. *Mol Ther Nucleic Acids* 2020;19:304-17.
70. Zhang D, Sun G, Zhang H, et al. Long non-coding RNA ANRIL indicates a poor prognosis of cervical cancer and promotes carcinogenesis via PI3K/Akt pathways. *Biomed Pharmacother* 2017;85:511-6.
71. Li R, Liu J, Qi J. Knockdown of long non-coding RNA CCAT1 suppresses proliferation and EMT of human cervical cancer cell lines by down-regulating Runx2. *Exp Mol Pathol* 2020;113:104380.
72. Qu B, Zhao AH, Nie XZ, et al. Up-regulation of long non-coding RNA MF12 functions as an oncogenic role in cervical cancer progression. *Eur Rev Med Pharmacol Sci* 2019;23: 4680-7.
73. Guo HM, Yang SH, Zhao SZ, et al. LncRNA NEAT1 regulates cervical carcinoma proliferation and invasion by targeting AKT/PI3K. *Eur Rev Med Pharmacol Sci* 2018;22:4090-7.
74. Zhang L, Ge S, Cao B. Long non-coding RNA MIAT promotes cervical cancer proliferation and migration. *J Biochem* 2020;168:183-90.
75. Yuan LY, Qin X, Li L, et al. Overexpression of LINC00037 represses cervical cancer progression by activating mTOR signaling pathway. *J Cell Physiol* 2019;234:13353-60.
76. Eoh KJ, Paek J, Kim SW, et al. Long non-coding RNA, steroid receptor RNA activator (SRA), induces tumor proliferation and invasion through the NOTCH pathway in cervical cancer cell lines. *Oncol Rep* 2017;38:3481-8.
77. Wang C, Zou H, Yang H, et al. Long noncoding RNA plasmacytoma variant translocation 1 gene promotes the development of cervical cancer via the NFkappaB pathway. *Mol Med Rep* 2019;20:2433-40.
78. Shen X, Zhao W, Zhang Y, et al. Long Non-Coding RNA-NEAT1 Promotes Cell Migration and Invasion via Regulating

miR-124/NF-kappaB Pathway in Cervical Cancer. *Onco Targets Ther* 2020;13:3265-76.

79. Jiang H, Liang M, Jiang Y, et al. The lncRNA TDRG1 promotes cell proliferation, migration and invasion by targeting miR-326 to regulate MAPK1 expression in cervical cancer. *Cancer Cell Int* 2019;19:152.

80. Liu X, Yang Q, Yan J, et al. LncRNA MNX1-AS1 promotes the progression of cervical cancer through activating MAPK pathway. *J Cell Biochem* 2019;120:4268-77.

81. Wang XW, Zhang W. Long non-coding RNA cancer susceptibility candidate 2 inhibits the cell proliferation, invasion and angiogenesis of cervical cancer through the MAPK pathway. *Eur Rev Med Pharmacol Sci* 2019;23:3261-9.

82. Wei X, Zhou Y, Qiu J, et al. Low expression of TUG1 promotes cisplatin sensitivity in cervical cancer by activating the MAPK pathway. *J BUON* 2019;24:1020-6.

83. Wang DW, You D, Dong J, et al. Knockdown of long non-coding RNA LINC00518 inhibits cervical cancer proliferation and metastasis by modulating JAK/STAT3 signaling. *Eur Rev Med Pharmacol Sci* 2019;23:496-506.

84. Yang M, Wang M, Li X, et al. Wnt signaling in cervical cancer? *J Cancer* 2018;9:1277-86.

85. Gupta RA, Shah N, Wang KC, et al. Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature* 2010;464:1071-6.

86. Stamos JL, Weis WI. The beta-catenin destruction complex. *Cold Spring Harb Perspect Biol* 2013;5:a007898.

87. Arcaro A, Guerreiro AS. The phosphoinositide 3-kinase pathway in human cancer: genetic alterations and therapeutic implications. *Curr Genomics* 2007;8:271-306.

88. Martini M, De Santis MC, Braccini L, et al. PI3K/AKT signaling pathway and cancer: an updated review. *Ann Med* 2014;46:372-83.

89. Bossler F, Hoppe-Seyler K, Hoppe-Seyler F. PI3K/AKT/mTOR Signaling Regulates the Virus/Host Cell Crosstalk in HPV-Positive Cervical Cancer Cells. *Int J Mol Sci* 2019;20.

90. Liu Z, Luo S, Wu M, et al. LncRNA GHET1 promotes cervical cancer progression through regulating AKT/mTOR and Wnt/beta-catenin signaling pathways. *Biosci Rep* 2020;40.

91. Molinolo AA, Marsh C, El Dinali M, et al. mTOR as a

molecular target in HPV-associated oral and cervical squamous carcinomas. *Clin Cancer Res* 2012;18:2558-68.

92. Jain S, Ghanghas P, Rana C, et al. Role of GSK-3beta in Regulation of Canonical Wnt/beta-catenin Signaling and PI3-K/Akt Oncogenic Pathway in Colon Cancer. *Cancer Invest* 2017;35:473-83.

93. Brzozowa-Zasada M, Piecuch A, Dittfeld A, et al. Notch signalling pathway as an oncogenic factor involved in cancer development. *Contemp Oncol (Pozn)* 2016;20:267-72.

94. Rodrigues C, Joy LR, Sachithanandan SP, et al. Notch signalling in cervical cancer. *Exp Cell Res* 2019;385:111682.

95. Li H, Wang Y. Long Noncoding RNA (lncRNA) MIR22HG Suppresses Gastric Cancer Progression through Attenuating NOTCH2 Signaling. *Med Sci Monit* 2019;25:656-65.

96. Zhou S, Yu L, Xiong M, et al. LncRNA SNHG12 promotes tumorigenesis and metastasis in osteosarcoma by upregulating Notch2 by sponging miR-195-5p. *Biochem Biophys Res Commun* 2018;495:1822-32.

97. Zhu Y, Tong Y, Wu J, et al. Knockdown of LncRNA GHET1 suppresses prostate cancer cell proliferation by inhibiting HIF-1alpha/Notch-1 signaling pathway via KLF2. *Biofactors* 2019;45:364-73.

98. Qi C, Xiaofeng C, Dongen L, et al. Long non-coding RNA MACC1-AS1 promoted pancreatic carcinoma progression through activation of PAX8/NOTCH1 signaling pathway. *J Exp Clin Cancer Res* 2019;38:344.

99. Morrison DK. MAP kinase pathways. *Cold Spring Harb Perspect Biol* 2012;4.

100. Zhang Y, Cheng X, Liang H, et al. Long non-coding RNA HOTAIR and STAT3 synergistically regulate the cervical cancer cell migration and invasion. *Chem Biol Interact* 2018;286:106-10.

101. Maegdefessel L. The emerging role of microRNAs in cardiovascular disease. *J Intern Med* 2014;276:633-44.

102. Ciccacci C, Politi C, Novelli G, et al. Advances in Exploring the Role of Micrnas in Inflammatory Bowel Disease. *Microna* 2016;5:5-11.

103. Kwan JY, Psarianos P, Bruce JP, et al. The complexity of microRNAs in human cancer. *J Radiat Res* 2016;57 Suppl 1:i106-i11.

104. Bayoumi AS, Sayed A, Broskova Z, et al. Crosstalk

between Long Noncoding RNAs and MicroRNAs in Health and Disease. *Int J Mol Sci* 2016;17:356.

105. Du H, Chen Y. Competing endogenous RNA networks in cervical cancer: function, mechanism and perspective. *J Drug Target* 2019;27:709-23.

106. Huang Y, Luo H, Li F, et al. LINC00152 down-regulated miR-193a-3p to enhance MCL1 expression and promote gastric cancer cells proliferation. *Biosci Rep* 2018;38.

107. Zhu H, Zeng Y, Zhou CC, et al. SNHG16/miR-216-5p/ZEB1 signal pathway contributes to the tumorigenesis of cervical cancer cells. *Arch Biochem Biophys* 2018;637:1-8.

108. Wu W, Guo L, Liang Z, et al. Lnc-SNHG16/miR-128 axis modulates malignant phenotype through WNT/beta-catenin pathway in cervical cancer cells. *J Cancer* 2020;11:2201-12.

109. Jin XJ, Chen XJ, Zhang ZF, et al. Long noncoding RNA SNHG12 promotes the progression of cervical cancer via modulating miR-125b/STAT3 axis. *J Cell Physiol* 2019;234:6624-32.

110. Yuan LY, Zhou M, Lv H, et al. Involvement of NEAT1/miR-133a axis in promoting cervical cancer progression via targeting SOX4. *J Cell Physiol* 2019;234:18985-93.

111. Pan X, Cao YM, Liu JH, et al. MEG3 Induces Cervical Carcinoma Cells' Apoptosis Through Endoplasmic Reticulum Stress by miR-7-5p/STC1 Axis. *Cancer Biother Radiopharm* 2021;36:501-10.

112. Jin J, Chen X, Chen J, et al. Long noncoding RNA MACC1-AS1 is a potential sponge of microRNA-34a in cervical squamous cell carcinoma and upregulates cyclin-dependent kinase 6. *Oncol Lett* 2020;19:2339-45.

113. Hu P, Zhou G, Zhang X, et al. Long non-coding RNA Linc00483 accelerated tumorigenesis of cervical cancer by regulating miR-508-3p/RGS17 axis. *Life Sci* 2019;234:116789.

114. Feng Y, Qu L, Wang X, et al. LINC01133 promotes the progression of cervical cancer by sponging miR-4784 to up-regulate AHDC1. *Cancer Biol Ther* 2019;20:1453-61.

115. Zheng JJ, Du XJ, Wang HP, et al. Long non-coding RNA 00152 promotes cell proliferation in cervical cancer via regulating miR-216b-5p/HOXA1 axis. *Eur Rev Med Pharmacol Sci* 2019;23:3654-63.

116. Liao LM, Zhang FH, Yao GJ, et al. Role of Long Noncoding RNA 799 in the Metastasis of Cervical Cancer through Upregulation of TBL1XR1 Expression. *Mol Ther*

Nucleic Acids 2018;13:580-9.

117. Peng X, Gao J, Cai C, et al. LncRNA LINC01503 aggravates the progression of cervical cancer through sponging miR-342-3p to mediate FXYD3 expression. *Biosci Rep* 2020;40.

118. Dai J, Wei R, Zhang P, et al. Long Noncoding RNA ZFPM2-AS1 Enhances the Malignancy of Cervical Cancer by Functioning as a Molecular Sponge of microRNA-511-3p and Consequently Increasing FGFR2 Expression. *Cancer Manag Res* 2020;12:567-80.

119. Hou A, Zhang Y, Fan Y, et al. LncRNA MAGI2-AS3 Affects Cell Invasion and Migration of Cervical Squamous Cell Carcinoma (CSCC) via Sponging miRNA-233/EPB41L3 Axis. *Cancer Manag Res* 2020;12:4209-16.

120. Wang AH, Jin CH, Cui GY, et al. MIR210HG promotes cell proliferation and invasion by regulating miR-503-5p/TRAF4 axis in cervical cancer. *Aging (Albany NY)* 2020;12:3205-17.

121. Zhang J, Zhou M, Zhao X, et al. Long noncoding RNA LINC00173 is downregulated in cervical cancer and inhibits cell proliferation and invasion by modulating the miR-182-5p/FBXW7 axis. *Pathol Res Pract* 2020;216:152994.

122. Zhu Y, Zhang X, Wang L, et al. FENRRR suppresses cervical cancer proliferation and invasion by targeting miR-15a/b-5p and regulating TUBA1A expression. *Cancer Cell Int* 2020;20:152.

123. Zhu L, Zhang Q, Li S, et al. Interference of the long noncoding RNA CDKN2B-AS1 upregulates miR-181a-5p/TGFbeta1 axis to restrain the metastasis and promote apoptosis and senescence of cervical cancer cells. *Cancer Med* 2019;8:1721-30.

124. Hu Y, Ma Y, Liu J, et al. LINC01128 expedites cervical cancer progression by regulating miR-383-5p/SFN axis. *BMC Cancer* 2019;19:1157.

125. Liu Y, Guo R, Qiao Y, et al. LncRNA NNT-AS1 contributes to the cisplatin resistance of cervical cancer through NNT-AS1/miR-186/HMGB1 axis. *Cancer Cell Int* 2020;20:190.

126. Guo Q, Li L, Bo Q, et al. Long noncoding RNA PITPNA-AS1 promotes cervical cancer progression through regulating the cell cycle and apoptosis by targeting the miR-876-5p/c-MET axis. *Biomed Pharmacother* 2020;128:110072.

127. Chen P, Wang R, Yue Q, et al. Long non-coding RNA TTN-AS1 promotes cell growth and metastasis in cervical cancer via miR-573/E2F3. *Biochem Biophys Res Commun*

- 2018;503:2956-62.
128. Zhao J, Yang T, Li L. LncRNA FOXP4-AS1 Is Involved in Cervical Cancer Progression via Regulating miR-136-5p/CBX4 Axis. *Onco Targets Ther* 2020;13:2347-55.
129. Zhang J, Wang Q, Quan Z. Long non-coding RNA CASC9 enhances breast cancer progression by promoting metastasis through the mediation of miR-215/TWIST2 signaling associated with TGF-beta expression. *Biochem Biophys Res Commun* 2019;515:644-50.
130. Shi C, Yang Y, Yu J, et al. The long noncoding RNA LINC00473, a target of microRNA 34a, promotes tumorigenesis by inhibiting ILF2 degradation in cervical cancer. *Am J Cancer Res* 2017;7:2157-68.
131. Zhang H, Xue B, Wang S, et al. Long noncoding RNA TP73 antisense RNA 1 facilitates the proliferation and migration of cervical cancer cells via regulating microRNA607/cyclin D2. *Mol Med Rep* 2019;20:3371-8.
132. Guan MM, Rao QX, Huang ML, et al. Long Noncoding RNA TP73-AS1 Targets MicroRNA-329-3p to Regulate Expression of the SMAD2 Gene in Human Cervical Cancer Tissue and Cell Lines. *Med Sci Monit* 2019;25:8131-41.
133. Xu J, Zhang J. LncRNA TP73-AS1 is a novel regulator in cervical cancer via miR-329-3p/ARF1 axis. *J Cell Biochem* 2020;121:344-52.
134. Ohms S, Rangasamy D. Silencing of LINE-1 retrotransposons contributes to variation in small noncoding RNA expression in human cancer cells. *Oncotarget* 2014;5:4103-17.
135. Liu Z, Wu M, Shi H, et al. DDN-AS1-miR-15a/16-TCF3 feedback loop regulates tumor progression in cervical cancer. *J Cell Biochem* 2019;120:10228-38.
136. Zhou Q, Xie Y, Wang L, et al. LncRNA EWSAT1 upregulates CPEB4 via miR-330-5p to promote cervical cancer development. *Mol Cell Biochem* 2020;471:177-88.
137. Li GC, Xin L, Wang YS, et al. Long Intervening Noncoding 00467 RNA Contributes to Tumorigenesis by Acting as a Competing Endogenous RNA against miR-107 in Cervical Cancer Cells. *Am J Pathol* 2019;189:2293-310.
138. Guo H, Yang S, Li S, et al. LncRNA SNHG20 promotes cell proliferation and invasion via miR-140-5p-ADAM10 axis in cervical cancer. *Biomed Pharmacother* 2018;102:749-57.
139. Zhu Y, Wu Y, Yang L, et al. Long non-coding RNA activated by transforming growth factor-beta promotes proliferation and invasion of cervical cancer cells by regulating the miR-144/ITGA6 axis. *Exp Physiol* 2019;104:837-44.
140. Zhang Q, Zheng J, Liu L. The long noncoding RNA PCGEM1 promotes cell proliferation, migration and invasion via targeting the miR-182/FBXW11 axis in cervical cancer. *Cancer Cell Int* 2019;19:304.
141. Hu T, Zhang Q, Gao L. LncRNA CAR10 Upregulates PDPK1 to Promote Cervical Cancer Development by Sponging miR-125b-5p. *Biomed Res Int* 2020;2020:4351671.
142. Xu Y, Zhou W, Zhang C, et al. Long non-coding RNA RP11-552M11.4 favors tumorigenesis and development of cervical cancer via modulating miR-3941/ATF1 signaling. *Int J Biol Macromol* 2019;130:24-33.
143. Yao T, Lu R, Zhang J, et al. Growth arrest-specific 5 attenuates cisplatin-induced apoptosis in cervical cancer by regulating STAT3 signaling via miR-21. *J Cell Physiol* 2019;234:9605-15.
144. Cui L, Nai M, Zhang K, et al. LncRNA WT1-AS inhibits the aggressiveness of cervical cancer cell via regulating p53 expression via sponging miR-330-5p. *Cancer Manag Res* 2019;11:651-67.
145. Zhang Y, Na R, Wang X. LncRNA WT1-AS up-regulates p53 to inhibit the proliferation of cervical squamous carcinoma cells. *BMC Cancer* 2019;19:1052.
146. Dai SG, Guo LL, Xia X, et al. Long non-coding RNA WT1-AS inhibits cell aggressiveness via miR-203a-5p/FOXN2 axis and is associated with prognosis in cervical cancer. *Eur Rev Med Pharmacol Sci* 2019;23:486-95.
147. Sun J, Chu H, Ji J, et al. Long non-coding RNA HOTAIR modulates HLA-G expression by absorbing miR-148a in human cervical cancer. *Int J Oncol* 2016;49:943-52.
148. Liu M, Jia J, Wang X, et al. Long non-coding RNA HOTAIR promotes cervical cancer progression through regulating BCL2 via targeting miR-143-3p. *Cancer Biol Ther* 2018;19:391-9.
149. Zheng P, Yin Z, Wu Y, et al. LncRNA HOTAIR promotes cell migration and invasion by regulating MKL1 via inhibition miR206 expression in HeLa cells. *Cell Commun Signal* 2018;16:5.

150. Ou L, Wang D, Zhang H, et al. Decreased Expression of miR-138-5p by lncRNA H19 in Cervical Cancer Promotes Tumor Proliferation. *Oncol Res* 2018;26:401-10.
151. Liang J, Zhang S, Wang W, et al. Long non-coding RNA DSCAM-AS1 contributes to the tumorigenesis of cervical cancer by targeting miR-877-5p/ATXN7L3 axis. *Biosci Rep* 2020;40.
152. Duan H, Li X, Chen Y, et al. LncRNA RHPN1-AS1 promoted cell proliferation, invasion and migration in cervical cancer via the modulation of miR-299-3p/FGF2 axis. *Life Sci* 2019;239:116856.
153. Gao F, Feng J, Yao H, et al. LncRNA SBF2-AS1 promotes the progression of cervical cancer by regulating miR-361-5p/FOXM1 axis. *Artif Cells Nanomed Biotechnol* 2019;47:776-82.
154. Liu D, Huang K, Wang T, et al. NR2F2-AS1 accelerates cell proliferation through regulating miR-4429/MBD1 axis in cervical cancer. *Biosci Rep* 2020;40.
155. Chang S, Sun L, Feng G. SP1-mediated long noncoding RNA POU3F3 accelerates the cervical cancer through miR-127-5p/FOXD1. *Biomed Pharmacother* 2019;117:109133.
156. Rui X, Xu Y, Huang Y, et al. lncRNA DLG1-AS1 Promotes Cell Proliferation by Competitively Binding with miR-107 and Up-Regulating ZHX1 Expression in Cervical Cancer. *Cell Physiol Biochem* 2018;49:1792-803.
157. Yu Y, Shen HM, Fang DM, et al. LncRNA HCP5 promotes the development of cervical cancer by regulating MACC1 via suppression of microRNA-15a. *Eur Rev Med Pharmacol Sci* 2018;22:4812-9.
158. Jin X, Chen X, Hu Y, et al. LncRNA-TCONS_00026907 is involved in the progression and prognosis of cervical cancer through inhibiting miR-143-5p. *Cancer Med* 2017;6:1409-23.
159. Bai X, Wang W, Zhao P, et al. LncRNA CRNDE acts as an oncogene in cervical cancer through sponging miR-183 to regulate CCNB1 expression. *Carcinogenesis* 2020;41:111-21.
160. Fan MJ, Zou YH, He PJ, et al. Long non-coding RNA SPRY4-IT1 promotes epithelial-mesenchymal transition of cervical cancer by regulating the miR-101-3p/ZEB1 axis. *Biosci Rep* 2019;39.
161. Chen WJ, Xiong L, Yang L, et al. Long Non-Coding RNA LINC01783 Promotes the Progression of Cervical Cancer by Sponging miR-199b-5p to Mediate GBP1 Expression. *Cancer Manag Res* 2020;12:363-73.
162. Zhao S, Yu M, Wang L. LncRNA miR503HG Regulates the Drug Resistance of Recurrent Cervical Squamous Cell Carcinoma Cells by Regulating miR-155/Caspase-3. *Cancer Manag Res* 2020;12:1579-85.
163. Chen X, Xiong D, Yang H, et al. Long noncoding RNA OPA-interacting protein 5 antisense transcript 1 upregulated SMAD3 expression to contribute to metastasis of cervical cancer by sponging miR-143-3p. *J Cell Physiol* 2019;234:5264-75.
164. Yang J, Jiang B, Hai J, et al. Long noncoding RNA opa-interacting protein 5 antisense transcript 1 promotes proliferation and invasion through elevating integrin alpha6 expression by sponging miR-143-3p in cervical cancer. *J Cell Biochem* 2019;120:907-16.
165. Ji N, Wang Y, Bao G, et al. LncRNA SNHG14 promotes the progression of cervical cancer by regulating miR-206/YWHAZ. *Pathol Res Pract* 2019;215:668-75.
166. Wu F, Zhou D, Cui Y, et al. Long non-coding RNA UCA1 modulates the glycolysis of cervical cancer cells by miR-493-5p/HK2. *Int J Clin Exp Pathol* 2018;11:3943-51.
167. Shao S, Wang C, Wang S, et al. LncRNA STXBP5-AS1 suppressed cervical cancer progression via targeting miR-96-5p/PTEN axis. *Biomed Pharmacother* 2019;117:109082.
168. Feng Y, Zou W, Hu C, et al. Modulation of CASC2/miR-21/PTEN pathway sensitizes cervical cancer to cisplatin. *Arch Biochem Biophys* 2017;623-624:20-30.
169. Zhu Y, Liu B, Zhang P, et al. LncRNA TUSC8 inhibits the invasion and migration of cervical cancer cells via miR-641/PTEN axis. *Cell Biol Int* 2019;43:781-8.
170. Fan Y, Sheng W, Meng Y, et al. LncRNA PTENP1 inhibits cervical cancer progression by suppressing miR-106b. *Artif Cells Nanomed Biotechnol* 2020;48:393-407.
171. Ou L, Xiang TY, Hao XY, et al. Reduced long non-coding RNA PTENP1 contributed to proliferation and invasion via miR-19b/MTUS1 axis in patients with cervical cancer. *Eur Rev Med Pharmacol Sci* 2020;24:4132-44.
172. Zhang X, Zhao X, Li Y, et al. Long noncoding RNA SOX21-AS1 promotes cervical cancer progression by competitively sponging miR-7/VDAC1. *J Cell Physiol* 2019;234:17494-504.
173. Yang J, Liang B, Hou S. TMPO-AS1 promotes cervical cancer progression by upregulating RAB14 via sponging miR-577. *J Gene Med* 2019;21:e3125.

174. Gang X, Yuan M, Zhang J. Long Non-Coding RNATMPO-AS1 Promotes Cervical Cancer Cell Proliferation, Migration, and Invasion by Regulating miR-143-3p/ZEB1 Axis. *Cancer Manag Res* 2020;12:1587-99.
175. Song H, Liu Y, Jin X, et al. Long non-coding RNA LINC01535 promotes cervical cancer progression via targeting the miR-214/EZH2 feedback loop. *J Cell Mol Med* 2019;23:6098-111.
176. Dou X, Zhou Q, Wen M, et al. Long Noncoding RNA FOXD2-AS1 Promotes the Malignancy of Cervical Cancer by Sponging MicroRNA-760 and Upregulating Hepatoma-Derived Growth Factor. *Front Pharmacol* 2019;10:1700.
177. Ma Z, Gu G, Pan W, et al. LncRNA PCAT6 Accelerates the Progression and Chemoresistance of Cervical Cancer Through Up-Regulating ZEB1 by Sponging miR-543. *Onco Targets Ther* 2020;13:1159-70.
178. Li Y, Wang H, Huang H. Long non-coding RNA MIR205HG function as a ceRNA to accelerate tumor growth and progression via sponging miR-122-5p in cervical cancer. *Biochem Biophys Res Commun* 2019;514:78-85.
179. Wu F, Sui Y, Wang Y, et al. Long Noncoding RNA SNHG7, a Molecular Sponge for microRNA-485, Promotes the Aggressive Behavior of Cervical Cancer by Regulating PAK4. *Onco Targets Ther* 2020;13:685-99.
180. Zhao D, Zhang H, Long J, et al. LncRNA SNHG7 Functions as an Oncogene in Cervical Cancer by Sponging miR-485-5p to Modulate JUND Expression. *Onco Targets Ther* 2020;13:1677-89.
181. Shen H, Wang L, Xiong J, et al. Long non-coding RNA CCAT1 promotes cervical cancer cell proliferation and invasion by regulating the miR-181a-5p/MMP14 axis. *Cell Cycle* 2019;18:1110-21.
182. Lu W, Wan X, Tao L, et al. Long Non-Coding RNA HULC Promotes Cervical Cancer Cell Proliferation, Migration and Invasion via miR-218/TPD52 Axis. *Onco Targets Ther* 2020;13:1109-18.
183. Xu J, Yang B, Wang L, et al. LncRNA BBOX1-AS1 upregulates HOXC6 expression through miR-361-3p and HuR to drive cervical cancer progression. *Cell Prolif* 2020;53:e12823.
184. Liu C, Tian X, Zhang J, et al. Long Non-coding RNA DLEU1 Promotes Proliferation and Invasion by Interacting With miR-381 and Enhancing HOXA13 Expression in Cervical Cancer. *Front Genet* 2018;9:629.
185. Wang Q, Ding J, Nan G, et al. LncRNA NOC2L-4.1 functions as a tumor oncogene in cervical cancer progression by regulating the miR-630/YAP1 pathway. *J Cell Biochem* 2019;120:16913-20.
186. Yang J, Hou S, Liang B. LINC00319 promotes migration, invasion and epithelial-mesenchymal transition process in cervical cancer by regulating miR-3127-5p/RPP25 axis. *In Vitro Cell Dev Biol Anim* 2020;56:145-53.
187. Xie F, Xie G, Sun Q. Long Noncoding RNA DLX6-AS1 Promotes the Progression in Cervical Cancer by Targeting miR-16-5p/ARPP19 Axis. *Cancer Biother Radiopharm* 2020;35:129-36.
188. Zhu H, Zheng T, Yu J, et al. LncRNA XIST accelerates cervical cancer progression via upregulating Fus through competitively binding with miR-200a. *Biomed Pharmacother* 2018;105:789-97.
189. Liu X, Xie S, Zhang J, et al. Long Noncoding RNA XIST Contributes to Cervical Cancer Development Through Targeting miR-889-3p/SIX1 Axis. *Cancer Biother Radiopharm* 2020;35:640-9.
190. Chen X, Xiong D, Ye L, et al. Up-regulated lncRNA XIST contributes to progression of cervical cancer via regulating miR-140-5p and ORC1. *Cancer Cell Int* 2019;19:45.
191. Song L, Wang L, Pan X, et al. lncRNA OIP5-AS1 targets ROCK1 to promote cell proliferation and inhibit cell apoptosis through a mechanism involving miR-143-3p in cervical cancer. *Braz J Med Biol Res* 2020;53:e8883.
192. Wang L, Zhong Y, Yang B, et al. LINC00958 facilitates cervical cancer cell proliferation and metastasis by sponging miR-625-5p to upregulate LRRC8E expression. *J Cell Biochem* 2020;121:2500-9.
193. Zhao H, Zheng GH, Li GC, et al. Long noncoding RNA LINC00958 regulates cell sensitivity to radiotherapy through RRM2 by binding to microRNA-5095 in cervical cancer. *J Cell Physiol* 2019;234:23349-59.
194. Li H, Hong J, Wijayakulathilaka W. Long non-coding RNA SNHG4 promotes cervical cancer progression through regulating c-Met via targeting miR-148a-3p. *Cell Cycle* 2019;18:3313-24.
195. Guo Q, Zhang Q, Lu L, et al. Long noncoding RNA RUSC1-AS1 promotes tumorigenesis in cervical cancer by acting as a competing endogenous RNA of microRNA-744 and

- consequently increasing Bcl-2 expression. *Cell Cycle* 2020; 19: 1222-35.
196. Hou A, Zhang Y, Zheng Y, et al. LncRNA terminal differentiation-induced ncRNA (TINCR) sponges miR-302 to upregulate cyclin D1 in cervical squamous cell carcinoma (CSCC). *Hum Cell* 2019;32:515-21.
197. Zhao H, Hu GM, Wang WL, et al. LncRNA TDRG1 functions as an oncogene in cervical cancer through sponging miR-330-5p to modulate ELK1 expression. *Eur Rev Med Pharmacol Sci* 2019;23:7295-306.
198. Guo M, Lin B, Li G, et al. LncRNA TDRG1 promotes the proliferation, migration, and invasion of cervical cancer cells by sponging miR-214-5p to target SOX4. *J Recept Signal Transduct Res* 2020;40:281-93.
199. Feng Y, Ma J, Fan H, et al. TNF-alpha-induced lncRNA LOC105374902 promotes the malignant behavior of cervical cancer cells by acting as a sponge of miR-1285-3p. *Biochem Biophys Res Commun* 2019;513:56-63.
200. Barr JA, Hayes KE, Brownmiller T, et al. Long non-coding RNA FAM83H-AS1 is regulated by human papillomavirus 16 E6 independently of p53 in cervical cancer cells. *Sci Rep* 2019;9:3662.
201. Sharma S, Munger K. Expression of the Long Noncoding RNA DINO in Human Papillomavirus-Positive Cervical Cancer Cells Reactivates the Dormant TP53 Tumor Suppressor through ATM/CHK2 Signaling. *mBio* 2020;11:202. Zhang M, Song Y, Zhai F. ARFHPV E7 oncogene, lncRNA HOTAIR, miR-331-3p and its target, NRP2, form a negative feedback loop to regulate the apoptosis in the tumorigenesis in HPV positive cervical cancer. *J Cell Biochem* 2018;119: 4397-407.
203. Yang L, Yi K, Wang H, et al. Comprehensive analysis of lncRNAs microarray profile and mRNA-lncRNA co-expression in oncogenic HPV-positive cervical cancer cell lines. *Oncotarget* 2016;7:49917-29.
204. Sharma S, Munger K. Expression of the cervical carcinoma expressed PCNA regulatory (CCEPR) long noncoding RNA is driven by the human papillomavirus E6 protein and modulates cell proliferation independent of PCNA. *Virology* 2018; 518:8-13.
205. Iancu IV, Anton G, Botezatu A, et al. LINC01101 and LINC00277 expression levels as novel factors in HPV-induced cervical neoplasia. *J Cell Mol Med* 2017;21:3787-94.
206. He H, Liu X, Liu Y, et al. Human Papillomavirus E6/E7 and Long Noncoding RNA TMPOP2 Mutually Upregulated Gene Expression in Cervical Cancer Cells. *J Virol* 2019;93.
207. Zhou D, Wu F, Cui Y, et al. Long non-coding RNA-OIS1 inhibits HPV-positive, but not HPV-negative cervical squamous cell carcinoma by upregulating MTK-1. *Oncol Lett* 2019;17:2923-30.