



International Journal of Molecular and Cellular Medicine p-ISSN: 2251-9637 o-ISSN: 2251-9645 Cellular and Molecular Biology Research Center

Potential Otubain1 Inhibitor, an Approach for a Treatment against Breast Cancer

Andrea Muñoz-Ayala, D Victor G. García-González, D Angel Pulido-Capiz, D Brenda Chimal-Vega, D Ruth A. García-Villarreal, D José L. Vique-Sánchez*

School of Medicine Campus Mexicali, Autonomous University of Baja California, Mexicali, 21000, BC, México.

Article type:	ABSTRACT
Original Article	The develop of new anticancer drug continues worldwide and one of the new therapeutic
	targets to reach it is Otubain 1 (OTUB1), since OTUB1 has functions related to prognosis in
	a variety of tumors and is strongly related to tumor proliferation, migration, and apoptosis by
	their functions on deubiquitinating. This study uses OTUB1's active site to develop a specific
	pharmacological treatment to regulate the OTUB1 functions. The aim of this research was to
	evaluate the effects of ten compounds (OT1 - OT10), that previously were selected by
	molecular docking to develop a new anticancer drug to decrease the OTUB1 functions in the
	cancer processes. We evaluated the cytotoxic effect of OT1 – OT10 compounds on MCF-7,
	BT474 and MDA-MB 231 cells by MTT assay, and we determined characteristics of
	apoptosis by western blot analysis. Then, the best compound (OT5) was analyzed by
	molecular docking, molecular dynamics and theoretical toxicity for describing the
	interactions of OT5 compound with the OTUB1's active site. We proposed that the OT5
	compound has a high probability to be selective against OTUB1, with an apoptosis
	(regulating caspase-8) and cytotoxic effect on some cancer lines; IC50 for MCF-7: 97 μM
Received:	and MDA-MB 231:147 $\mu\text{M},$ as well as we described that this compound could have specific
2024.07.29	interactions in the catalytic domain of OTUB1, modifying this protein's activity, decreasing
Revised:	the OTUB1 functions, and probably safe for humans. These results show the high potential
2024.09.07	of this compound for promoting the development of this compound as a new drug against
Accepted:	cancer.
2024.09.08	Keywords: Apoptosis, caspase-8, MCF-7, MDA-MB 231, drug

Cite this article: Muñoz-Ayala A, *et al.* Potential Otubain1 Inhibitor, an Approach for a Treatment against Breast Cancer. *International Journal of Molecular and Cellular Medicine*. 2024; 13(4):350-360. DOI: 10.22088/IJMCM.BUMS.13.4.350

*Corresponding: José L. Vique-Sánchez Address: School of Medicine Campus Mexicali, Autonomous University of Baja California, Mexicali, 21000, BC, México. E-mail: jvique@uabc.edu.mx



Publisher: Babol University of Medical Sciences

This work is published as an open access article distributed under the terms of the Creative Commons Attribution 4.0 License (http://creativecommons.org/licenses/by-nc/4). Non-commercial uses of the work are permitted, provided the original work is properly cited.

Introduction

The development of a new anticancer drug continues worldwide, we previously reported ten compounds (OT1-OT10) selected according to pharmacological characteristics and Lipinsky's rules, and this could interact with the Otubain 1 (OTUB1) (1), this protein is a promising therapeutic target against cancer, due to the functional complexity of OTUB1 in autophagy and immune response, researchers need to be more careful when evaluating the prognostic value of OTUB1 in different cancers and for poor prognosis (2), it is a deubiquitinating enzyme that belongs to the OTU family of cysteine proteases (3.4), OTUB1 has multiple functions, this enzyme can decrease the ubiquitination state of a protein by removing ubiquitin (canonical activity) or prevent ubiquitin conjugation (non-canonical activity) (5,6). Examples of regulating processes are immune response, factors that regulate interferon and NF-KB (7,8), and regulating many cancerassociated signaling pathways including Mitogen-activated protein kinase (MAPK), Epithelialmesenchymal transition (EMT), Forkhead box protein M1 (FOXM1), Ras homologous A (RHOA), mechanistic target of rapamycin complex 1 (mTORC1), c-Maf, cellular inhibitor of apoptosis-1 (c-IAP1), and P53 to promote tumor cell survival, proliferation, invasiveness, prognosis, and therapeutic resistance (9-13). Furthermore, OTUB1 has been reported to be present in many cancer tissues and has been negatively associated with the poor condition of cancer patients, because OTUB1 is highly expressed in many cancers and is known to inhibit the degradation of molecules that regulate cell proliferation due to the abovementioned (5, 9, 14, 15). Otherwise, there are studies over the inhibition of the OTUB1 (2.9), like in the c- Maf axis by extracts that induces myeloma cell apoptosis, due to OTUB1 stabilizes c-Maf by preventing its polyubiquitination and enhances c-Maf transcriptional activity (13), as well as being identified that the expression of OTUB1 is very important for melanoma-cell survival (9).

On the other hand, OTUB1 could regulate apoptosis in cancer cell, and the regulation of the TNF-related apoptosis-inducing ligand (TRAIL) expression (9), thus for any development of anticancer drug is important to generate apoptosis (16), and the extrinsic apoptotic pathway could be activated by activating the members of the tumor necrosis factor alpha (TNF α) receptor super-family, which are usually called the death receptors (i.e., Fas, also called CD95/Apo-1, DR4 and DR5, TNF receptors, TRAIL) (17). OTUB1 is related with these receptors, for example, once that the level of TRAIL is increased, the oligomerization leading to the formation of pro-caspase-8 and activating the death-inducing signaling complex is identified (9). Thus, OTUB1 could regulate TRIAL, since it has been identified that inhibiting OTUB1 increases the function of TRAIL, which leads to the activation of caspase-8, which is related to regulating cell death, through the intrinsic or extrinsic pathway and that apoptosis-initiating caspases (2,8,9, and 10), and thus avoid necroptosis (18,19). The *caspase-8* may be at different levels of expression according to the type of cancer (20,21), for example in MCF-7 and Esophageal carcinoma (ESCA) with overexpression of caspase-8, in addition, are studies where it is reported that in MCF-7 the caspase-8 activation is an initiating event in DHAinduced apoptotic cell death (22), even other studies relating to an OTUB1 inhibitory effect (13,23), both studies using OTUB1 as therapeutic target (12) with promising results that demonstrated that OTUB1 might be a good therapeutic target.

In this study, we evaluated ten compounds reported (1), that could interact near the catalytic site of OTUB1 (Asp88, Cys91 and His265) (3,5,6), these compounds could decrease the deubiquitinating function

of OTUB1 for developing a cancer drug (5,9,11,12,24,25), by molecular docking, molecular dynamics simulation, cytotoxicity, MTT, and western blot assays.

Materials and methods

Synthetic compounds

We use the compounds that previously reported OT1 – OT10 (1), these compounds were bought in Chembridge Corp. (26) (ID- Chembridge: OT1-7988434, OT2-7501037, OT3-5194547, OT4-7986805, OT5-7013154, OT6-7699311, OT7-5634266, OT8-7987933, OT9-5181121, and OT10-7661200). Purity of the compound reported by the manufacturer was greater than 95% checked by LC/MS, the stock solutions for compounds were prepared in DMSO (1-10 mM).

Cell Viability Assay

Cell culture reagents were purchased from Thermo-Fisher (Carlsbad, CA, U.S.A.), tissue culture plates and other plastic materials were obtained from Corning Inc. Cell lines BT-474, MDA-MB-231, and MCF-7 were obtained from American Type Culture Collection (ATCC). Cells were cultured following the provider's recommendations, and the cytotoxic effect was determined by MTT assay as described previously; MTT was obtained from Merck (Darmstadt, Germany) (27). Briefly, the cell lines were seeded (8 - 20 x 10³ cells per well). First, we used 100 µM concentration of each compound (OT1 - OT10) in 0.4 % DMSO incubated for 48 h to determine the cytotoxic effect, and if some compound had cytotoxic effect was determined the IC50 using concentrations between 10, 20, 50, 100, and 170 µM in 0.4 % DMSO incubated for 24 h. After 24 h, the medium was aspirated, 20 µL of 5 mg/mL MTT reagent was added to each well and incubated for 4 h. The excessive liquid was gently removed and 100 μ L of DMSO (0.4 %) were added to the wells. Then, the plate was read at 570 nm using the microtiter plate reader (µQuant, Bio-Tek Instrument, USA). The assays were performed in triplicate in three independent studies and the percentage of viability was calculated according to the formula: % viability= [mean Optical Density (O.D.) treated cells×100]/ (mean O.D. control cells). The concentration leading to 50% inhibition viability (IC50) was calculated by non-linear regression analysis (percentage of viability vs. log concentration) with GraphPad Prism software ver. 8.0 software (GraphPad, CA, USA).

Western blot to identify caspase-8 in Apoptosis effect

MCF-7 cells were seeded into a 6-well plate (400,000 cells per well) until they reached 90-95% confluence. DMEM supplemented medium was replaced with DMEM experimental medium for 2h. Later, cells were incubated with DMSO, 90 μ M and 170 μ M of OT5 compound for 24 h. Cells were washed with PBS and lysed with lysis buffer (150 mM NaCl, 10 mM Tris, pH 7.4, 1% Triton X- 100, 0.5% NP40, EDTA, 0.2 mM sodium orthovana-date, 0.3 μ M aprotinin,130 μ M bestatin, 14 μ M E- 64, and 1 μ M leupeptin). Proteins were separated by 10% SDS-PAGE and transferred to PVDF membranes. Membranes were blocked at 37 °C for 1h in TBS/0.1% tween 5% low fat milk. Anti-pro-caspase-8 (sc-56070) and anti- β actin (sc-47778) primary antibodies (Santa Cruz Biotechnology, USA). Anti-mouse secondary antibodies coupled to Horseradish peroxidase from Thermo-Fisher. Chemiluminescence detection was performed using Immobilon Western kit (Millipore, MA, U.S.A.) and X-ray film for some blots, for others Bio-Rad ChemiDoc XRS+ was used and a digital image was obtained.

Description of the molecular interaction of cytotoxic compounds with OTUB1 by docking

Otubain 1 structure from the Protein Data Bank (28) (OTUB1, PDB code 3VON) was used for molecular interaction using Molecular Operating Environment (MOE) following procedures previously reported (29-31) and Protein-Ligand Interaction Profiler (29,32,33,34) to describe the interaction between compounds and OTUB1.

Molecular dynamics simulations of cytotoxic compounds with OTUB1

The complex of OTUB1 with compound OT5 generated by docking was used for molecular dynamics simulation (MD). It was performed using Dynamics – MOE and OPLS-AA forcefield. At the beginning, in two consecutives steps of 100 ps reaching a temperature of 300 K and a pressure of 1 bar. Finally, a 10 ns MD was carried out. Furthermore, the interactions were analyzed.

Statistical analysis

Data's results were expressed as mean \pm standard deviation (SD) of minimum three independent experiments. Averages and standard deviations were calculated in Excel (365 Microsoft, USA). IC50 for cytotoxicity of compounds were calculated using GraphPad Prism 8 Software (La Jolla, CA, USA).

Results

Cytotoxic effect of OT1 – OT10 compounds

In this study, we tested ten compounds (OT1 – OT10) that we reported previously to interact with OTUB1 (1). After the initial concentration at 100 μ M of each compound, we determined only one compound (OT5 compound) with cytotoxic effect at MCF-7 (ER+ cells), MDA-MB 231 (TNBC cells), and BT-474 (HER2+ cells) (Figure 1).



Fig.1. BT-474 cell line after 48 h of incubation. A) Group control with DMSO as vehicle, B) Group of OT5 compound with 100 μM. **IC50 of OT5 compound**

We determined the IC50 using the results of the percentage of viability by MTT. Concentrations of OT5 compound between 10, 20, 50, 100, and 170 μ M were used (Table 1). With these results, we determined an IC50 of 97.2 ± 1.13 μ M (MCF-7 cells) and 147.4 ± 1.09 μ M (MDA-MB-231 cells) for 24 h of incubation, using the Graph Pad Prism 8.

Table 1. Concentrations used of OT5 compound and viability percentage in <i>in vitro</i> cultures of MCF-7 and MDA-MB-231 cells.						
Concentration (µM)	Viability percentage of MCF-7	Viability percentage of MDA-MB-231				
0	100	100				
10	90	91				
20	88	84				
50	80	76				
100	48	69				
170	22	39				

Apoptosis Assay

We made WB to determine pro-Caspase-8, and we identified that caspase-8 increased by OT5 compound with 90 μ M (near of IC50), and with 170 μ M was identified cleaved caspase-8 (Figure 2) after 24 h of incubation using MCF-7 cells. These results were related with the apoptosis effect by OT5 compound.



Fig.2. Western Blot assay for pro-caspase-8 and β -actin after 24 h of incubation in MCF-7 cells. Line 1: control with DMSO, line 2: 90 μ M of OT5 compound and Line 3: 170 μ M of OT5 compound.

Interaction of OT5 compound with OTUB1

According to the interactions report (1), OT5 compound could have interaction in the region between Lys84, Asp88, Cys91, and His265 amino acids (Figure 3). The main interactions are in Asn45, Lys84, Arg86, Pro87, Asp88, Gln165, Cys212, Arg262, Pro263, Gly264, and His265 in OTUB1. This reported particularly a greater interaction with Arg86, Pro263 and His265 (mainly hydrogen bonding interaction in the conformers analyzed), with these results, the catalytic site could be blocked, which is essential for the deubiquitinating functions of OTUB1. The details of the interaction between OTUB1 with conformers of each compound are shown in the previous report (1).

Molecular dynamics simulation

To study in more detail the complex OTUB1-OT5, 10 ns of molecular dynamics simulation was performed. The root mean square deviation (RMSD) was analyzed, the data showed that the inhibitor formed

a stable complex almost during simulation time, after 10 ns were stable in the main interactions, notably the interactions with Lys84, Arg86 and Pro286 (Figure 4).



Fig.3. OTUB1 (orange) with 22 conformers of OT5 compound (white) in the active site.



Fig.4. Interactions at 0, 5 and 10 ns frame from molecular dynamics simulation between OT5 compound and OTUB1.

The overall RMSD after 10 ns was 1.518 A, and specifically in the amino acids with the main interactions, the RMSD were between 0.627 to 1.287 A (Table 2).

Table 2. Main interactions and RMSD of OT5 with OTUB1 between $0 - 10$ ns.							
Time	Ligand	Receptor (Residues in OTUB1)	Interaction	Distance	RMSD		
0 ns	Ν	PRO263	H-donor	3.16	-		
	6-ring	LYS84	pi-cation	3.77	-		
5 ns	Ν	PRO263	H-donor	2.8	0.800		
	6-ring	LYS84	pi-cation	3.6	0.886		
		ARG86			0.795		
10 ns	Ν	PRO263	H-donor	2.77	1.207		
		LYS84			1.287		
	6-ring	ARG86	pi-H	4.17	0.627		
	6-ring	ARG86	pi-H	4.16			

Discussion

In this study, we evaluated ten compounds directed to the OTUB1's active site to develop a specific pharmacological treatment to regulate the OTUB1 functions (mainly as deubiquitinating), someone of these related to cancer progression, and immune response (35), that have increased the development of specific inhibitors; for example, the 61 compounds with great results over lung cancer (36). Thus, our study focuses on determining the OTUB1 inhibitors effect over breast cancer. We determined that one compound (OT5) has results enough to continue the development of this anticancer drug. We propose that the OT5 compound could be regulating the deubiquitylating function of OTUB1 (3, 35).

The *in vitro* results showed that OT5 compound has a cytotoxic effect and an apoptosis effect related to regulate the caspase-8 determined by western blot. We determined the cytotoxic effect on different cell lines (MCF-7, MDA-MB 231, and BT474) using 100 µM of OT5 compound, and it generated signals of apoptosis and death cell, by loss of monolayer and changes morphological in cells (Figure 1) (9,37,38,39). As we had mentioned, it is important to generate apoptosis in the studies to develop new anticancer drugs, and the caspase-8 is a key protein of crosstalk signaling in a variety of cancers (40), and one of them, an anticancer drug reported an increase of caspase-8 level after 24 to 72 h incubation (41), and MCF-7 cells are related to the caspase-8 level increase with the apoptosis (19,42,43,44). Our results showed that OT5 compound has an effect over caspase-8 that could justify the apoptosis effect by this compound after 24 h (Figure 2). In this way, OT5 compound could be promoting the apoptosis, and this effect is related to an report in melanoma, where in this study was generating/promoting the apoptosis by decreasing OTUB1 functions, and it increased the expression of TRAIL and this increases apoptosis by caspase-8 (9), and it could be related to our results (Figure 2), where with a low concentration of OT5 compound increased the concentration of caspase-8 and identified as death cell, this effect could be related to an apoptotic effect of an anticancer compound, where there is an increase in the pro-caspase-8 (55 kDa) and active-caspase-8 in some cancer lines (9,45), for example in human bladder cancer (T24 cells) (46), colon cancer cells (DLDI) (47) and in MDA-MB 231 cells, so, in these studies, the caspase-8 concentration was maintained or increased, and all of them were related to apoptosis (38). On the other hand, OTUB1 has an important relation with c-MYC. Consequently, OTUB1 increases c-MYC protein levels, which increase glycolysis and tumor growth, and this is related in breast cancer cells (MCF-7 and MDA-MB 231) (48). Therefore, the activation of c-MYC contributes to the progression of multiple types of cancer (35).

To justify the cytotoxic and apoptosis effect, the *in silico* results of OT5 compounds showed that the main interactions between OT5 and OTUB1 are with amino acids in a region considered as potential therapeutic target (active site) (3,5,6) (Figures 3 and 4) that could regulate the c- Maf axis to generate apoptosis (12) (13). The results of the molecular dynamics simulation showed that the interactions between the OT5 compound with the active site (Figure 5) are stable, and we describe the main interactions of OT5 compound with OTUB1, the interactions are very similar either in the molecular docking and in the molecular dynamics simulation; in the OTUB1's active site (Table 2), these interactions could block the accessibility to the catalytic site (Asp88, Cys91 and His265) and decreases some function of OTUB1. In this way, an example of OTUB1 inhibitor is the 61 compound tested in lung cancer, it interacts near the OTUB1's active site (36). Thus, according to some functions of OTUB1, it is reported that Asp88 and Cys91 are necessary to OTUB1/c-Maf axis (12-13), and the OT5 compound could generate a disfunction of OTUB1, due to this region essential for the deubiquitinating functions of OTUB1. Thus, we related our results with other studies where OTUB1 was inhibited, it reduced their functions and the caspase-8 was activated, and all those studies with an effect on the control of apoptosis (18, 19, 22, 40, 41, 42, 43, 44).



Fig.5. OTUB1 structure surface, the results of the molecular dynamics simulation after 10 ns. A) It shows the main amino acids' interactions between the OT5 compound and the active site are stables. B) It shows the region of active site (white) and the colocation of OT5 compound (pink), for blocking the access to active site.

Moreover, we expand the theoretical toxicity studies (49-51), and OT5 compound toxicity results are potentially secure to be used in humans, as well as to propose the OT5 compound for pharmacophore modeling studies (52-54), and reach to improve the anticancer effect of new molecules.

As conclusion, we propose the OT5 compound with high probability to be selective against OTUB1, with an apoptosis and cytotoxic effect on some cancer lines. We describe that this compound could have specific interactions in the catalytic domain of OTUB1, modifying this protein's activity, decreasing the OTUB1 functions, and probably safe for humans. Therefore, this study demonstrates the potential of OT5

compound as anticancer drug, and other studies should be performed to continue the development of this new anticancer drug.

Acknowledgments

The authors are very grateful to SNII-CONAHCYT, FMM-UABC and Dr. José Manuel Avendaño Reyes.

References

1. Galindo-Hernández O, García-Salazar LA, García-González VG, et al. Potential inhibitors of the OTUB1 catalytic site to develop an anti-cancer drug using in-silico approaches. Reports Biochem Mol Biol 2023; 11:684–93.

2. Zhang W, Qiu W. OTUB1 recruits tumor infiltrating lymphocytes and is a prognostic marker in digestive cancers. Front Mol Biosci 2020; 7:212.

3. Balakirev MY, Tcherniuk SO, Jaquinod M, et al. Otubains: a new family of cysteine proteases in the ubiquitin pathway. EMBO Rep 2003; 4:517–22.

4. Wiener R, DiBello AT, Lombardi PM, et al. E2 ubiquitin-conjugating enzymes regulate the deubiquitinating activity of OTUB1. Nat Struct Mol Biol 2013; 20:1033–9.

5. Saldana M, VanderVorst K, Berg AL, et al. Otubain 1: a non-canonical deubiquitinase with an emerging role in cancer. Endocr Relat Cancer 2019; 26:R1–14.

6. Wu Q, Huang Y, Gu L, et al. OTUB1 stabilizes mismatch repair protein MSH2 by blocking ubiquitination. J Biol Chem 2021; 296: 100466.

7. Jahan AS, Biquand E, Muñoz-Moreno R, et al. OTUB1 is a key regulator of RIG-I-dependent immune signaling and is targeted for proteasomal degradation by influenza A NS1. Cell Rep 2020; 30:1570–84.

8. Mulas F, Wang X, Song S, et al. The deubiquitinase OTUB1 augments NF-κB-dependent immune responses in dendritic cells in infection and inflammation by stabilizing UBC13. Cell Mol Immunol 2021; 18:1512–27.

9. Lee B-S, Kang SU, Huang M, et al. OTUB1 knockdown promotes apoptosis in melanoma cells by upregulating TRAIL expression. BMB Rep 2021; 54:608–13.

10. Sun X-X, Dai M-S. Deubiquitinating enzyme regulation of the p53 pathway: a lesson from Otub1. World J Biol Chem 2014; 5:75–84.

11. Sun X-X, Challagundla KB, Dai M-S. Positive regulation of p53 stability and activity by the deubiquitinating enzyme Otubain 1. EMBO J 2012; 31:576–92.

12. Xu Y, Xu M, Tong J, et al. Targeting the Otub1/c-Maf axis for the treatment of multiple myeloma. Blood 2021; 137:1478–90.

13. Sun T, Xu Y, Xu Z, et al. Inhibition of the Otub1/c-Maf axis by the herbal acevaltrate induces myeloma cell apoptosis. Cell Commun Signal 2021; 19:24.

14. Liao Y, Wu N, Wang K, et al. OTUB1 promotes progression and proliferation of prostate cancer via deubiquitinating and stabling cyclin E1. Front Cell Dev Biol 2021; 8.

15. Liu T, Jiang L, Tavana O, Gu W. The deubiquitylase OTUB1 mediates ferroptosis via stabilization of SLC7A11. Cancer Res 2019; 79:1913–24.

16. Ghotaslou A, Samii A, Boustani H, et al. AMG-232, a new inhibitor of MDM-2, enhances doxorubicin efficiency in pre-B acute lymphoblastic leukemia cells. Reports Biochem Mol Biol 2022; 11:111–24.

17. Khosravi-Far R. Death receptor signals to the mitochondria. Cancer Biol Ther 2004; 3:1051-7.

Boice A, Bouchier-Hayes L. Targeting apoptotic caspases in cancer. Biochim Biophys Acta - Mol Cell Res 2020; 1867:118688.
Jelínek M, Balušíková K, Schmiedlová M, et al. The role of individual caspases in cell death induction by taxanes in breast cancer cells. Cancer Cell Int 2015; 15:8.

20. Stupack DG. Caspase-8 as a therapeutic target in cancer. Cancer Lett 2013; 332:133-40.

21. Uhlén M, Fagerberg L, Hallström BM, et al. Tissue-based map of the human proteome. Science 2015; 347.

22. Kang KS, Wang P, Yamabe N, et al. Docosahexaenoic acid induces apoptosis in MCF-7 cells in vitro and in vivo via reactive oxygen species formation and caspase 8 activation. PLoS One 2010; 5:e10296.

23. Xu Y, Sun T, Zeng K, et al. Anti-bacterial and anti-viral nanchangmycin displays anti-myeloma activity by targeting Otub1 and c-Maf. Cell Death Dis 2020; 11:818.

24. Doherty LM, Mills CE, Boswell SA, et al. Integrating multi-omics data reveals function and therapeutic potential of deubiquitinating enzymes. Elife 2022; 11:e72879

25. Sivakumar D, Stein M. Binding of SARS-CoV covalent non-covalent inhibitors to the SARS-CoV-2 papain-like protease and ovarian tumor domain deubiquitinases. Biomolecules 2021; 11:802.

26. ChemBridge Corp. EXPRESS-Pick Stock. Chemical structures available in hit2lead.com: https:// www. hit2lead.com/ search.asp? db= SC& SearchPage=id. Accessed 2024.

27. Mosmann T. Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. J Immunol Methods 1983; 65:55-63.

28. RCSB PDB. Crystal structure of the ubiquitin protease [Online Database]. Available from: <u>https://www.rcsb.org/3d-view/3VON</u>. Accessed October 15, 2023.

29. Benítez-Cardoza CG, Vique-Sánchez JL. Potential inhibitors of the interaction between ACE2 and SARS-CoV-2 (RBD), to develop a drug. Life Sci 2020; 256:117970.

30. Halgren TA. Merck molecular force field. I. Basis, form, scope, parameterization, and performance of MMFF94. J Comput Chem 1996; 17:490-519.

31. Vique-Sánchez JL. Potential inhibitors interacting in Neuropilin-1 to develop an adjuvant drug against COVID-19, by molecular docking. Bioorg Med Chem 2021; 33:116040.

32. Soga S, Shirai H, Kobori M, et al. Use of amino acid composition to predict ligand-binding sites. J Chem Inf Model 2007; 47:400-6.

33. Adasme MF, Linnemann KL, Bolz SN, et al. PLIP 2021: expanding the scope of the protein–ligand interaction profiler to DNA and RNA. Nucleic Acids Res 2021; 49:W530–4.

34. Salentin S, Schreiber S, Haupt VJ, et al. PLIP: fully automated protein–ligand interaction profiler. Nucleic Acids Res 2015; 43:W443-7.

35. Wu M, Sun L, Song T. OTUB1-mediated inhibition of ubiquitination: a growing list of effectors, multiplex mechanisms, and versatile functions. Front Mol Biosci 2024; 10:1261273..

36. Tan L, Shan H, Han C, et al. Discovery of potent OTUB1/USP8 dual inhibitors targeting proteostasis in non-small-cell lung cancer. J Med Chem 2022; 65:13645-59.

37. Balvan J, Krizova A, Gumulec J, et al. Multimodal holographic microscopy: distinction between apoptosis and oncosis. PLoS One 2015; 10:e0121674.

38. Guvenir Celik E, Eroglu O, et al. Combined treatment with ruxolitinib and MK-2206 inhibits the JAK2/STAT5 and PI3K/AKT pathways via apoptosis in MDA-MB-231 breast cancer cell line. Mol Biol Rep 2023; 50:319-29.

39. Mirsanei JS, Nazari M, Shabani R, et al. Does gold-silver core-shell nanostructure with alginate coating induce apoptosis in human lymphoblastic tumoral (Jurkat) cell line? Rep Biochem Mol Biol 2023; 12:233-40.

40. Chai J, Lei Y, Xiang X, et al. High expression of caspase-8 as a predictive factor of poor prognosis in patients with esophageal cancer. Cancer Med 2023; 12:7651-66.

41. Sharifi S, Barar J, Hejazi MS, et al. Doxorubicin changes Bax/Bcl-xL ratio, caspase-8 and -9 in breast cancer cells. Adv Pharm Bull 2015; 5:351-9.

42.Gui C, Zhang C, Xiong X, et al. Total flavone extract from Ampelopsis megalophylla induces apoptosis in the MCF-7 cell line. Int J Oncol 2021; 58:409-18.

43. Jamalzadeh L, Ghafoori H, Aghamaali M, et al. Induction of apoptosis in human breast cancer MCF-7 cells by a semi-synthetic derivative of artemisinin: a caspase-related mechanism. Iran J Biotechnol 2017; 15:157-65.

44. Anguiano-Sevilla LA, Lugo-Cervantes E, Ordaz-Pichardo C, et al. Apoptosis induction of Agave lechuguilla Torrey extract on human lung adenocarcinoma cells (SK-LU-1). Int J Mol Sci 2018; 19:3765.

45. Kuo HM, Tseng CC, Chen NF, et al. MSP-4, an antimicrobial peptide, induces apoptosis via activation of extrinsic Fas/FasLand intrinsic mitochondria-mediated pathways in one osteosarcoma cell line. Mar Drugs 2018; 16:8.

46. Park C, Choi EO, Hwangbo H, et al. Induction of apoptotic cell death in human bladder cancer cells by ethanol extract of Zanthoxylum schinifolium leaf, through ROS-dependent inactivation of the PI3K/Akt signaling pathway. Nutr Res Pract 2022; 16:330.

47.Hu S, Yao X, Hao Y, Pan A, Zhou X. 8-Gingerol regulates colorectal cancer cell proliferation and migration through the EGFR/STAT/ERK pathway. Int J Oncol 2020; 56:390-397

48. Han X, Ren C, Lu C, Qiao P, et al. Deubiquitination of MYC by OTUB1 contributes to HK2 mediated glycolysis and breast tumorigenesis. Cell Death Differ 2022; 29:1864–73.

49. Daina A, Michielin O, Zoete V. SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules. Sci Rep 2017; 7:42717.

50. Banerjee P, Eckert AO, Schrey AK, et al. ProTox-II: a webserver for the prediction of toxicity of chemicals. Nucleic Acids Res 2018; 46:W257–63.

51. Dong J, Wang NN, Yao ZJ, et al. ADMETlab: a platform for systematic ADMET evaluation based on a comprehensively collected ADMET database. J Cheminform 2018; 10:29.

52.Khalid I, Jafar TH, Unar A, et al. In-Silico Identification of Anticancer Compounds; Ligand-Based Pharmacophore Approach against EGFR Involved in Breast Cancer. Adv Breast Cancer Res 2021; 10:120–32.

53. Kumar S, Ali I, Abbas F, et al. In-silico design, pharmacophore-based screening, and molecular docking studies reveal that benzimidazole-1,2,3-triazole hybrids as novel EGFR inhibitors targeting lung cancer. J Biomol Struct Dyn 2024:42:9416-9438.

54.Shaikh GM, Murahari M, Thakur S, et al. Studies on ligand-based pharmacophore modeling approach in identifying potent future EGFR inhibitors. J Mol Graph Model 2022; 112:108114.