

Molecular detection of Integron genes and pattern of antibiotic resistance in *Pseudomonas aeruginosa* strains isolated from intensive care unit, Shahid Beheshti Hospital, North of Iran

Fatemeh Moradian kouchaksaraei¹, Elaheh Ferdosi Shahandashti², Zahra Molana³, Masomeh Moradian kouchaksaraei⁴, Fariba Asgharpour⁵, Ali Mojtabaei⁶, Ramazan Rajabnia^{3*}

1. Department of Microbiology Science and Research Branch Islamic Azad University, Guilan, Iran.
2. Department of Microbiology and Immunology, Babol University of Medical Sciences, Babol, Iran.
3. Infectious Diseases and Tropical Medicine Research Center, Babol University of Medical Sciences, Babol, Iran.
4. Department of Microbiology Science and Research Branch Islamic Azad University, Fars, Iran.
5. Faculty of Para-Medicine; Babol University of Medical Sciences, Babol, Iran.
6. Department of Microbiology, Faculty of Medicine, Guilan University Campus, Rasht, Iran

Submitted 23 January 2013; Accepted 11 February 2013

Pseudomonas aeruginosa is one of the most important pathogens that causes nosocomial infections and shows high level of antibiotic resistance. Integrons are one of the transposable elements in bacteria and their role in antibiotic resistance has been well demonstrated. The aim of this study was a molecular characterization of the integron genes and the determination of the resistance or sensitivity pattern to ceftizoxime, ceftazidime, cephalexin, amikacin, ofloxacin, imipenem, cefepime, ticarcillin, gentamicin, ciprofloxacin, cefazolin and ceftriaxone antibiotics in *P. aeruginosa* strains isolated from Intensive Care Units (ICU), Shahid Beheshti Hospital, North of Iran. This cross-sectional study was performed from 2011 to 2012. Totally, fifty four *P. aeruginosa* strains were isolated from ICU at Shahid-Beheshti hospital, Babol, North of Iran. The bacteria were diagnosed based on mobility, pigment production, growth in 42° C, oxidase and catalase tests. PCR analysis was carried out to detect integron genes using hep 35 and hep 36 primers. Also, disk diffusion method was performed to evaluate antibiotic susceptibility of the bacteria using ceftizoxime, ceftazidime, cephalexin, amikacin, ofloxacin, imipenem, cefepime, ticarcillin, gentamicin, ciprofloxacin, cefazolin and ceftriaxone antibacterial reagents. This study revealed that 20 (37%) *P. aeruginosa* isolates had integron genes. The antibiotic susceptibility test showed that 52 (96.3%) of the isolates were multidrug-resistant. 12 out of 54 isolated bacteria were resistant to all antibiotics tested. All bacteria were resistant to cefepime (100%) and the highest resistance rate was seen to ceftazidime 92.6% followed by cefazolin 96.3%. The lowest resistance rate was observed to ciprofloxacin 38.9%, ofloxacin 44.4%, amikacin 46.3% and ticarcillin 48.1%. According to this study, *P. aeruginosa* isolates showed high level of antibiotic resistance and the presence of integrons in these strains can explain the influence of these genes in resistance creation. There was a significant association between resistance to cefotaxime, amikacin, ofloxacin, imipenem, ticarcillin, gentamicin and the presence of integrons.

Key words: *Pseudomonas aeruginosa*, integrons, drug resistance

* Corresponding author: Infectious Diseases and Tropical Medicine Research Center, Babol University of Medical Sciences, Babol, Iran.
E-mail: ramazan69@yahoo.com

P*seudomonas aeruginosa* is a non-fermenting aerobic Gram negative microorganism known as a pathogen leading to severe infections in hospitals. This organism is naturally resistant to many antimicrobial agents and it also can acquire the resistance against available antibiotics through multiple mechanisms (1). In the recent years due to the resistance of these pathogenic bacteria to different antibiotics, treatment of their infections has encountered a number of difficulties and many patients died because of these types of infections. The resistance mechanisms of these bacteria can be explained by either the mutation of genes or vertical transmissions through transformation or conjugation. Therefore the antibiotic resistance genes can be transported by plasmids and transposons between cells (2, 3).

Lately a new group of transportable genetic elements has been identified which can transfer the antibiotic resistance genes. This genetic element has been called integron and can surround genes and transfer them while they are in the gene cassettes. Integrons are classified into four classes according to their integrase genes, and the class 1 integron is the prevalent one (4, 5, 6).

Class 1 integron is composed of a 5'-conserved segment (5'CS) including the *intI* and *attI* genes and *Pant* promoter and 3'-conserved segment (3'CS) encoding resistance to sulfonamides (*SulI*) and disinfectants (7, 8). Class 2 integron is similar to class 1 and it contains integrase gene (*intI2*), one recombinant location (*aatI 2*) and a cassette gene without *SulI* in its 3' terminal region. The class 2 integron is less abundant than class 1 integron among Gram negative bacteria with antimicrobial resistance (9). Integron class 3 has *intI3* gene, *attI3* site and promoter and it is structurally identical to class 1 integron (10). Due to an important role of Integrons in antimicrobial resistance, a wide range of studies has been done on this gene. In Iran, an experiment has been performed in Imam Hospital of Uromieh that studied the influence of class 1

integron on the resistance of *P. aeruginosa* during 2007 to 2008 (11). Also the presence of class 1 and 2 integrons was studied in south of China during 2001 to 2005 (1).

In our previous study we investigated class 1 integron in *Pseudomonas aeruginosa* isolates from different places and devices of ICU in Shahid Beheshti Hospital during 2008 to 2009 (12). The aim of the present study was to perform a molecular analysis of all classes of integrons with hep primers in *Pseudomonas aeruginosa* strains isolated during 2011-2012 from the intensive care unit (ICU), and to determine their antibiotic resistance pattern as well as comparing these recent findings with those reported in our previous study.

Materials and Methods

Bacterial strains

This cross - sectional study was performed on 54 *P. aeruginosa* isolated from equipment and devices of ICU of Shahid - Beheshti hospital of Babol during 2011-2012. After collection, samples were cultured in EMB and blood agar media (Merck.Germany) and incubated at 37°C for 24 hours. The gram negative bacilli were analyzed using differential tests in order to identify the *P. aeruginosa* strains. These tests include: mobility, pigment production, growth at 42°C, oxidase and catalase tests.

Antimicrobial susceptibility testing

The susceptibility of the *P. aeruginosa* strains to cephalexine (CTX), amikacin (AN), ofloxacin (OFX), imipenem (IPM), cefepime (FEP), ticarcillin (TIC), gentamicin (GM), ceftazidime, (CAZ), ciprofloxacin (CP), cefazolin (CZ), ceftriaxone (CRO) and ceftizoxime (CT) antibiotics was determined by disc diffusion method (Padtan Teb Co, Iran) and the data were analyzed according to clinical and laboratory standards institute (CLSI 2006).

DNA extraction and polymerase chain reaction

DNA of isolates was extracted using High

Molecular detection of Integron genes and pattern of antibiotic

pure PCR Template preparation Kit (Roche Co, Germany). The PCR was performed in 50 μ l volumes that contained 10 μ l of extracted DNA, 50 pMole of each primer (Copenhagen, Denmark), 30 μ l distilled water, 5 μ l of 10X buffer, 1.5mMole MgCl₂, 0.2 mMole dNTPs and 1.5 U of Taq DNA polymerase enzyme (Copenhagen, Denmark). The presence of integrons was detected by hep primers in each isolate but this primer does not determine the class of integron. The hep 35 with the sequence of 5' TGCAGGTYAARGATBTKGATT 3' and hep 36 with the sequence of 5' CARCACATGCG-TRTARAT 3' were used for amplification of integrons gene (13).

Amplification of a 491bp fragment was conducted in a thermocycler as follows: a primary denaturation at 94°C for 5 minutes followed by 30 cycles of denaturation at 94°C for 30s, annealing at 55°C for 30s and extension at 72°C for 45s and one cycle of final extension at 72°C for 10 min. After performing PCR reaction, electrophoresis of PCR product was conducted in 1.5% agarose gel (Cinagene Co, Iran) and the results were evaluated in the presence of 100 bp DNA size marker (Fermentas Co, Ukraine).

Statistical analysis

The data were analyzed using the SPSS software version 18. The Chi-square test was

employed to calculate the P value in terms of resistant, intermediate and susceptible numbers of integron-positive and negative isolates.

Results

Fifty four isolated *P. aeruginosa* strains from Shahid Beheshti hospital ICU of Babol were tested for their resistance or susceptibility pattern to ceftizoxime, ceftazidime, cephalexine, amikacin, ofloxacin, imipenem, cefepime, ticarcillin, gentamicin, ciprofloxacin, cefazolin and ceftriaxone antibiotics by disc diffusion method. The highest resistances were against cefepime 100%, ceftazidime 92.6% and cefazolin 96.3% and the lowest were against ciprofloxacin 38.9%, ofloxacin 44.4%, amikacin 46.2% and ticarcillin 48.1% (Fig.1).

Furthermore, multi-drug resistant *P. aeruginosa* strains were studied. Twelve strains of 54 isolated *P. aeruginosa* were resistant to all antibiotics. Also 52 strains (96.3%) had shown resistance to 3 or more antibiotics and the second frequent group of this study was resistance to CT, CTX, CZ, CAZ, and FEP. In addition, the lowest frequency distribution of multi-drug resistance was seen in one isolate with resistance to two antibiotics Table 1; and only one isolate was susceptible to all antibiotics.




Fig 1. The antibiotic resistance and susceptibility of *P. aeruginosa* strains isolated from Shahid Beheshti hospital ICU of Babol to different antibiotics

Table 1: multi- drug resistance in *P. aeruginosa* strains isolated from Shahid Beheshti hospital ICU of Babol by disc diffusion method.

Antibiotics	Number of antibiotic resistant isolates	Antibiotics No.	Total number of resistant isolates (%)
FEP-CAZ-CZ-CT-CTX-CRO-AN-GM-TIC-IPM-OFX-CIP	12	12	12 (22.2%)
FEP-CAZ-CT-CTX-CRO-AN-CZ-IPM-TIC-CIP-OFX	1		
FEP-CAZ-CT-CTX-CRO-GM-TIC-CZ-AN-IPM-OFX	2		
FEP-CAZ-CT-CTX-CRO-CZ-OFX-GM-TIC-CIP-IPM	2	11	10 (18.5%)
FEP-CAZ-CZ-CT-CTX-IPM-GM-CIP-TIC-CRO-AN	3		
FEP-CZ-CAZ-CT-CTX-IPM-GM-CIP-CRO-AN-OFX	1		
FEP-CZ-CAZ-CTX-IPM-GM-CIP-CRO-AN-OFX-TIC	1		
FEP-CAZ-CTX-CRO-GM-TIC-IPM-CZ-CT-AN-	2	10	2 (3.7%)
FEP-CAZ-CT-CTX-CRO-TIC-GM-CZ-TIC-	1	9	1 (1.9%)
FEP-CZ-GM-CAZ-CRO-OFX-IPM	1		
FEP-CZ-AN-GM-CAZ-OFX-CT-	1	7	3 (5.5%)
FEP-CZ-CAZ-CRO-OFX-IPM-CT	1		
FEP-CZ-CAZ-CRO-CTX-CT	1		
FEP-CZ-AN-GM-CZ-CTX	1	6	2 (3.7%)
FEP-CAZ-CZ-GM-AN	1		
FEP-TIC-CRO-CTX-CAZ	1	5	9 (18.5%)
FEP-CZ-OFX-CTX-IPM	1		
FEP-CAZ-CZ-CTX-CT	6		
FEP-CZ-CAZ-CT	4		
FEP-CZ-CAZ-CTX	1	4	6 (11.2%)
FEP-CZ-AN-CT	1		
FEP-CAZ-TIC-	1		
FEP-CAZ-CZ	5	3	7 (12.9%)
FEP-CTX-CZ	1		
FEP-CZ	1	2	1 (1.9%)

PCR analysis, using hep35 and hep36 primers, amplified an expected band on agrose gel (491 bp) in 20 out of 54 isolates of *P. aeruginosa* and therefore 37% of the studied isolates had the integron genes (Fig. 2). Furthermore, all the integron positive strains were multi-drug resistant, and among them nine (45%) and four (20%) isolates were respectively resistant to twelve and eleven antibiotics which used in the current study. (Table 2). Table 2 shows antibiotic susceptibility of integron-positive and integron-negative strains of *P. aeruginosa*. 57% of isolates resistant to Ceftriaxone, 62% of isolates resistant to Ciprofloxacin, 59% of isolates resistant to Imipenem, 56% of isolates resistant to Amikacin

and 57% of isolates resistant to Gentamicin had integron genes (Table 2).


**Fig .2.** Gel-Electrophoresis of the PCR products of integrons by hep primers (1 & 2) and control samples (positive control: C+, negative control: C-).

Table 2: Antibiotic susceptibility of Integron-Positive and Integron-Negative strains of *P. aeruginosa* isolated from ICU.

Antibiotics	disc diffusion method				Integron Positive			Integron Negative			P value	
	n=54				n=20			n=34				
	R % (n)	I % (n)	S % (n)	R % (n)	I % (n)	S % (n)	R % (n)	I % (n)	S % (n)			
imipenem	53.7% (29)	1.9% (1)	44.4% (24)	85% (17)	0% (0)	15% (3)	35.3% (12)	2/9% (1)	61.8% (21)	0.01		
cefepime	100% (54)	0% (0)	0% (0)	100% (20)	0% (0)	0% (0)	100% (34)	0% (0)	0% (0)	NS		
ticarcillin	48.1% (26)	3.8% (2)	48.1% (26)	80% (16)	0% (0)	20% (4)	29.4% (10)	5/9% (2)	64.7% (22)	0.00		
ofloxacin	44.4% (24)	5.6% (3)	50% (27)	75% (15)	0% (0)	25 (5)	25% (9)	9/4% (3)	65.6% (22)	0.01		
amikacin	46.2% (25)	1.9% (1)!	51.9% (28)	70% (14)	5% (1)	25% (5)	32.4% (11)	0% (0)	67.6% (23)	0.04		
gentamicin	55.5% (30)	5.6% (3)	38.9% (21)	85% (17)	0% (0)	15% (3)	38.2% (13)	8.8% (3)	53% (18)	0.02		
cefazolin	96.3% (52)	0% (0)	3.7% (2)	100% (20)	0% (0)	0% (0)	94.1% (32)	0% (0)	5.9% (2)	NS		
ceftriaxone	51.8% (28)	24.1% (13)	24.1% (13)	80% (16)	10% (2)	10% (2)	35.2% (12)	32.4% (11)	32.4% (11)	0.07		
ceftizoxime	74.1% (40)	0% (0)	25.9% (14)	90% (18)	0% (0)	10% (2)	64.7% (22)	0% (0)	35.3% (12)	0.05		
cefotaxime	66.6% (36)	13% (7)	20.4% (11)	85% (17)	0% (0)	15% (3)	55.9% (19)	20.6% (7)	23.5% (8)	0.04		
ciprofloxacin	38.9% (21)	5.6% (3)	55.5% (30)	65% (13)	0% (0)	35% (7)	23.5% (8)	8.8% (3)	67.7% (23)	0.05		
Ceftazidime	92.6% (50)	3.7% (2)	3.7% (2)	100% (20)	0% (0)	0% (0)	88.2% (30)	5/9% (2)	5/9% (2)	NS		

R: resistant; I: intermediate; S: sensitive

Discussion

In this study the presence of integron genes in *P. aeruginosa* strains was investigated and antimicrobial susceptibility test was performed. A high proportion of isolated strains showed high resistance to investigated antibiotics. The presence of integron genes among resistant *P. aeruginosa* isolates confirms the role of this gene in resistance of strains and validates the origin of these bacteria from one common colony. Hence, there is a high risk of infection in hospitalized patients of ICU and due to increase of antibiotic resistance in isolated strains; their treatment is becoming a matter of concern. Therefore some strategies should be employed to inhibit the increasing resistance of these strains and the spread of resistance genes

between strains and prevent the colonization of these bacteria in hospital and especially in its ICU.

In our previous study we investigated class 1 integron in *Pseudomonas aeruginosa* isolates from ICU during 2008 to 2009 (12) and found that 39.4% of isolates had *intl* gene among which 24.2% were multidrug-resistant. While in this study 96.3% were multidrug-resistant. This difference could be due to the increased resistance or difference in methods used in susceptibility testing.

In this investigation the rate of susceptibility to ceftizoxime, ceftriaxone, ciprofloxacin, ofloxacin, was 25.9%, 24.1%, 55.5%, and 50%, respectively. These data are quite in line with Fuladi et al. results that reported 23% susceptibility to ceftizoxime and 29% to ceftriaxone but they

found 36% susceptibility to ciprofloxacin and 70% to ofloxacin (14). In addition, 96.3 % of isolated strains showed multi-drug resistance (MDR) which is higher than 87.05% of MDR reported in Mirsalehian et al.'s study (15).

Here, the resistance of isolated *P. aeruginosa* to ticarcillin, cefazolin, cephalexin, ceftriaxone, cefepime, amikacin, gentamicin, ciprofloxacin antibiotics were 48.1%, 96.3%, 66.6%, 51.8%, 100%, 46.2%, 55.5% respectively, while Chang and his colleagues in 2009 studied *klebsiella pneumoniae*, another cause of infection in hospital, and reported 9.2%, 21%, 3.7%, 5.8%, 0.3%, 9.2%, 17.4% and 14.8% rate of resistance to the same antibiotics (16). These data disclose the high rate of resistance and decreased rate of susceptibility of strains to different antibiotics in Iran which can be owing to hygienic statement, epidemiology of the region and different strains of bacteria in the studied region (14).

Also in a study that has been performed in United States during 2001 to 2003, most of multi-drug resistant strains (29.7%) were resistant to 4 antibiotics including ceftazidime, imipenem, gentamicin, ciprofloxacin or ofloxacin while in our investigation 22.2% of isolated strains were resistant to 12 antibiotics and 11.1% of specimens showed resistance to cefepime, ceftazidime, cefazolin, cephalexin, ceftizoxime. These data reveal the increasing rate of antibiotics resistance in Iran and all over the world (17).

Several studies in Iran have disclosed the critical condition of multi-drug resistance strains especially *P. aeruginosa* in hospital infections (18). Accordingly, studying the resistance pattern of these bacteria is an important subject. Increasing the resistance of *P. aeruginosa* to multiple antibiotics cause multiple drug regimens recruitment for those infections.

Several factors are involved in multi-drug resistance of *P. aeruginosa* among which change of membrane's permeability, plasmids and integrons

are important (19). The presence of integrons was detected by hep primers in each isolate, but this primer does not determine the class of integrons. The frequency of integrons in this study was 37% while Ruiz Martinez and his colleagues in Spain and Bing Gu et al. in China reported respectively 34.2% and 40.7% integrons frequency (20, 21).

In our investigation, 18 isolates (33.3%) were resistant to imipenem, ceftazidime, cefepime, ciprofloxacin and amikacin antibiotics among which, 14 (77.7%) had integrons; While a clinical study in Turkey on 67 isolates of *P. aeruginosa* from ICU showed that 79.1% were sensitive to imipenem, ceftazidime, cefepime, ciprofloxacin and amikacin antibiotics and only 14 (20.8%) were resistant to all putative antibiotics and in this group only two strains had integron (22). The different pattern of antibiotics resistance in Iran in comparison with other countries can be explained in part by its geographical diversity, the place of strains isolation and antibiotic usage.

In this research, the resistance of integron positive strains to gentamicin, ticarcillin, ciprofloxacin, imipenem, amikacin, ceftriaxone, ofloxacin, ceftazidime, cefazolin and cefepime antibiotics was 85%, 80%, 65%, 85%, 70%, 80%, 75%, 100%, 100% and 100%, respectively. These data showed that the presence of integron is significantly associated with resistance to gentamicin, ticarcillin, imipenem, amikacin, cefotaxime and ofloxacin antibiotics. Moreover, all of the integron positive strains were resistant to cefepime and ceftazidime but there was no significant association between the presence of integrons and the resistance to these two antibiotics.

This investigation can be compared with the study of Jianguo Chen and his colleagues in China where 38% of 71 isolated strains had integron and 88.9% of integron positive specimens were resistant to ceftriaxone, 92.6% to gentamicin, 48.1% to ceftazidime, 59.3% to ciprofloxacin, 25.9% to imipenem, 63% to cefepime, 66.7% to amikacin

Molecular detection of Integron genes and pattern of antibiotic

and 92.6% to cephotaxim (23). These results indicate that the presence of integrons can be effective in antibiotic resistance of *Pseudomonas aeruginosa* cases.

In our study, all 20 integron positive specimens showed multi-drug resistance and this may confirm the role of integrons in resistance of these bacteria to different antibiotics.

Since different groups of antibiotics have diverse effect on *Pseudomonas*, other mechanisms of resistance may be involved in this process (24, 25). In our study, we find only one isolate sensitive to all antibiotics. Decreasing the rate of sensitivity of these bacteria to multiple antibiotics can lead to many difficulties concerning the treatment of such infections.

Our data revealed that high rate of resistance of *Pseudomonas* to cefepime, ceftazidime, ticarcillin, imipenem, ofloxacin, cefazolin and ceftriaxone makes these antibiotics as an inappropriate drug for treatment of *Pseudomonas*' infections. It is possible that in the near future with the expansion of multdrug resistance, ciprofloxacin, ticarcillin and amikacin antibiotics lose their influence on *Pseudomonas*. In our investigation we observed some specimens that were resistant to different antibiotics inspite being integron negative. Therefore we should also look for other factors that make these bacteria resistant to antibiotics.

Acknowledgements

We thank the research council of Babol University of Medical Sciences for the financial support, department of Microbiology and Immunology Babol University of Medical Sciences and Department of Microbiology Science and Research Branch Islamic Azad University of guilan.

References

1. Xu Z, Li L, Shirtliff ME, et al. Occurrence and characteristics of class 1 and 2 integrons in *Pseudomonas aeruginosa* isolates from patients in southern China. *J Clin Microbiol* 2009;47:230-4.
2. Carattoli A. Importance of integrons in the diffusion of resistance. *Vet Res* 2001;32:243-59.
3. Stokes HW, Hall RM. A novel family of potentially mobile DNA elements encoding site-specific gene-integration functions: integrons. *Mol Microbiol* 1989;3:1669-83.
4. Partridge SR, Recchia GD, Stokes HW, et al. Family of class 1 integrons related to In4 from Tn1696. *Antimicrob Agents Chemother* 2001;45:3014-20.
5. Collis CM, Hall RM. Expression of antibiotic resistance genes in the integrated cassettes of integrons. *Antimicrob Agents Chemother* 1995;39:155-62.
6. Collis CM, Kim MJ, Stokes HW, et al. Binding of the purified integron DNA integrase IntI1 to integron- and cassette-associated recombination sites. *Mol Microbiol* 1998;29:477-90.
7. Peng CF, Lee MF, Fu HT, et al. Characterization of class 1 integrons and antimicrobial resistance in CTX-M-3-producing *Serratia marcescens* isolates from southern Taiwan. *Jpn J Infect Dis* 2007;60:250-6.
8. Penteado AP, Castanheira M, Pignatari AC, et al. Dissemination of bla(IMP-1)-carrying integron In86 among *Klebsiella pneumoniae* isolates harboring a new trimethoprim resistance gene dfr23. *Diagn Microbiol Infect Dis* 2009;63:87-91.
9. Marquez C, Labbate M, Ingold AJ, et al. Recovery of a functional class 2 integron from an *Escherichia coli* strain mediating a urinary tract infection. *Antimicrob Agents Chemother* 2008;52:4153-4.
10. Collis CM, Kim MJ, Partridge SR, et al. Characterization of the class 3 integron and the site-specific recombination system it determines. *J Bacteriol* 2002;184:3017-26.
11. Yousefi S, Nahaei M, Farajnia S, et al. Class 1 integron and Imipenem Resistance in Clinical Isolates of *Pseudomonas aeruginosa*: Prevalence and Antibiotic Susceptibility. *Iran J Microbiol* 2010;2:115-21.
12. Rajabnia R, Asgharpour F, Ferdosi Shahandashti E. Class 1 Integron in *Pseudomonas aeruginosa* Isolates From Different Places and Devices of ICU in Babol, Iran. *Jundishapur J Microbiol* 2013;6:138-43.
13. Farshad S, Japoni A, Hosseini M. Low distribution of integrons among multidrug resistant *E. coli* strains isolated from children with community-acquired urinary tract infections in

Shiraz, Iran. Pol J Microbiol 2008;57:193-8.

14. Imani Fouladi A, Satari M, Pourbabaei AA, et al. Relation Between Quinolones And Beta – Lactamase Resistance With Feature Of Producing Capsules In *Pseudomonas aeruginosa* Isolated From Urine. Medical Sciences Journal of Islamic Azad University 2009;19:97-103.
15. Mirsalehian A, Feizabadi M, Nakhjavani FA, et al. Detection of VEB-1, OXA-10 and PER-1 genotypes in extended-spectrum beta-lactamase-producing *Pseudomonas aeruginosa* strains isolated from burn patients. Burns 2010;36:70-4.
16. Chang CY, Fang YT, Tsai SM, et al. Characterization of class 1 integrons and gene cassettes in clinical isolates of *Klebsiella pneumoniae* from Taiwan. Diagn Micr Infec Dis 2009;65:214-6.
17. McGowan JE, Jr. Resistance in nonfermenting gram-negative bacteria: multidrug resistance to the maximum. Am J Med 2006;119:S29-36; discussion S62-70.
18. Fazeli H, Moslehi Tekantapeh Z, G.H.R. I, et al. Determination of Drug Resistance Patterns And Detection of bla-vim Gene in *Pseudomonas aeruginosa* Strains Isolated from Buren Patients In The Emam Mosa Kazem Hospital, Esfahan, Iran. Iranian Journal of Microbiology 2010;3:1-8.
19. Tenover FC. Mechanisms of antimicrobial resistance in bacteria. Am J Infect Control 2006;34:S3-S10.
20. Gu B, Tong MQ, Zhao WJ, et al. Prevalence and characterization of class I integrons among *Pseudomonas aeruginosa* and *Acinetobacter baumanii* isolates from patients in Nanjing, China. Journal of Clinical Microbiology 2007;45:241-3.
21. Ruiz-Martinez L, Lopez-Jimenez L, Fuste E, et al. Class 1 integrons in environmental and clinical isolates of *Pseudomonas aeruginosa*. Int J Antimicrob Agents 2011;38:398-402.
22. Budak F, Kasap M, Kolayli F, et al. Integron-associated resistance genes among multidrug-resistant *Pseudomonas aeruginosa* isolated from clinical specimens. Turk J Med Sci 2012;42:149-56.
23. Chen JG, Su ZL, Liu YZ, et al. Identification and characterization of class 1 integrons among *Pseudomonas aeruginosa* isolates from patients in Zhenjiang, China. Int J Infect Dis 2009;13:717-21.
24. Lister PD, Wolter DJ, Hanson ND. Antibacterial-resistant *Pseudomonas aeruginosa*: clinical impact and complex regulation of chromosomally encoded resistance mechanisms. Clin Microbiol Rev 2009;22:582-610.
25. Xavier DE, Picao RC, Girardello R, et al. Efflux pumps expression and its association with porin down-regulation and beta-lactamase production among *Pseudomonas aeruginosa* causing bloodstream infections in Brazil. BMC Microbiol 2010;10:217.