Summer 2018, Vol 7, No 3

DOI: 10.22088/IJMCM.BUMS.7.3.185

Inhibition of Interleukin-1 Receptor-Associated Kinases 1/4, Increases Gene Expression and Serum Level of Adiponectin in Mouse Model of Insulin Resistance

Athena Rajaie, Mostafa Allahyari, Mahdieh Nazari-Robati, Hossein Fallah*

Department of Clinical Biochemistry, Afzalipour School of Medicine, Kerman University of Medical Sciences, Kerman, Iran.

Submmited 17 August 2018; Accepted 8 October 2018; Published 14 October 2018

Insulin resistance is a feature of most patients with type 2 diabetes mellitus. Epidemiological evidence suggest a correlation between inflammation and insulin resistant states such as obesity, but the underlying mechanisms are largely unknown. Interleukin-1 receptor-associated kinases (IRAK) play a central role in inflammatory responses by regulating the expression of various inflammatory genes in immune cells. This study was aimed to investigate the effect of IRAK inhibitor on gene transcription and serum concentration of adiponectin in insulin-resistant mice. Experimental mice were randomly divided into 6 groups: the healthy control group was fed a regular chow diet while other groups were fed with a high-fat diet for 12 weeks. After this period, the animals were treated with IRAK inhibitor, pioglitazone, both IRAK and pioglitazone, and DMSO, for two weeks. Adiponectin gene expression level was analyzed by real-time PCR. Additionally, serum adiponectin levels were measured by ELISA. Homeostasis model assessment-adiponectin (HOMA-AD) as an insulin sensitivity index was calculated. IRAK inhibitor and pioglitazone increased significantly the expression of adiponectin gene. Also, adiponectin concentration in the control group (9.67±1.1 µg/ml) increased to 25.34±2.04 µg/ml in pioglitazone treatment group. IRAK inhibitor also increased adiponectin concentration (18.24±1.53 µg/ml) but did not show a synergistic effect with pioglitazone when administered simultaneously (26.66±2.5 μg/ml). HOMA-AD was 0.33±0.04 in pioglitazone treated group, 0.6±0.13 in IRAK inhibitor group, and 0.31±0.03 in animals that received IRAKi and pioglitazone. Our findings suggest that increased adiponectin secretion from adipose tissue mediated by IRAK inhibitor may increase the insulin sensitivity in an animal model of insulin resistance.

Key words: Insulin resistance, inflammation, adiponectin, IRAK inhibitor

Insulin resistance (IR) is a complicated condition in which three primary metabolic tissues that are sensitive to insulin; skeletal muscle, liver, and white adipose tissue (WAT) become less sensitive to insulin and its downstream metabolic actions

under normal serum glucose concentrations (1). IR is the condition in which a cell, tissue, or organism fails to respond appropriately to a given dose of insulin (2).

IR accompanies a wide range of pathological

^{*}Corresponding author: Department of Clinical Biochemistry, Afzalipour School of Medicine, Kerman University of Medical Sciences, Kerman, Iran. E-mail: hf59ma@gmail.com

conditions, including obesity, lipodystrophy, sepsis, steroid use, growth hormone excess, polycystic ovarian syndrome, cancer, neuro-degenerative disease, hypertension, hyperglycemia, and metabolic syndrome (1) and even some physiological conditions, such as pregnancy (2).

Obesity, characterized as a state of chronic low-grade inflammation caused by over- nutrition, is a major cause of decreased insulin sensitivity, which makes obesity a major risk factor for IR (1).

Factors released from adipose tissue that could contribute to the development of IR and B-cell dysfunction, include tumor necrosis factor α (TNF- α), free fatty acids (FFAs), adiponectin, resistin, leptin, agonists of the peroxisome proliferator-activated receptor γ (PPAR γ) (3), interleukin (IL)-1 β , IL-6, monocyte chemoattractant protein-1 (MCP-1), nuclear factor kappa B (NF κ -B), c-Jun N-terminal kinase (JNK), macrophage, high-sensitivity C-reactive protein (hs-CRP), the JAK-STAT signaling pathway (1).

Adiponectin, an adipocyte-specific secretory protein carrying 244 amino acids with 18 signal residues (4) is an adipokine that is specifically and abundantly expressed in adipose tissue, and directly sensitizes the body to insulin. The adiponectin receptors, AdipoR1 and AdipoR2, which mediate the antidiabetic metabolic actions of adiponectin, have been cloned, and are downregulated in obesity-linked IR (5).

Adiponectin, the adipocyte hormone with the highest plasma concentration, is considered a modulator of carbohydrate and lipid metabolism and a marker of insulin sensitivity. Although mainly produced in adipose tissue, serum adiponectin concentrations are negatively correlated with the amount of visceral adiposity (6).

Numerous clinical studies demonstrated an inverse relationship between serum adiponectin levels and overproduction of pro-inflammatory markers such as TNF- α and CRP (7). Given that IL-6 is pro-inflammatory, it is widely accepted that

like TNF- α , IL-6 negatively impacts obesity-induced IR (8).

Thiazolidinediones, including pioglitazone, constitute a new class of oral antidiabetic drugs that are widely used as insulin-sensitizing agents through the activation of PPAR-γ, thereby regulating the transcription of specific genes involved in adipogenesis and IR (9).

Inflammation plays an important role in the development of IR via various cytokines and molecular pathways, and may therefore be targeted with appropriate interventions to prevent IR (1). Interleukin 1 receptor-associated kinase 1(IRAK1) mediates pro-inflammatory signaling receptor/toll-like receptors, via IL-1 which may contribute to IR. IL-1-R and toll-like receptors interact with MyD88 to activate IRAK-4 which phosphorylates and activates IRAK-1. Downstream IRAK-1, **TNF** from receptor associated factor 6 (TRAF6) interacts with Mitogen-activated protein kinase kinase kinase 7 (MAP3K7) to activate the IkB kinase (IKK) complex leading to phosphorylation of IkBa that then dissociates from NF-κB. NF-κB enters the nucleus to promote the transcription of many proinflammatory genes including interferons and cytokines, such as IL-1 and TNF-α (10). IRAK-1 phosphorylate insulin receptor substrate 1 (IRS-1) at Ser24, which impairs binding of IRS-1 to phosphatidylinositol 3-kinase (PI3K), thereby impairing metabolic actions of insulin including insulin-stimulated translocation glucose transporter type 4 (GLUT4) (1).

IRAK-1, which impairs insulin signaling, thereby induces IR. Understanding the detailed mechanisms of this cross-talk between proinflammatory signaling pathways and insulin signaling pathways may help to improve the diagnosis of pre-diabetic states, and also suggest new strategies to treat diabetes (10). Potent and selective small molecule inhibitors targeting IRAK-4 kinase activity are currently being pursued as

potential therapeutics for autoimmune and inflammatory diseases (11).

IRAK inhibitor can suppress the inflammation pathways and may therefore increase adiponectin expression and secretion by decreasing the level of pro-inflammatory cytokines. Thereupon, we aimed to evaluate the adiponectin gene expression level under treatment with IRAK inhibitor.

Materials and methods

Animals

In this experimental study, male C57BL/6 mice, (mean weight 19 g, 6 weeks-of-age) were obtained from the animal house of Pasteur Institute of Iran (IPI). The Ethics Committee of Kerman University of Medical Sciences approved the study procedure. The animals were maintained in a temperature-controlled room (22°C) on a 12-h light-dark cycle. One week after arrival, mice were randomly divided into 6 groups (n= 8 each) and 5 groups were fed high-fat diet while one group received continuous feeding of a normal diet for 12 weeks. High-fat diet was composed of 365 g regular chow, 250 g casein, 310 g fat, 60 g minerals and vitamins, 60 g cholesterol, 3 g methionine, and 2 g cholic acid per Kg (12).

Treatment

After 12 weeks, the animals were treated for two additional weeks as follows: healthy control group received normal diet without any drug; control group received high-fat diet without any drug; IRAK inhibitor (IRAKi) group received high-fat diet with intra peritoneal (i.p.) injection of 2.12 mg/kg IRAKi 3 times weekly (13). IRAKi was diluted in DMSO (5 mM) and further dissolved in sterile PBS (pH 7.2). Pioglitazone (Pio) group received high-fat diet and intra gastric infection of pioglitazone (10 mg/kg/day) (9). IRAKi+Pio group received high-fat diet with pioglitazone and IRAKi as mentioned above. The vehicle group was fed with high-fat diet and DMSO.

Blood and tissue collection

At the end of the treatment period (2 weeks), animals were fasted for 12 h overnight. A separate group of animals was killed by cardiac puncture under terminal anesthesia induced by diethyl ether (14). Visceral adipose tissue samples were harvested. All samples were flash frozen in liquid nitrogen and subsequently stored at -75 °C until evaluation.

Measurement of serum adiponectin

Serum samples were separated from blood, by centrifuge at 3000 rpm for 10 min at 4 °C. Adiponectin level was determined with a double-antibody radioimmunoassay (ZellBio GmbH kit, Germany). Homeostasis model assessment of adiponectin (HOMA-ID) was calculated using the equation: [(insulin (U/ml) × glucose (mmol/l))/22.5×adiponectin (µg/ml)] (15).

Total RNA extraction

Total RNA was extracted from visceral adipose tissue samples using a RNeasy Mini Kit (Favorgen Biotech Corp, Taiwan) incorporating DNase digestion. Extracted total RNA was quantified using a NanoDrop spectrophotometer (ND-1000 nanodrop). RNA quality was assessed by ethidium bromide staining of 18S and 28S ribosomal RNA after electrophoresis on 1.5% agarose gel. Complementary cDNA templates for real-time PCR assays were prepared from Moloney Murine Leukemia Virus Reverse Transcriptase (M-MLV RT), oligodT primer (Favorgen Biotech Corp, Taiwan) and total RNA (2.5 μg).

Real Time PCR

Real-time quantitative PCR amplification was conducted with the mic Biomolecular systems (BMS, Australia) using YTA Super SYBR Green qPCR MasterMix 2X (Favorgen Biotech Corp, Taiwan) according to the protocol recommended by the manufacturer. PCR was run as follows: initial activation of Taq DNA polymerase at 94 °C for 3 min, 5 s at 95 °C for denaturing, 5 s at 55 °C for annealing, 10 s at 72 °C for elongation, and 40 PCR cycles were performed. All experiments were run in

duplicate and non-template controls and dissociation curves were used to detect primer—dimer conformation and nonspecific amplification. According to the similarity of PCR efficiency for adiponectin and *GAPDH*, difference in gene expression was calculated by 2-^{ΔΔCT} method.

Primer sequences are described in Table 1 (Pishgam Biotech Co, Iran).

Statistical analysis:

All data were analyzed using SPSS software, Version 22. For determination of difference between groups, nonparametric Kruskal-Wallis test was used. All data were presented as Mean \pm SE. P< 0.05 was considered statistically significant. For determination of correlation between HOMA-IR and adiponectin, Spearman's Rho test was used.

Table 1. Sequence of primers		
Gene	Primers 5' → 3'	Amplicon size (bp)
Adiponectin	F:ATGGCAGAGATGGCACTCCT	162
	R:CATAAGCGGCTTCTCCAGGC	
GAPDH	F:ACCATCTTCCAGGAGCGAGA	
	R:GAAGGGGCGGAGATGATGAC	152




Fig. 1. Transcription level of adiponectin gene. Each IRAK inhibitor or pioglitazone increased the expression level of adiponectin gene in high-fat diet mice. IRAKi: IRAK inhibitor; pio: pioglitazone. *** P<0.0001, ** P<0.01, * P<0.05 in comparison with control group.

Results

Transcription level of adiponectin gene

RT-qPCR results showed that IRAK inhibitor significantly increased the expression of adiponectin gene (P<0.0001). Also, pioglitazone increased adiponectin gene expression (P=0.0027), although these agents did not show any incremental effect on each other. Results of adiponectin gene expression evaluation are summarized in figure 1.

Adiponectin serum concentration

High-fat diet resulted in a decrease of serum adiponectin concentration (9.67±1.1 µg/ml) in comparison with healthy control group (24.02±1.24 ug/ml) that received normal diet. However, pioglitazone treatment increased adiponectin concentration to 25.34±2.04 µg/ml. IRAK inhibitor adiponectin concentration also increased (18.24±1.53 μg/ml) but could not exert a synergistic effect on pioglitazone when administred simultaneously (26.66±2.5 μg/ml) (figure 2).




Fig. 2. Serum concentration of adiponectin. IRAK inhibitor or pioglitazone increased the serum concentration level of adiponectin in high- fat diet mice. IRAK: IRAK inhibitor; pio: pioglitazone.*** P< 0.0001, ** P< 0.01, ** P< 0.05 in comparison with control group.




Fig. 3. Effects of IRAKi on HOMA-AD. IRAK inhibitor or pioglitazone decreased HOMA-AD in high-fat diet mice. IRAKi: IRAK inhibitor; pio: pioglitazone. *** P< 0.0001, ** P< 0.01, ** P< 0.05 in comparison with control group.




Fig. 4. Correlation between HOMA-IR and adiponectin. Correlation coefficient= -0.527; P< 0.0001.

HOMA-AD: Index of insulin sensitivity

As shown in figure 3, HOMA-AD was 0.33±0.04 in pioglitazone treated group, 0.6±0.13 in IRAK inhibitor group, and 0.31±0.03 in animals that received IRAKi and pioglitazone. These results were significantly lower than the control group (2.27±0.28, P<0.0001). HOMA-AD in healthy animals was 0.10±0.007 that is indicative of high insulin sensitivity of these animals.

HOMA-IR and adiponectin correlation

There was a significant (P<0.0001) and negative correlation (correlation coefficient=-0.527) between adiponectin concentration and HOMA-IR (figure 4) shows this correlation.

Discussion

In obesity state, adipose tissue is developed and its macrophages are increased. Low-grade inflammation increases blood free fatty acid levels which results from obesity, and can induce IR and type 2 diabetes (16, 17). Obesity also is related to a decrease of adiponectin concentration. Adiponectin is an insulin sensitizer adipokine (18). IRAK inhibitor, as an interference in inflammation pathways, can increase insulin sensitivity. Therefore, we evaluated the effects of this factor in comparison with pioglitazone on adiponectin gene expression at mRNA and protein levels.

Our results showed that IRAK inhibitor and pioglitazone increased the transcription level of adiponectin. Also, adiponectin concentration increased in serum of insulin-resistant mice by IRAK inhibitor and pioglitazone.

In Toll-Like Receptor signaling pathways which are activated by lipids, lipoproteins, lipopolysaccharides, and proteins, MYD88 is complexed with IRAK4 and IRAK1. IRAK4 activates IRAK1. Activated IRAK1 is released from the complex and through a specific signaling cascade, finally activates NF-κB, a key factor in inflammation response, which increases the transcription of some pro-inflammatory cytokines

(17, 19). IRAK inhibitor by suppression of this signaling pathway can induce anti-inflammatory functions (11) and decrease IR which results from inflammation.

There is a negative correlation between BMI and adiponectin concentration so obesity along with its related inflammation decreases adiponectin production (20, 21). Adiponectin production is controlled by several factors. Pro-inflammatory cytokines that increased in IR, control negatively the production of adiponectin. In contrast, PPARγ increases the expression of adiponectin in adipose tissue (22). This is compatible with our results that pioglitazone as a PPARγ agonist and IRAK inhibitor as an anti-inflammatory factor increased the level of adiponectin.

Overexpression of IRAK can induce inflammation and decrease adiponectin level. So, it is not surprising that IRAK inhibitor increased adiponectin concentration (23). However, deletion of the IRAK1 gene can increase insulin sensitivity without any effect on adiponectin level (10).

HOMA-AD is an insulin sensitivity index, and our results showed that IRAK inhibitor significantly increased the insulin sensitivity with an efficiency completely comparable with pioglitazone. Also, we showed a significant and negative correlation between HOMA-IR and adiponectin levels, suggesting that adiponectin plays a critical role in tolerance of glucose and increase of sensitivity to insulin. Furthermore, adiponectin can act as a predictor IR, so that increase in adiponectin level, decreases IR, and increase in IR is accompanied by adiponectin decrease.

In our study, pioglitazone and IRAK inhibitor did not show any synergistic or antagonist effect on each other, suggesting that each of them exert their effects in an independent manner. Pioglitazone as a PPAR γ agonist increases the expression of adiponectin while IRAK inhibitor increases the adiponectin level by suppression of proinflammatory cytokines (22, 24, 25).

Adiponectin through increasing of fatty acid oxidation and glucose uptake, increases insulin sensitivity (26). In addition, adiponectin has anti-inflammatory activity and decreases NF.κB dependent inflammation (18).

In conclusion, IRAK inhibitor increases adiponectin production and secretion from adipocytes. Adiponectin is an insulin sensitizer adipokine and for this reason, it is possible that IR decrease by IRAKi is mediated by adiponectin gene overexpression. IRAK inhibitor efficacy in reducing HOMA-AD is fully comparable with pioglitazone and further studies may lead to its introduction as a new antidiabetic agent.

Conflict of interest

Authors declare no conflict of interest.

References

- Chen L, Chen R, Wang H, et al. Mechanisms Linking Inflammation to Insulin Resistance. Int J Endocrinol 2015;2015;508409.
- Kang S, Tsai LT, Rosen ED. Nuclear Mechanisms of Insulin Resistance. Trends Cell Biol 2016;26:341-51.
- Arner P. The adipocyte in insulin resistance: key molecules and the impact of the thiazolidinediones. Trends Endocrinol Metab 2003;14:137-45.
- 4. Hayama S, Higuchi T, Miyakoshi H, et al. Analytical evaluation of a high-molecular-weight (HMW) adiponectin chemiluminescent enzyme immunoassay. Clin Chim Acta 2010:411:2073-8.
- Kadowaki T, Yamauchi T, Kubota N, et al. Adiponectin and adiponectin receptors in insulin resistance, diabetes, and the metabolic syndrome. J Clin Invest 2006;116:1784-92.
- 6. Sirbu AE, Buburuzan L, Kevorkian S, et al. Adiponectin expression in visceral adiposity is an important determinant of insulin resistance in morbid obesity. Endokrynol Pol 2018:69:252-8.
- Shetty S, Kusminski CM, Scherer PE. Adiponectin in health and disease: evaluation of adiponectin-targeted drug development strategies. Trends Pharmacol Sci 2009;30:234-9.
- 8. Kraakman MJ, Kammoun HL, Allen TL, et al. Blocking IL-6 trans-signaling prevents high-fat diet-induced adipose tissue

- macrophage recruitment but does not improve insulin resistance.

 Cell Metab 2015;21:403-16.
- 9. Elrashidy RA, Asker ME, Mohamed HE. Pioglitazone attenuates cardiac fibrosis and hypertrophy in a rat model of diabetic nephropathy. J Cardiovasc Pharmacol Ther 2012;17:324-33.
- 10. Sun XJ, Kim SP, Zhang D, et al. Deletion of interleukin 1 receptor-associated kinase 1 (Irak1) improves glucose tolerance primarily by increasing insulin sensitivity in skeletal muscle. J Biol Chem 2017;292:12339-50.
- 11. Wang Z, Wesche H, Stevens T, et al. IRAK-4 inhibitors for inflammation. Curr Top Med Chem 2009;9:724-37.
- 12. Wang CY, Liao JK. A mouse model of diet-induced obesity and insulin resistance. Methods Mol Biol 2012;821:421-33.
- 13. Rhyasen GW, Bolanos L, Fang J, et al. Targeting IRAK1 as a therapeutic approach for myelodysplastic syndrome. Cancer Cell 2013;24:90-104.
- 14. Williams LM, Campbell FM, Drew JE, et al. The development of diet-induced obesity and glucose intolerance in C57BL/6 mice on a high-fat diet consists of distinct phases. PLoS One 2014;9:e106159.
- 15. Vilela BS, Vasques AC, Cassani RS, et al. The HOMA-Adiponectin (HOMA-AD) Closely Mirrors the HOMA-IR Index in the Screening of Insulin Resistance in the Brazilian Metabolic Syndrome Study (BRAMS). PLoS One 2016;11:e0158751.
- Meshkani R, Adeli K. Hepatic insulin resistance, metabolic syndrome and cardiovascular disease. Clin Biochem 2009;42:1331-46.
- 17. Lauterbach MA, Wunderlich FT. Macrophage function in obesity-induced inflammation and insulin resistance. Pflugers Arch 2017;469:385-96.
- 18. Wang X, Chen Q, Pu H, et al. Adiponectin improves NF-kappaB-mediated inflammation and abates atherosclerosis progression in apolipoprotein E-deficient mice. Lipids Health Dis 2016;15:33.
- Kawasaki T, Kawai T. Toll-like receptor signaling pathways.
 Front Immunol 2014;5:461.
- 20. Frankenberg ADV, Reis AF, Gerchman F. Relationships between adiponectin levels, the metabolic syndrome, and type 2 diabetes: a literature review. Arch Endocrinol Metab 2017;61:614-22.
- 21. Zembala-Szczerba M, Jaworowski A, Huras H, et al. Low-

Grade Metabolically- Induced Inflammation Mediators Interleukin-6, Adiponectin, and TNF-alpha Serum Levels in Obese Pregnant Patients in the Perinatal Period. Med Sci Monit Basic Res 2017;23:1-7.

- 22. Shehzad A, Iqbal W, Shehzad O, et al. Adiponectin: regulation of its production and its role in human diseases. Hormones (Athens) 2012;11:8-20.
- 23. Ahmad R, Shihab PK, Thomas R, et al. Increased expression of the interleukin-1 receptor-associated kinase (IRAK)-1 is associated with adipose tissue inflammatory state in obesity. Diabetol Metab Syndr 2015;7:71.
- 24. Yan H, Xie H, Ying Y, et al. Pioglitazone use in patients with diabetes and risk of bladder cancer: a systematic review and meta-analysis. Cancer Manag Res 2018;10:1627-38.
- 25. Simons PJ, van den Pangaart PS, Aerts JM, et al. Proinflammatory delipidizing cytokines reduce adiponectin secretion from human adipocytes without affecting adiponectin oligomerization. J Endocrinol 2007;192:289-99.
- 26. Achari AE, Jain SK. Adiponectin, a Therapeutic Target for Obesity, Diabetes, and Endothelial Dysfunction. Int J Mol Sci 2017;18.